


# **Gauge Approach and Quantization Methods in Gravity Theory**

Vladimir N. Ponomarev  
Andrei O. Barvinsky  
Yuri N. Obukhov



NAUKA



IBRAE

Nuclear Safety Institute of the Russian Academy of Sciences

Vladimir N. Ponomarev

Andrei O. Barvinsky

Yuri N. Obukhov

# **Gauge Approach and Quantization Methods in Gravity Theory**



MOSCOW NAUKA 2017

УДК 530.12

ББК 22.31

П56

**Рецензенты:**

доктор физико-математических наук, профессор В.Ч. Жуковский,  
доктор физико-математических наук, профессор П.С. Кондратенко

**Gauge approach and quantization methods in gravity theory** / Ponomarev V. N., Barvinsky A. O., Obukhov Yu. N. ; Nuclear Safety Institute of the Russian Academy of Sciences — М. : Nauka, 2017. — 360 p. : ill. — ISBN 978-5-02-040047-4 (bound).

The book gives an overview of the geometrical gauge approach to the gravity theory and the methods of quantization of the gravitational field. Gauge-theoretic formalism (universal principle of the local invariance and the mechanism of spontaneous breaking of the gauge symmetry) forms the basis for the modern understanding of fundamental physical interactions and is successfully confirmed by the experimental discoveries of the gauge bosons and the Higgs particle. The carefully selected material of the book provides a minimal but sufficient mathematical introduction to the methods of the gauge gravitational theory, and gives a concise but exhaustive description of all specific physical consequences. In order to describe the many dramatic changes which took place recently in understanding the concepts, issues and motivations of quantum gravitational physics, a special emphasis is put on a comparison of three different quantization methods of the gravitational field – the covariant approach, the Dirac-Wheeler-DeWitt quantization and the method of Arnowitt-Deser-Misner. The basics of the canonical quantization is explained along with the detailed exposition of the path integral approach, and the introduction to the modern BRST technique is given to demonstrate the consistency of the Arnowitt-Deser-Misner and the Dirac-Wheeler-DeWitt quantization schemes.

The book addresses the physicists who specialize in the gravity theory and the high-energy physics, and it can be recommended to the graduate students and senior undergraduate university students in theoretical physics and mathematics.

**Калибровочный подход и методы квантования в теории гравитации** / Пономарев В. Н., Барвинский А. О., Обухов Ю. Н. ; Институт проблем безопасного развития атомной энергетики РАН — М. : Наука, 2017. — 360 с. : ил. — ISBN 978-5-02-040047-4 (в пер.).

Формализм калибровочной теории (универсальный принцип локальной инвариантности и механизм спонтанного нарушения калибровочной симметрии) лежит в основе современного понимания фундаментальных физических взаимодействий и успешно подтверждается экспериментальными открытиями калибровочных бозонов и частицы Хиггса. В книге дается обзор геометрического калибровочного подхода к теории гравитации и методов квантования гравитационного поля. Тщательно подобранный материал книги обеспечивает минимальное, но достаточно математическое введение в методы калибровочной теории гравитации и дает краткое, но исчерпывающее описание конкретных физических следствий калибровочного подхода. Чтобы осветить многочисленные драматические изменения, произошедшие в последнее время в осмыслении понятий, проблем и целей квантово-гравитационной физики, особое внимание уделяется сравнению трех различных методов квантования гравитационного поля – ковариантного подхода, квантования Дирака-Уилера-ДеВитта и метода Арновитта-Дезера-Мизнера. В книге объясняются основы канонического квантования наряду с подробным изложением формализма континуальных интегралов, и дается введение в современную технику БРСТ, с тем чтобы продемонстрировать согласованность схем квантования Арновитта-Дезера-Мизнера и Дирака-Уилера-ДеВитта.

Книга адресована физикам, которые специализируются в теории гравитации и физике высоких энергий, и ее можно рекомендовать аспирантам и студентам старших курсов по теоретической физике и математике.

ISBN 978-5-02-040047-4

© Ponomarev V. N., Barvinsky A. O., Obukhov Yu. N., 2017

© Nuclear Safety Institute of the Russian Academy of Sciences, 2017

© Design and Layout. Publishing House «Nauka», 2017

# Contents

<b>Preface</b>	<b>viii</b>
<b>1 Lagrangian description of gravity theories of Hilbert-Einstein type</b>	<b>1</b>
1.1. Geometrical spacetime structures.....	1
1.2. Hilbert variational principle and field equations of the general relativity .....	8
1.3. First-order formalism and field equations of the Einstein-Cartan theory .....	11
<b>2 Canonical formalism of gravity theories</b>	<b>17</b>
2.1. Systems with singular Lagrangians.....	17
2.2. Peculiarities of the canonical formalism in curved spacetime. ....	30
2.3. Geometry of $(3 + 1)$ -decomposition of spacetime.....	31
2.4. Canonical formalism for the fields on the curved spacetime.....	34
<b>3 Dynamics of gravity theories of Hilbert-Einstein type</b>	<b>45</b>
3.1. The problem of the “frozen” formalism.....	45
3.2. Arnowitt-Deser-Misner procedure of selection of physical degrees of freedom .....	46
3.3. Asymptotically flat and closed worlds.....	49
<b>4 Torsion effects on the structure and evolution of gravitating systems</b>	<b>57</b>
4.1. Matter fields in the Einstein-Cartan theory.....	57
4.2. Conformal invariance and spacetime torsion .....	71
4.3. Pre-Friedman stage of Universe’s evolution and spacetime torsion.....	74
4.4. Production of scalar particles by cosmological torsion field .....	80

<b>5</b>	<b>Kinematics of gauge theories of gravity</b>	<b>87</b>
5.1.	Special aspects of gauge approach in gravitation .....	87
5.2.	Bundle of frames and generalized affine connection .....	89
5.3.	Spontaneous symmetry breaking and non-linear realizations .....	97
5.4.	Kinematics of the gauge gravity theory.....	100
<b>6</b>	<b>Gauge gravity models with dynamical (<math>\Gamma</math>-S)-interaction</b>	<b>107</b>
6.1.	Principles for construction of dynamics .....	107
6.2.	Choice of the action in the gauge gravity theory .....	109
6.3.	Physical consequences of the gauge gravity theory .....	114
6.4.	Model description of microscopic gravitational interactions.....	121
<b>7</b>	<b>Quantization of gravity</b>	<b>129</b>
7.1.	Methods of quantum gravity .....	129
7.2.	Arnowitt-Deser-Misner quantization method.....	133
7.3.	Dirac-Wheeler-DeWitt quantum geometrodynamics .....	138
7.4.	Dirac quantization from Batalin-Fradkin-Vilkovisky formalism .....	142
7.5.	Semiclassical approximation.....	152
7.6.	Problems and prospects of quantum gravity and cosmology .....	160
	<b>Appendices</b>	<b>163</b>
A1.	Geometry of manifolds.....	163
A2.	Spinor analysis on an arbitrary manifold.....	169
A3.	Basic notations.....	173
A4.	Comments on the literature and general remarks .....	174
	<b>References</b>	<b>179</b>
	<b>Subject index</b>	<b>197</b>
	<b>Bibliography on gauge gravity theory</b>	<b>201</b>



## VLADIMIR N. PONOMAREV

Vladimir Ponomarev was born in Moscow in 1945. He has studied physics at the Moscow State Lomonosov University and at the Warsaw University. His first scientific adviser was Andrzej Trautman, who supervised his internship at the Institute of Theoretical Physics of the Warsaw University in 1969-72. After defending his PhD thesis (scientific advisors: Andrzej Trautman and Dmitry Ivanenko) in 1974, he worked at the Department of Theoretical Physics at Moscow State Lomonosov University till 1982, when he was elected as the head of the Department of Physics for Natural Sciences of the Moscow State Pedagogical University. In 1985, he was awarded the title of full professor. In 1988, he took part in the organization of the Nuclear Safety Institute, where he became the deputy director. He published about 150 scientific papers on the classical and quantum theory of gravity, as well as coauthored the monograph “Geometrodynamical methods and the gauge approach in the theory of gravitational interactions” (1985, Moscow) together with A.O. Barvinsky and Yu.N. Obukhov. During the period of 1991-2009, he was involved in the work on social-economic transformations within the former USSR, including the service in the Government of the Russian Federation as a Deputy Minister. In 2009, he resumed the work in the Nuclear Safety Institute as the Deputy Director for Strategic Development and Innovations, along with continuing the activities in expert groups under the Government and Parliament of the Russian Federation. Throughout his 50-year professional career, he has been active in teaching, also in the recent times as an invited professor at several universities and academic institutions, including the Moscow Physical-Technical University.



## ANDREI O. BARVINSKY

Andrei Barvinsky was born in 1955. After graduating from the Physics Department of Moscow State University he has got his PhD in theoretical physics in the field of quantum gravity theory and held a postdoctoral fellowship at the University of Alberta, Canada. Since then he is a leading scientific researcher at the Theory Department of the Lebedev Physics Institute of the Russian Academy of Sciences. The areas of his scientific interests are quantum field theory, quantum gravity and cosmology. In these areas he has more than one hundred publications including the pioneering contributions to the quantum theory of gauge constrained systems, background formalism and the effective action method in quantum field theory and Higgs inflation theory in cosmology of the very early quantum Universe. His teaching activity included supervision of many PhD students and professorship at the Ludwig-Maximillan University in Munich, Germany. He has an ongoing wide scientific collaboration with the physicists of the Landau Institute for Theoretical Physics, the Physics Department of the Ludwig-Maximillan University in Munich, University of Cologne, Department of Physics and Astronomy of the University of British Columbia, the University of Bologna and the Theory Division of CERN. He is also one of the organizers of the long series of the Sakharov and Ginzburg conferences on physics held for more than twenty years at the Theory Department of the Lebedev Physics Institute in Moscow.



## YURI N. OBUKHOV

Yuri Obukhov was born in Berlin in 1956 and graduated from the Department of Physics of the Moscow State Lomonosov University. He received the PhD doctoral degree in 1983 from the Moscow State University, where he later worked for many years. He spent the first post-doctoral term in 1986-87 in the relativity group of A. Trautman at the Institute of Theoretical Physics of the Warsaw University. After he was awarded a Humboldt Fellowship and visited the University of Cologne in 1992-94, a long and fruitful collaboration with Friedrich Hehl has started, and subsequently Yuri Obukhov has held several times research positions at the Institute for Theoretical Physics of the University of Cologne. He was also twice (2002 and 2006) a visiting professor at the Institute for Theoretical Physics of the University of Sao Paulo (UNESP), and a research fellow at the Mathematics Department of the University College London (2009-2011). Since 2013 he is a senior scientist at the Nuclear Safety Institute of the Russian Academy of Sciences. Along with research, Yuri Obukhov took part in educational activities, teaching at different universities the lecture courses on classical and quantum field theory, classical gravity theory, relativistic cosmology, and supervising numerous master and PhD students. The scientific interests of Yuri Obukhov encompass the relativistic quantum theory, electrodynamics, classical gravity theory and cosmology. His original results obtained in these areas were published in more than 150 papers. In 2003, together with Friedrich Hehl he published a book “Foundations of Classical Electrodynamics: Charge, Flux, and Metric” (Birkhauser, Boston).

# Preface

Modern understanding of gravitational phenomena is based on the concept of the spacetime geometry. This is drastically different from Newton's theory in which the Euclidean space and an absolute time represented a fixed arena for independent physical processes of different kind: mechanical, electric, magnetic, gravitational. Eventually, the development of ideas of Minkowski and Einstein culminated in a unification of space and time into a four-dimensional spacetime manifold. Its geometry became dynamical: it is no longer assumed to be fixed but depends on the motion of matter. In a broad sense, gravitation can be viewed as a geometrodynamics, i.e., as a theory of a dynamical geometry of spacetime. The beautiful and powerful methods of differential geometry are used in Einstein's general relativity (GR) to describe the gravitational phenomena in terms of the properties of the four-dimensional spacetime manifold. Later, it was recognized that the three other physical interactions (electromagnetic, weak and strong) can also be geometrized by making use of the Yang-Mills gauge-theoretic approach, and a natural question arose whether the gravity theory can be consistently formulated in the gauge framework. The answer is not that simple as it might appear at the first sight. The subtle point is that the Standard Model is based on the fundamental symmetry groups acting in the internal spaces, whereas the gravity is obviously related with the symmetry of the spacetime itself.

In the recent decades, a growing interest of researchers has been attracted to the problem of construction of a unified theory of fundamental interactions. Such a theory, capable to explain the hierarchical structure of physical forces in nature, most probably could be based on the universal principle of the local invariance and on the resulting formalism of the gauge fields. A significant progress in this area is manifest in the development of the unified models of weak and electromagnetic interactions so successfully confirmed by the experimental discoveries of the gauge bosons and the Higgs particle. However, these models do not take the gravity into account, and therefore they should be considered only as approximations to the unification of all interactions which could take place on extremely small scales, or at the very high energies and high temperatures of the early Universe, when the gravitational effects are expected to be significant. Thus, the next step in the construction of a genuine unified theory is a consistent

inclusion of gravity in the general scheme of fundamental physical interactions, reconciling the principles of general relativity theory with those of the quantum theory and of gauge field theory.

This book gives an overview of the geometrical gauge approach to the gravity theory (Chapters 1, 4-6) and methods of the gravitational field quantization (Chapters 2, 3 and 7). Taking into account the long history of the subject and the volume of the relevant literature, the material for the book was carefully selected to provide a minimal but sufficient mathematical introduction to the methods of the gauge gravitational theory (Chapter 4), and to give a concise but exhaustive description of all specific physical consequences (Chapters 3 and 5). The special feature of our book is the use of the invariant language to describe the  $(3 + 1)$ -decomposition of the spacetime manifold (Chapter 2). A particular emphasis is put on a comparison of three different quantization methods of the gravitational field – the covariant approach, the Dirac-Wheeler-DeWitt quantization and the method of Arnowitt-Deser-Misner (Chapter 3).

In this regard, it is worth emphasizing that many dramatic changes took place recently in understanding the concepts, issues and motivations of quantum gravitational physics. Precision cosmology in the form of the cosmic microwave background (CMB) observations actually rendered quantum gravity the status of an experimental subject, inflation theory acquired a dominant role in the picture of the early Universe and we have been put face to face with the mysteries of the dark matter and dark energy phenomena. On top of this progress quantum geometrodynamics method and quantum cosmology have lost a status of acute direction in theoretical physics and, moreover, a widely accepted viewpoint prevailed that this approach is essentially discredited due to numerous but rather unproductive applications of the Wheeler-DeWitt equation. This opinion is, however, very erroneous, in particular, because the analogous situation in quantum field theory is treated completely differently – nobody calls in question the status of the Schrödinger equation, even though almost all field-theoretical results were attained by the relativistic invariant  $S$ -matrix method. Similarly with the Wheeler-DeWitt equation – it lies at the foundation of quantum gravity theory of the Universe as a whole, even though its direct application to concrete problems might be ineffective and should be replaced by such advanced, though technical, tools as path integration. This is the main objective of Chapter 7.

The idea to write this book was born after the reprint volume with commentaries by Friedrich Hehl and Milutin Blagojević “Gauge Theory of Gravitation” was published in 2013, referring to our monograph “Geometrodynamical methods and gauge approach in the theory of gravitational interactions” which appeared in Russian in 1985. An original plan was to translate that old book into English. However, soon it became clear to the authors that after the 30 years some of the material became outdated and should be removed, whereas important new results which appeared in the meantime (also due to the authors) should be included. A need of the serious changes became obvious, and a new deeply revised book was written, now in English. In particular, the discussion of


the classical gauge models and the quantum gravity methods has undergone an essential revision. At the same time, we decided to preserve the general structure of the monograph and to keep the overall presentation style which was, in our opinion, quite successfully chosen in the old book.

We did not intend to describe in detail the modern mathematical methods of the classical gravity theory, which can be found in [2, 3, 4]. The summary of the relevant notions and definitions is given in Appendix A3. At the same time, all the derivations and constructions necessary for understanding of the book are given in full. We also did not try to provide a comprehensive review of the literature on the subject. This is to some extent compensated by the comments to the literature presented at the end of the book in Appendix A4. Moreover, we compiled an exhaustive bibliography of more than 3000 publications on the gauge gravity theory, its physical aspects and related mathematical results. Since no special preselection was made, the readers should clearly understand that some items in this bibliography may be erroneous or misleading, however, we do hope that none of the important publications on the gauge gravity was overlooked, thus making this bibliography a useful resource on the subject.

The book addresses the physicists who specialize in the gravity theory and the high-energy physics, and it can be recommended to the graduate students and senior undergraduate university students in physics and mathematics, who are familiar with the classical and quantum field theory at the level of the textbooks “Classical field theory” of L.D. Landau and E.M. Lifshitz and “Introduction to the theory of quantized fields” of N.N. Bogoliubov and D.V. Shirkov.

The book is written jointly by the authors. Chapters 1, 4-6 are based on original works of Yu.N. Obukhov and V.N. Ponomarev. The material presented in Chapters 2, 3, and 7 is based on the results obtained by A.O. Barvinsky and V.N. Ponomarev.

To help the reader, we give an approximate scheme of dependence of the chapters:



The authors express their profound gratitude to all colleagues who took part in the discussion of results presented in the book. We are sincerely grateful to the scientific and administrative staff of the Nuclear Safety Institute of the Russian Academy of Sciences (IBRAE RAN) for the constant and generous support and for the assistance in preparing the manuscript for publication. One of the authors (A.O.B.) is grateful for hospitality to the Theory Division of CERN where his work on the book was accomplished. Y.N.O. acknowledges a partial support from the Russian Foundation for Basic Research (Grant No. 16-02-00844-A).

*Authors*

# 1

## Lagrangian description of gravity theories of Hilbert-Einstein type

### 1.1. Geometrical spacetime structures

In the framework of the geometrodynamical approach, we model the spacetime as a smooth four-dimensional manifold  $M_4$ . On the latter, one can introduce various geometrical structures to describe the gravitational field. In this section, we present a brief overview of the most important ones: the metric and the affine connection structure. The summary of the main notions, definitions and methods of the modern differential geometry can be found in Appendix A1, for a more detailed exposition see [2]-[4]. Our basic notations are given on page 173.

#### *Metric structure (g-structure)*

A differentiable manifold  $M_4$  is called a Riemannian space  $V_4$ , when the metric is defined on it, that is when a smooth tensor field  $g_{\mu\nu}(x^\lambda)$ , such that  $g_{\mu\nu}(x^\lambda) = g_{\nu\mu}(x^\lambda)$ , introduces the metricity relations on the manifold: for any two infinitely close points  $x$  and  $x + dx$ , the interval (“distance”) between them is defined by

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu.$$

The metric defines a scalar product on the manifold by assigning to any two vectors  $A$  and  $B$  a real number  $AB = g_{\mu\nu} A^\mu B^\nu$ , and thereby introduces for each vector  $V$  its length  $l^2(V) = g_{\mu\nu} V^\mu V^\nu$ . By definition, the metric is assumed to be non-degenerate in the sense that the determinant  $g = \det g_{\mu\nu} \neq 0$ . This allows to determine the matrix  $g^{\mu\nu}(x^\lambda)$  inverse to  $g_{\mu\nu}(x^\lambda)$  such that  $g^{\mu\nu} g_{\nu\lambda} = \delta_\lambda^\mu$ , thereby establishing an isomorphism between the covariant and contravariant tensor

spaces. In local coordinates, this isomorphism is described by the operation of raising and lowering of indices. The metric is called Lorentzian if its signature is  $\pm 2$ . The Lorentzian metric structure forms the basis of the general relativity theory (GR).

### *Connection structure ( $\Gamma$ -structure)*

Independently of the metric structure, one can define the structure of an affine connection on  $M_4$  manifold. The linear (affine) connection  $\Gamma$  introduces the local isomorphism of tangent spaces at different points  $x \in M_4$  of the manifold, by specifying a rule that maps a vector  $V$  (an element of the tangent space) at a point  $x$  into a vector  $V'$  at an infinitely near point  $x + dx$ . In the local coordinates, this law is written as

$$\delta V^\mu = V'^\mu - V^\mu = -\Gamma^\mu_{\nu\alpha} V^\nu dx^\alpha.$$

Thus, equivalently, we say that the affine connection  $\Gamma$  is introduced on  $M_4$  manifold, when the global (i.e., given in each chart of the atlas of the manifold) field  $\Gamma^\mu_{\nu\alpha}(x)$  is defined on it. When changing one coordinate chart  $\{x\}$  to another one  $\{x'\}$ , the quantities  $\Gamma^\mu_{\nu\alpha}(x)$ , which are called coefficients of the linear (affine) connection, are transformed as

$$\Gamma'^\alpha_{\beta\mu}(x') = \left[ \frac{\partial x^\alpha}{\partial x'^\rho} \frac{\partial x'^\sigma}{\partial x^\beta} \Gamma^\rho_{\sigma\nu}(x) + \frac{\partial x^\alpha}{\partial x'^\rho} \frac{\partial}{\partial x^\nu} \left( \frac{\partial x'^\rho}{\partial x^\beta} \right) \right] \frac{\partial x'^\nu}{\partial x^\mu}.$$

By defining of the connection structure, one can introduce the notion of a covariant derivative of a tensor. The covariant derivative with respect to an arbitrary connection  $\Gamma$  is denoted by  $\nabla$  and is defined as follows. Suppose that in the local coordinates, the tensor field has components  $T^{\alpha_1 \dots \alpha_p}_{\beta_1 \dots \beta_q}$ . Then

$$\begin{aligned} \nabla_\mu T^{\alpha_1 \dots \alpha_p}_{\beta_1 \dots \beta_q} = & \partial_\mu T^{\alpha_1 \dots \alpha_p}_{\beta_1 \dots \beta_q} + \Gamma^{\alpha_1}_{\rho\mu} T^{\rho\alpha_2 \dots \alpha_p}_{\beta_1 \dots \beta_q} \dots + \Gamma^{\alpha_p}_{\rho\mu} T^{\alpha_1 \dots \alpha_{p-1}}_{\beta_1 \dots \beta_q} \\ & - \Gamma^\sigma_{\beta_1\mu} T^{\alpha_1 \dots \alpha_p}_{\sigma \dots \beta_q} \dots - \Gamma^\sigma_{\beta_q\mu} T^{\alpha_1 \dots \alpha_p}_{\beta_1 \dots \sigma}. \end{aligned}$$

The connection also defines the notion of a parallel transport of tensors. Let  $\gamma(t)$  be a smooth curve in  $M_4$ , specified by the parametric equations  $\gamma = \{x^\mu(t)\}$ . Then we define the covariant derivative of a tensor field  $T$  along this curve,  $DT/dt$ . In the local coordinates, it is equal to  $DT/dt = \frac{dx^\mu}{dt} \nabla_\mu T$ .

The manifold  $M_4$  with the connection structure defined on it is called an affinely connected space (denoted by  $L_4$ ), and it is characterized by a number of geometrical objects which we briefly consider below.

In general, we assume that both structures—the metric and the connection—are introduced on the manifold. By definition, they are independent. We call such a space a generalized metric-affine spacetime  $G_4$ .

Let us consider the parallel transport of a vector  $V$  from a point  $x \in G_4$  along some closed curve  $\gamma \subset G_4$ . When returning back to  $x$ , one finds that in general case the final (parallelly transported) vector  $V'$  does not coincide with the original one  $V$ . The difference is manifest in the three effects: the vector  $V'$

is rotated with respect to  $V$ , its length is changed, and the image of the contour  $\gamma$  in the tangent space turns out to be broken by a non-zero vector.

These effects are determined by the following basic geometrical objects of  $G_4$ :  
The curvature tensor of the connection  $\Gamma$

$$R^\alpha{}_{\beta\mu\nu} = \partial_\mu \Gamma^\alpha{}_{\beta\nu} - \partial_\nu \Gamma^\alpha{}_{\beta\mu} + \Gamma^\alpha{}_{\rho\mu} \Gamma^\rho{}_{\beta\nu} - \Gamma^\alpha{}_{\rho\nu} \Gamma^\rho{}_{\beta\mu}; \quad (1.1)$$

the homothetic curvature tensor

$$\Omega_{\mu\nu} = \partial_\mu \Gamma_{\cdot\nu} - \partial_\nu \Gamma_{\cdot\mu}, \quad (1.2)$$

where  $\Gamma_{\cdot\mu} := \Gamma^\lambda{}_{\lambda\mu}$ ;

and the torsion tensor [20]

$$Q^\alpha{}_{\mu\nu} = \Gamma^\alpha{}_{[\mu\nu]} = \frac{1}{2}(\Gamma^\alpha{}_{\mu\nu} - \Gamma^\alpha{}_{\nu\mu}). \quad (1.3)$$

In the local coordinates, the aforementioned effects of the parallel transport along a closed loop read as follows:

- the rotation of a vector:

$$\delta V^\alpha \approx R^\alpha{}_{\beta\mu\nu} V^\beta ds^{\mu\nu}, \quad (1.4)$$

- the change of the length:

$$\delta l \approx l(V) \Omega_{\mu\nu} ds^{\mu\nu}, \quad (1.5)$$

- the contour image in the tangent space is broken by the vector:

$$\xi^\alpha \approx 2Q^\alpha{}_{\mu\nu} ds^{\mu\nu}. \quad (1.6)$$

Here  $ds^{\mu\nu}$  is the surface element spanned by the closed contour  $\gamma$ .

We introduce now another fundamental tensor which is an important characteristic of  $G_4$  manifold, even being not entirely independent of the tensors introduced above. This is the nonmetricity tensor that measures the (in)compatibility of the independent metric and connection. It is equal to the covariant derivative of the metric  $g_{\alpha\beta}$ :

$$K_{\mu\alpha\beta} = \nabla_\mu g_{\alpha\beta}. \quad (1.7)$$

We say that the metric and connection are compatible when  $K_{\mu\alpha\beta} = 0$  vanishes on  $M_4$ , i.e., when the metricity condition is fulfilled

$$\nabla_\mu g_{\alpha\beta} = 0. \quad (1.8)$$

Let us consider some properties of the fundamental tensors. A closer inspection of the aforementioned non-integrable effects (1.4)-(1.6) shows that they are due to the fact that covariant derivatives  $\nabla$  do not commute with each other. For any vector  $V$  we find

$$(\nabla_\mu \nabla_\nu - \nabla_\nu \nabla_\mu) V^\alpha = R^\alpha{}_{\beta\mu\nu} V^\beta + 2Q^\beta{}_{\mu\nu} \nabla_\beta V^\alpha.$$

This result can be generalized to arbitrary tensors. In particular, by considering the commutator of covariant derivatives of the metric, we obtain the useful identities relating the curvature tensors (1.1) and the nonmetricity (1.7):

$$2\nabla_{[\mu}K_{\nu]\alpha\beta} = 2\nabla_{[\mu}\nabla_{\nu]}g_{\alpha\beta} = -2R_{(\alpha\beta)\mu\nu} + 2Q^{\rho}_{\mu\nu}K_{\rho\alpha\beta}. \quad (1.9)$$

As a result, we find that the symmetric part of the curvature tensor in the first pair of indices  $R_{(\alpha\beta)\mu\nu}$  is nontrivial when  $K_{\alpha\mu\nu} \neq 0$ . Contracting eq. (1.9) with  $g^{\alpha\beta}$ , we recover the homothetic curvature

$$g^{\alpha\beta}R_{\alpha\beta\mu\nu} = R^{\alpha}_{\alpha\mu\nu} = \Omega_{\mu\nu} = -\frac{1}{2}(\partial_{\mu}K_{\nu} - \partial_{\nu}K_{\mu}), \quad (1.10)$$

where  $K_{\mu} = K_{\mu\alpha\beta}g^{\alpha\beta}$  is the vector of nonmetricity (or otherwise, the Weyl vector field [5]).

The equation (1.7) can be solved with respect to the connection  $\Gamma^{\lambda}_{\mu\nu}$  so that the latter is expressed in terms of the metric, torsion and nonmetricity:

$$\Gamma^{\lambda}_{\mu\nu} = \{\overset{\lambda}{\mu\nu}\} + D^{\lambda}_{\mu\nu}. \quad (1.11)$$

Here

$$\{\overset{\lambda}{\mu\nu}\} = \frac{1}{2}g^{\lambda\sigma}(\partial_{\nu}g_{\mu\sigma} + \partial_{\mu}g_{\nu\sigma} - \partial_{\sigma}g_{\mu\nu})$$

is the Christoffel symbol, and

$$D^{\lambda}_{\mu\nu} = Q^{\lambda}_{\mu\nu} + Q_{\mu\nu}{}^{\lambda} + Q_{\nu\mu}{}^{\lambda} - \frac{1}{2}g^{\rho\lambda}(K_{\nu\mu\rho} + K_{\mu\nu\rho} - K_{\rho\mu\nu})$$

is the connection defect tensor (also called a distortion tensor). The connection  $\{\overset{\lambda}{\mu\nu}\}$  is the only torsion-less connection which is compatible with the metric structure. It is called a Riemannian or a metric connection.

An arbitrary connection with a non-zero torsion, but compatible with metric, is called the Riemann-Cartan connection. Its explicit structure is obtained from (1.11) if we put  $K_{\alpha\mu\nu} = 0$ :

$$\Gamma^{\lambda}_{\mu\nu} = \{\overset{\lambda}{\mu\nu}\} + Q^{\lambda}_{\mu\nu} + Q_{\mu\nu}{}^{\lambda} + Q_{\nu\mu}{}^{\lambda} = \{\overset{\lambda}{\mu\nu}\} + T^{\lambda}_{\mu\nu}. \quad (1.12)$$

The combination

$$T^{\lambda}_{\mu\nu} = Q^{\lambda}_{\mu\nu} + Q_{\mu\nu}{}^{\lambda} + Q_{\nu\mu}{}^{\lambda}$$

is usually called a contortion tensor. From now on we will confine ourselves to the Riemann-Cartan spacetime  $U_4$  with the metric and the Riemann-Cartan connection (1.12).

The fundamental tensors in  $U_4$  have a large number of independent components (24 for the torsion and 36 for the curvature), and it is convenient to decompose these geometrical objects into the  $L_6$ -irreducible pieces (where  $L_6 = SO(3, 1)$  is the Lorentz group). The decomposition of the torsion tensor into the three  $L_6$ -irreducible parts reads as follows:

$$Q^{\lambda}_{\mu\nu} = \bar{Q}^{\lambda}_{\mu\nu} + \frac{2}{3}\delta^{\lambda}_{[\nu}Q_{\mu]} + g^{\lambda\sigma}\varepsilon_{\mu\nu\rho\sigma}\check{Q}^{\rho}. \quad (1.13)$$

Here  $\bar{Q}^\lambda{}_{\mu\nu}$  is the traceless part of the torsion tensor, with the properties:  $\bar{Q}^\lambda{}_{\mu\lambda} = 0$  and  $\varepsilon^{\lambda\mu\nu\rho}\bar{Q}_{\mu\nu\rho} = 0$ ; the torsion trace vector is  $Q_\mu := Q^\lambda{}_{\mu\lambda}$ ; and  $\check{Q}_\mu = \frac{1}{6}\varepsilon_{\mu\nu\lambda\sigma}Q^{\nu\lambda\sigma}$  is the torsion pseudotrace vector ( $\varepsilon_{\lambda\mu\nu\chi}$  is the totally anti-symmetric Levi-Civita tensor:  $\varepsilon_{\lambda\mu\nu\chi} = \varepsilon_{[\lambda\mu\nu\chi]}$ ).

In a similar way, one can decompose the curvature tensor in  $U_4$  spacetime into the 6 irreducible pieces [26]:

$$\begin{aligned} R_{\alpha\beta\mu\nu}(\Gamma) &= C_{\alpha\beta\mu\nu}(\Gamma) + \frac{1}{2} \{g_{\alpha\mu}R_{\beta\nu}(\Gamma) - g_{\alpha\nu}R_{\beta\mu}(\Gamma) - g_{\beta\mu}R_{\alpha\nu}(\Gamma) \\ &\quad + g_{\beta\nu}R_{\alpha\mu}(\Gamma)\} - \frac{1}{6} (g_{\alpha\mu}g_{\beta\nu} - g_{\alpha\nu}g_{\beta\mu}) R(\Gamma) \\ &\quad + \frac{1}{12} \varepsilon_{\alpha\beta\mu\nu}D(\Gamma) + \frac{1}{4} \{ \varepsilon_{\alpha\beta\mu}{}^\lambda D_{(\lambda\nu)}(\Gamma) - \varepsilon_{\alpha\beta\nu}{}^\lambda D_{(\lambda\mu)}(\Gamma) \\ &\quad + \varepsilon_{\mu\nu\beta}{}^\lambda D_{(\lambda\alpha)}(\Gamma) - \varepsilon_{\mu\nu\alpha}{}^\lambda D_{(\lambda\beta)}(\Gamma) \}. \end{aligned} \quad (1.14)$$

Here  $C_{\mu\lambda\nu\chi}(\Gamma)$  is the non-Riemannian analog of the Weyl tensor with the same algebraic properties

$$\begin{aligned} C_{\lambda\mu\nu\chi}(\Gamma) &= C_{\nu\chi\lambda\mu}(\Gamma), \quad C^\lambda{}_{\mu\lambda\nu}(\Gamma) = 0, \quad C_{[\lambda\mu\nu]\chi}(\Gamma) = 0, \\ C_{\lambda\mu\nu\chi}(\Gamma) &= C_{[\lambda\mu]\nu\chi}(\Gamma) = C_{\lambda\mu[\nu\chi]}(\Gamma), \end{aligned}$$

$R_{\mu\nu}(\Gamma) = R^\lambda{}_{\mu\lambda\nu}(\Gamma)$  is the generalized Ricci tensor,  $R(\Gamma) = g^{\mu\nu}R_{\mu\nu}(\Gamma)$  is the curvature scalar, and  $D^\mu{}_\nu(\Gamma) = \frac{1}{2}\varepsilon^{\mu\alpha\beta\lambda}R_{\alpha\beta\lambda\nu}(\Gamma)$  so that  $D(\Gamma) = D^\mu{}_\mu(\Gamma)$ . The list of the 6 irreducible curvature part includes the generalized Weyl tensor, the curvature scalar, the curvature pseudoscalar, the skew-symmetric part of the Ricci tensor, and the two traceless symmetric tensors:

$$\begin{aligned} C_{\alpha\beta\mu\nu}(\Gamma), \quad R(\Gamma), \quad R_{(\mu\nu)}(\Gamma) - \frac{1}{4}R(\Gamma)g_{\mu\nu}, \\ R_{[\mu\nu]}(\Gamma), \quad D(\Gamma), \quad D_{(\mu\nu)}(\Gamma) - \frac{1}{4}D(\Gamma)g_{\mu\nu}. \end{aligned}$$

The total number of components is, as expected:  $36 = 20 + 1 + 9 + 6 + 1 + 9$ . The second line represents the essentially non-Riemannian irreducible parts. When the torsion (1.3) vanishes, eq. (1.14) reduces to the standard decomposition for Riemann-Christoffel curvature tensor [27] into the three pieces shown in the first line.

It is worthwhile to note that the skew-symmetric tensor  $D_{[\alpha\beta]}(\Gamma)$  does not represent an independent part. It is expressed in terms of the antisymmetric part of the Ricci tensor:  $D_{[\mu\nu]}(\Gamma) = -\frac{1}{2}\varepsilon_{\mu\nu\alpha\beta}R^{[\alpha\beta]}(\Gamma)$ .

The Riemann-Cartan curvature (1.12) is antisymmetric in the first pair of indices, as it is obvious from (1.9), and it satisfies the following identities:

$$R^\alpha{}_{[\mu\nu]\lambda}(\Gamma) = -2\nabla_{[\mu}Q^\alpha{}_{\nu\lambda]} + 4Q^\alpha{}_{\beta[\mu}Q^\beta{}_{\nu\lambda]} \quad (1.15)$$

– the generalized Ricci identity;

$$\begin{aligned} R_{\alpha\beta\mu\nu}(\Gamma) - R_{\mu\nu\alpha\beta}(\Gamma) &= \frac{3}{2} \{ R_{[\alpha\beta\mu]\nu}(\Gamma) - R_{[\alpha\beta\nu]\mu}(\Gamma) \\ &\quad - R_{[\mu\nu\alpha]\beta}(\Gamma) + R_{[\mu\nu\beta]\alpha}(\Gamma) \}, \end{aligned} \quad (1.16)$$

– the “commutator of pairs of indices” identity;

$$\nabla_{[\mu} R^{\alpha\beta}{}_{\nu\lambda]}(\Gamma) = 2R^{\alpha\beta}{}_{\sigma[\mu}(\Gamma) Q^{\sigma}{}_{\nu\lambda]} \quad (1.17)$$

– the generalized Bianchi identity.

Contracting a pair of indices (upper and lower) in (1.15), we derive a so-called contracted Ricci identity

$$R_{[\mu\nu]} = (\nabla_\lambda - 2Q_\lambda) \left( Q^\lambda{}_{\mu\nu} + 2\delta_{[\mu}^\lambda Q_{\nu]} \right).$$

Similarly, contracting the two pairs of indices in (1.17), we find the contracted Bianchi identities

$$(\nabla_\sigma - 2Q_\sigma) G^\sigma{}_\mu(\Gamma) + 2Q^\rho{}_{\mu\sigma} G^\sigma{}_\rho(\Gamma) + (Q^\sigma{}_{\alpha\beta} + 2\delta_{[\alpha}^\sigma Q_{\beta]}) R^{\alpha\beta}{}_{\sigma\mu}(\Gamma) = 0, \quad (1.18)$$

where

$$G_{\lambda\nu}(\Gamma) = R_{\lambda\nu}(\Gamma) - \frac{1}{2} g_{\lambda\nu} R(\Gamma)$$

is the generalized Einstein tensor.

With the help of the geometrical objects introduced above, one can establish a natural classification of the spacetime theories in accordance with their underlying metric-affine structures, see Table 1.1.

### *Tetrad structure*

Along with the so-called world geometrical structures which we described above, in the gravity theory one effectively uses their local Lorentz analogues. In the latter approach, the main object is tangent bundle  $T(M)$ , and the related bundle of orthonormal frames  $AO(M)$  which is the principal fibre bundle with the Lorentz  $L_6$  structure group.

All the geometrical objects of the spacetime theory had previously been defined in the local coordinates on the manifold with respect to the world or coordinate frame  $\partial_\mu \equiv \partial/\partial x^\mu$  in the tangent space  $T_x(M)$ . However, it is often convenient (and when considering spinors in  $M_4$ , it is necessary) to define all the objects with respect to arbitrary orthonormal basis in  $T(M)$ ,  $\{e_a\}$ ,  $a = 0, 1, 2, 3$ . Such a frame is orthonormal in the sense that the scalar products of its legs are equal  $(e_a \cdot e_b) = \eta_{ab} = \text{diag}(-1, +1, +1, +1)$  – the Minkowski metric. We call such a basis a local Lorentz frame, and accordingly all the geometrical objects are also called the local Lorentz objects when they are considered with respect to this frame. In order to distinguish the components with respect to  $\{e_a\}$  from the components with respect to  $\{\partial_\mu\}$ , we will use the Latin indices  $(a, b, c, \dots)$  for the local Lorentz frames, while the Greek indices  $(\mu, \nu, \lambda, \dots)$  will label the world quantities with respect to  $\{\partial_\mu\}$ .

The transition from a coordinate frame to an arbitrary orthonormal frame is described by the tetrad coefficients  $h_\mu^a$ . By definition

$$\partial_\mu = h_\mu^a e_a.$$

Table 1.1: Classification of metric-affine theories

Geometrical objects	Type of space	Example of theories
$R_{..} = 0 \quad Q_{..} = 0 \quad \Omega_{..} = 0$	Minkowski space $M_4$	Special relativity theory
$R_{..} = 0 \quad Q_{..} \neq 0 \quad \Omega_{..} = 0$	Weitzenböck (or absolute parallelism) space $P_4$	Translational gauge gravity theory [34]
$R_{..} \neq 0 \quad Q_{..} = 0 \quad \Omega_{..} = 0$	Riemann space $V_4$	General relativity theory (GR) [27]
$R_{..} \neq 0 \quad Q_{..} = 0 \quad \Omega_{..} \neq 0$	Weyl space $W_4$	Weyl's gravity theory [5]
$R_{..} \neq 0 \quad Q_{..} \neq 0 \quad \Omega_{..} = 0$	Riemann-Cartan space $U_4$	Einstein-Cartan gravity theory (ECT) [7, 6]
$R_{..} \neq 0 \quad Q_{..} \neq 0 \quad \Omega_{..} \neq 0$	Generalized metric-affine space $G_4$	Various asymmetric field theories, in particular Einstein-Schrödinger [9]

These quantities are closely related to the spacetime metric, namely

$$g_{\mu\nu} = h_\mu^a h_\nu^b \eta_{ab}. \quad (1.19)$$

In the bundle of the Lorentz frames, one can define a connection  $\Gamma^a{}_{b\mu}$  that introduces their parallel transport. The latter is naturally induced by the world affine connection  $\Gamma^\lambda{}_{\mu\nu}$ . The coefficients of the local Lorentz connection read as follows:

$$\Gamma^a{}_{b\mu} = h_\alpha^a h_b^\beta \Gamma^\alpha{}_{\beta\mu} + h_\sigma^a \partial_\mu h_b^\sigma. \quad (1.20)$$

Here, the quantity  $h_\mu^a$  is defined as an inverse matrix to  $h_a^\mu$ :

$$h_\mu^a h_a^\nu = \delta_\mu^\nu, \quad h_\mu^a h_b^\mu = \delta_b^a.$$

The frame  $e_a$  is called nonholonomic, if the corresponding tetrad coefficients  $h_\mu^a$  are no partial derivatives of some four functions  $f^a(x)$ ,  $a = 0, 1, 2, 3$ , i.e., when  $h_\mu^a \neq \partial_\mu f^a$ . Geometrically, it means that there are no such local coordinates  $y^a = f^a(x)$ , with respect to which  $e_a$  becomes a coordinate basis. The degree of deviation from the field of integrable local Lorentz frames, for which such new coordinates exist, is measured by the anholonomy object

$$C^a{}_{\mu\nu} := \partial_\mu h_\nu^a - \partial_\nu h_\mu^a. \quad (1.21)$$

This object is not a tensor, and it is closely related to the tetrad form of the Riemannian connection induced by the metric. As a result, one can reformulate Einstein's GR in terms of the tetrad  $h_\mu^a$  and the anholonomy object  $C^a{}_{\mu\nu}$  [8].

Computing the commutator of covariant derivatives  $D_\mu v^a = \partial_\mu v^a + \Gamma^a{}_{b\mu} v^b$  with respect to the local Lorentz connection, we obtain the local Lorentz curvature

$$R^a{}_{b\mu\nu} = \partial_\mu \Gamma^a{}_{b\nu} - \partial_\nu \Gamma^a{}_{b\mu} + \Gamma^a{}_{c\mu} \Gamma^c{}_{b\nu} - \Gamma^a{}_{c\nu} \Gamma^c{}_{b\mu}.$$

It is easy to show that the latter is related to the curvature tensor of the world affine connection,

$$R^a{}_{b\mu\nu} = h^a_\alpha h_b^\beta R^\alpha{}_{\beta\mu\nu}.$$

Analogously, we can prove that it is possible to recast the torsion tensor into a generalized covariant “curl” of the tetrad:

$$D_\mu h_\nu^a - D_\nu h_\mu^a = -2h_\lambda^a Q^\lambda{}_{\mu\nu}.$$

The tetrad formalism represents another framework for the gravity theory. The metric and the affine connection  $(g_{\mu\nu}, \Gamma^\alpha{}_{\beta\mu})$  in this approach are replaced by the tetrad and the local Lorentz connection  $(h^a_\mu, \Gamma^a{}_{b\mu})$ . It will be shown later that there is close relation of the latter variables with the structure of the principal bundle of the Lorentz group and it turns out to be possible to develop a consistent gauge approach to the gravity theory using this formulation.

Apparently, the most convenient mathematical formalism for the gauge approach in gravity is the theory of connections in the bundles over the spacetime manifold  $M_4$ . See Chapter 5 for the further details.

## 1.2. Hilbert variational principle and field equations of the general relativity

Einstein [22] formulated the fundamental equations of GR describing the dynamics of the interacting gravitational and matter fields on the basis of an idea that matter (its energy-momentum tensor  $T^{\mu\nu}$ ) gives rise to the Riemannian curvature of spacetime. An additional requirement was that the covariant conservation law  $T^{\mu\nu}{}_{;\nu} = 0$  should be an automatic consequence of the field equations. The latter assumption almost uniquely fixed the Einstein tensor  $G_{\mu\nu}$  as an appropriate structure describing the geometry of the Riemannian spacetime by making use of the contracted Bianchi identity (1.18) for the case of the vanishing torsion  $Q^\lambda{}_{\mu\nu} = 0$ .

As a result, Einstein’s gravitational field equations read as follows:

$$G_{\mu\nu} = \kappa T_{\mu\nu}, \tag{1.22}$$

where the value of the Einstein gravitational constant  $\kappa = 8\pi G/c^4 = 2.07 \times 10^{-43} \text{ kg}^{-1} \text{ m}^{-1} \text{ s}^2$  was derived from the comparison of GR with Newton’s gravitation theory;  $G = 6.67 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$  is the Newtonian gravitational constant.

In a vacuum (in the absence of matter), these equations reduce to

$$G_{\mu\nu} = 0 \quad \text{or} \quad R_{\mu\nu} = 0. \tag{1.22a}$$

At the same time, the field equations (1.22a) were derived by Hilbert [23] on the basis of the fundamental dynamical principle – the principle of the least action,

$$S_{HE} = \frac{1}{2\kappa c} \int d^4x \sqrt{g} \mathcal{L}_{HE} = \frac{1}{2\kappa c} \int d^4x \sqrt{g} R.$$

Gravity models based on Lagrangian which is linear on curvature, are called the gravity theories of the Hilbert-Einstein type.

A short remark is in order concerning the dimensions. The metric components are dimensionless, whereas the connection coefficients have a 1/length dimension. Accordingly, the curvature and its contractions have a dimension of  $1/(\text{length})^2$ . Consequently, the dimension of the integral is  $[d^4x R] = \text{m}^2$ . Taking into account that  $[\frac{1}{\kappa c}] = \text{kg/s}$ , we find that the action  $[S_{HE}] = \text{kg m}^2/\text{s} = [\hbar]$  indeed has the dimension of the action (recall that the Planck constant is a “quantum of an action”).

By substituting the expression of the Christoffel symbols  $\{\lambda_{\mu\nu}\}$  in terms of the metric  $g_{\mu\nu}$  into  $\mathcal{L}_{HE}$ , we obtain [27]

$$\begin{aligned} \mathcal{L}_{HE} &= \mathcal{L}_{HE}[g_{\mu\nu,\lambda\kappa}, g_{\mu\nu,\kappa}, g_{\mu\nu}] = g^{\mu\nu} \left( 2\{\lambda_{\mu[\nu}, \lambda]\} + 2\{\sigma[\lambda\} \{\nu\}_{\mu}\} \right) \\ &= \frac{1}{2} (g_{\mu\nu,\sigma\kappa} + g_{\sigma\kappa,\mu\nu} - g_{\mu\kappa,\sigma\nu} - g_{\sigma\nu,\mu\kappa}) g^{\mu\kappa} g^{\sigma\nu} \\ &\quad + g^{\mu\kappa} g^{\sigma\nu} g_{\alpha\beta} (\{\sigma\kappa\} \{\mu\nu\} - \{\sigma\nu\} \{\mu\kappa\}). \end{aligned}$$

Commas denote the usual partial derivatives,  $_{,\lambda} := \partial_\lambda$ , etc. Let us vary the action  $S_{HE}$  with respect to the metric  $g_{\mu\nu}$ . Then

$$\begin{aligned} \delta_g S_{HE} &= \frac{1}{2\kappa c} \int_{\Omega} d^4x \sqrt{g} G_{\mu\nu} \delta g^{\mu\nu} \\ &\quad + \frac{1}{2\kappa c} \oint_{\partial\Omega} d\sigma_\lambda \sqrt{g} g^{\mu\kappa} [-\delta_g \{\lambda_{\mu\kappa}\} + \delta_\mu^\lambda \delta_g \{\rho_{\kappa\rho}\}] = 0, \quad (1.23) \\ d\sigma_\lambda &= \frac{1}{3!} \varepsilon_{\lambda\mu\nu\chi} dx^{[\mu} dx^\nu dx^{\chi]}. \end{aligned}$$

Einstein’s equations (1.22a) follow from (1.23), if and only if the surface term vanishes due to the conditions imposed by the respective boundary problem

$$\oint_{\partial\Omega} d\sigma_\lambda \sqrt{g} g^{\mu\kappa} [-\delta_g \{\lambda_{\mu\kappa}\} + \delta_\mu^\lambda \delta_g \{\rho_{\kappa\rho}\}] = 0, \quad (1.24)$$

that is when the variational principle is consistent with the boundary value problem of the resulting Euler-Lagrange equations. The boundary conditions specify the appropriate test functions, and thereby restrict the class of the variational problems [10].

Einstein’s equations are the second-order partial differential equations with respect to metric  $g_{\mu\nu}$  in the spacetime domain  $\Omega$ , hence an appropriate Dirichlet boundary value problem for them fixes the values of the metric at the boundary  $\partial\Omega$  of the spacetime domain,  $g_{\mu\nu}|_{\partial\Omega} = C_{\mu\nu}$ . However, the corresponding condition on the variations,  $\delta g_{\mu\nu}|_{\partial\Omega} = 0$  is clearly not sufficient to make the surface term (1.24) vanish, in general. The only exceptions are the gravitational fields of isolated matter configurations (the asymptotically flat metrics).

Strictly speaking, the statement that a physical theory is based on the variational principle  $\delta S = 0$  is valid only when the system of the Euler-Lagrange equations is solved under the boundary conditions obtained from the stationarity condition of the action  $S$ . In other words, the boundary conditions obtained from the stationarity conditions should not contradict the order of Euler-Lagrange equations, i.e., the corresponding Lagrangian is a proper (nondegenerate) one.

For a given boundary value problem, a proper (nondegenerate) Lagrangian is not linear in the higher derivatives of the field functions. Making use of this definition, we conclude that the Hilbert Lagrangian of the gravitational field  $\mathcal{L}_{HE} = R$  is degenerate [11].

Let us illustrate the above by the example from mechanics. Consider the two Lagrangians  $\mathcal{L}_1 = -\frac{1}{2}q\ddot{q}$ , and  $\mathcal{L}_2 = \frac{1}{2}\dot{q}^2$  ( $q$  is the generalized coordinate, the dot denotes the time differentiation). Variations of the respective action functionals are given by

$$\delta S_1 = - \int_{t_1}^{t_2} \ddot{q} \delta q dt - \frac{1}{2} q \delta \dot{q} \Big|_{t_1}^{t_2} + \frac{1}{2} \dot{q} \delta q \Big|_{t_1}^{t_2}, \quad (1.25)$$

$$\delta S_2 = - \int_{t_1}^{t_2} \ddot{q} \delta q dt + \frac{1}{2} \dot{q} \delta q \Big|_{t_1}^{t_2}. \quad (1.26)$$

Both actions yield the same Euler-Lagrange equation  $\ddot{q} = 0$ . However, eq. (1.25) tells us that in order to obtain the equation of motion, it is necessary to fix the values of the coordinate  $q(t_1)$ ,  $q(t_2)$ , and the values of the velocities  $\dot{q}(t_1)$ ,  $\dot{q}(t_2)$  at the boundary. But this is impossible in view of the order of these equations. The attempt in (1.25) to impose only two conditions, for example  $(q/\dot{q})|_{t_1}^{t_2}$ , does not lead to a physically meaningful result. As for (1.26), the equation of motion  $\ddot{q} = 0$  follows from the usual variational problem with the fixed endpoints  $\delta q(t_1) = \delta q(t_2) = 0$ , consistent with the correct boundary value problem.

In a sense,  $\mathcal{L}_1$  in this mechanical example is similar to the Hilbert Lagrangian  $\mathcal{L}_{HE}$ , and  $\mathcal{L}_2$  is similar to the so-called truncated Lagrangian

$$\mathcal{L}_D = \mathcal{G} = g^{\mu\nu} \left( -\{\sigma_\lambda\} \{\sigma_{\mu\nu}\} + \{\sigma_\nu\} \{\sigma_{\mu\lambda}\} \right),$$

see [40]. The latter is a proper Lagrangian that leads to the self-consistent boundary value problem  $G_{\mu\nu} = 0$ ,  $g_{\mu\nu}|_{\partial\Omega} = C_{\mu\nu} = \text{const}$ .

It is worthwhile to note that  $\mathcal{L}_D$  (not  $\mathcal{L}_{HE}$ ) up to a boundary term coincides with the Lagrangian in the gauge theory of the translation group [13]:

$$\mathcal{L}_{T_4} = -\frac{1}{4} C_{abc} C^{abc} - \frac{1}{2} C_{abc} C^{bac} + C_{ab}{}^b C^a{}_c{}_c.$$

Here  $C^a{}_{bc} = h_b^\mu h_c^\nu C^a{}_{\mu\nu}$ .

In Hawking's approach [14], the truncated action is written as

$$S_D = S_{HE} + 2 \int_{\partial\Omega} d^4x \sqrt{g} K,$$

where the ordinary Hilbert-Einstein action is explicitly supplemented by the “surface term”. The boundary term is an integral of  $K$  which is the difference of traces of the second fundamental form of boundary  $\partial\Omega$  for the metric  $g$  and for the flat space metric. The role of the surface term is to compensate the contribution of the second derivatives of the metric in  $R$ , so that the Lagrangian becomes quadratic in the first derivatives (which means a nondegenerate), as required in the path integration method.

Along with Hilbert’s method of derivation of Einstein’s field equations from the variational principle with the action linear in the curvature, the gravitational field equations can be obtained within the framework of the first-order formalism, commonly called a Palatini principle [28], which we present in the next section.

### 1.3. First-order formalism and field equations of the Einstein-Cartan theory

In the derivations above, the Riemannian character of the spacetime structure was essential. However, we now recall that the metric and connection are defined as a priori independent structures on a differentiable manifold. Therefore, from the geometrical point of view, it would be more natural to consider  $g_{\mu\nu}$  and  $\Gamma^\lambda_{\mu\nu}$  as the independent dynamical variables of the gravitational theory. This idea underlied the variational principle of Palatini, who did not impose the metricity condition (1.8) as a constraint equation on  $g_{\mu\nu}$  and  $\Gamma^\lambda_{\mu\nu}$ . However, he kept the torsion tensor equal zero:  $Q^\lambda_{\mu\nu} = 0$ .

In this case, the field equations are obtained from the independent variation of the Hilbert action  $S_{HE} = \frac{1}{2\kappa c} \int_{\Omega} d^4x \sqrt{g} R(g_{\mu\nu}, \Gamma^\lambda_{(\mu\nu),\alpha}, \Gamma^\lambda_{(\mu\nu)})$  with respect to the metric  $g_{\mu\nu}$  and the symmetric connection  $\Gamma^\lambda_{(\mu\nu)}$ . The variation of  $S_{HE}$  with respect to the metric yields

$$G_{\mu\nu}(g_{\mu\nu}, \Gamma^\lambda_{(\mu\nu),\alpha}, \Gamma^\lambda_{(\mu\nu)}) = 0, \quad (1.27)$$

and from the variation with respect to  $\Gamma^\lambda_{\mu\nu}$  one obtains the metricity condition (1.8). If the latter is solved with respect to  $\Gamma^\lambda_{\mu\nu}$  (under the condition  $Q^\lambda_{\mu\nu} = 0$ ) and the result is plugged back into (1.27), we obtain Einstein’s field equation (1.22a) as before.

The vanishing torsion condition  $Q^\lambda_{\mu\nu} = 0$  is the constraint equation for the dynamical variables  $(g_{\mu\nu}, \Gamma^\lambda_{\mu\nu})$ . This fact was taken into account from the beginning of the derivation of the field equations by assuming  $\Gamma^\lambda_{\mu\nu} = \Gamma^\lambda_{(\mu\nu)}$ . If the constraints are not resolved from the very beginning, they should be added to the original Lagrangian with the undetermined Lagrange multipliers. Then the problem of derivation of the field equations becomes a variational problem on the conditional extremum of the action [15, 16, 17]. If the constraints are imposed after the field equations are derived, the resulting system would not be equivalent to the case, when the field equations are obtained by taking into

account constraints even before the variation. The exceptions arise when the Lagrange multipliers vanish due to field equations<sup>1</sup>.

We thus have demonstrated that Einstein's field equations in vacuum can be obtained by the Palatini variational method provided the torsion is equal zero. As we show below, by removing the restriction  $Q^\lambda{}_{\mu\nu} = 0$  it is possible to obtain more general gravitational field theories.

Let us consider the gravitational field dynamics based on the Palatini principle [18], assuming the full independence of the metric tensor from the general affine connection. The variation of Hilbert-Einstein type action (the subscript  $P$  stands for "Palatini")

$$S_P = \frac{1}{2\kappa_C} \int d^4x \sqrt{g} \mathcal{L}_P = \frac{1}{2\kappa_C} \int d^4x \sqrt{g} R(\Gamma)$$

yields, in the absence of matter (in vacuum):

$$\frac{\delta S_P}{\delta g_{\mu\nu}} = -G^{(\mu\nu)}(\Gamma) = 0, \quad (1.28)$$

$$\frac{\delta S_P}{\delta \Gamma^\lambda{}_{\mu\nu}} = -\nabla_\lambda g^{\mu\nu} + \delta^\nu_\lambda \nabla_\rho g^{\mu\rho} + 2Q^{\nu\mu}{}_\lambda + g^{\mu\nu} D_\lambda - D^\mu \delta^\nu_\lambda = 0, \quad (1.29)$$

where  $D_\lambda = D^\rho{}_{\lambda\rho}$  is the connection defect trace.

By splitting (1.29) into the symmetric and antisymmetric parts, we obtain:

$$Q^\lambda{}_{\mu\nu} = -\frac{2}{3} \delta^\lambda_{[\mu} Q_{\nu]}, \quad (1.30)$$

$$\nabla_\lambda g_{\mu\nu} = \frac{4}{3} Q_\lambda g_{\mu\nu}. \quad (1.31)$$

By solving the last equation with respect to  $\Gamma^\lambda{}_{\mu\nu}$ , we obtain the relation between the connection, the metric and the torsion

$$\Gamma^\lambda{}_{\mu\nu} = \{\lambda{}_{\mu\nu}\} - \frac{2}{3} \delta^\lambda_\mu Q_\nu. \quad (1.32)$$

Substituting (1.32) in (1.28), we find

$$G_{\mu\nu} = 0. \quad (1.33)$$

Summarizing, the following results were obtained by using the consistent Palatini principle:

- For the metric, the field equations (1.33) coincide with Einstein's equations of GR in vacuum.
- The torsion is expressed in terms of its trace  $Q_\lambda$  (1.30) (i.e., the connection (1.32) is semi-symmetric [24]) and remains arbitrary since the contraction of the equation (1.30) yields an identity.

---

<sup>1</sup>If the Lagrangian is supplemented by the term  $\lambda^{\lambda\mu\nu} Q_{\lambda\mu\nu}$  containing the Lagrange multipliers  $\lambda$ , the result is not changed, since  $\lambda^{\lambda\mu\nu} = 0$  due to the field equation.

- The covariant derivative of the metric (the nonmetricity) does not vanish (1.31) (the semi-metric parallel transport [24]).

However, the results obtained are not satisfactory since the torsion trace  $Q_\lambda$  is not fixed, therefore, all the main geometrical characteristics of the manifold are not determined. To find the way out of this situation, it is necessary to consider the variation principle with a different Lagrangian (see Chapter 5).

If we consider the theory with a Hilbert-Einstein Lagrangian, but require the preservation of vector lengths under the parallel transport, we obtain the Einstein-Cartan theory of gravity. The constraints (1.8) can be resolved from the very beginning and taken into account in action [7] or, without resolving them explicitly, they can be added to the Lagrangian with the undetermined Lagrange multipliers, and then the problem of derivation of the field equations becomes a variational problem on the constrained extremum of action [15, 16]. We consider the second procedure, including now into the dynamical scheme, along with the gravity, the matter fields interacting with the gravitational field. The gravitational interaction is introduced into the matter Lagrangian  $\mathcal{L}_m$  on the basis of the minimal coupling principle by replacing the Minkowski metric  $\eta_{\mu\nu}$  with the metric of the curved space  $g_{\mu\nu}(x^\lambda)$ , and of the partial derivatives – with the covariant derivatives of the connection  $\Gamma^\lambda{}_{\mu\nu}$ .

The total action reads

$$S_{ECT} = \int d^4x \sqrt{g} \left\{ \frac{1}{2\kappa c} R(\Gamma) + \frac{1}{c} \mathcal{L}_m(\Phi^A, \nabla_\mu \Phi^A, g_{\mu\nu}) + \frac{1}{2c} \Lambda^{\lambda\mu\nu} \nabla_\lambda g_{\mu\nu} \right\}, \quad (1.34)$$

where  $\Phi^A$  are the matter field variables ( $A$  is the generalized index),  $\Lambda^{\lambda\mu\nu}$  are the Lagrange multipliers. The matter Lagrangian depends on the connection  $\Gamma^\alpha{}_{\beta\mu}$  via the covariant derivatives only

$$\nabla_\mu \Phi^A = \partial_\mu \Phi^A - \Gamma^\alpha{}_{\beta\mu} \Omega_\alpha{}^{\beta A}{}_B \Phi^B. \quad (1.35)$$

Here  $\Omega_\alpha{}^{\beta A}{}_B$  are the generators of the general coordinate transformations:

$$x^\mu \longrightarrow x^\mu + \xi^\mu(x), \quad \Phi^A \longrightarrow \Phi^A - (\partial_\beta \xi^\alpha) \Omega_\alpha{}^{\beta A}{}_B \Phi^B.$$

By varying (1.34) with respect to  $g_{\mu\nu}$ ,  $\Gamma^\lambda{}_{\mu\nu}$ ,  $\Lambda^{\lambda\mu\nu}$ , and  $\Phi^A$ , we obtain the following field equations:

$$\frac{\delta S_{ECT}}{\delta g^{\mu\nu}} = \frac{1}{2c} \left\{ \frac{1}{\kappa} G_{(\mu\nu)}(\Gamma) + (\nabla_\lambda - 2Q_\lambda) \Lambda^\lambda{}_{\mu\nu} - T_{\mu\nu} \right\} = 0, \quad (1.36)$$

$$\frac{\delta S_{ECT}}{\delta \Gamma^\alpha{}_{\mu\nu}} = \frac{1}{c} \left\{ -\frac{1}{\kappa} (Q^\nu{}_{\alpha\beta} + 2\delta^\nu_{[\alpha} Q_{\beta]}) g^{\beta\mu} + N_\alpha{}^{\mu\nu} - \Lambda^{\nu\mu}{}_\alpha \right\} = 0, \quad (1.37)$$

$$\frac{\delta S_{ECT}}{\delta \Lambda^{\lambda\mu\nu}} = \frac{1}{2c} \nabla_\lambda g_{\mu\nu} = 0, \quad (1.38)$$

$$\frac{\delta S_{ECT}}{\delta \Phi^A} = \frac{1}{c} \left\{ \frac{\partial \mathcal{L}_m}{\partial \Phi^A} - (\nabla_\lambda - 2Q_\lambda) \frac{\partial \mathcal{L}_m}{\partial \nabla_\lambda \Phi^A} \right\} = 0, \quad (1.39)$$

where (1.36) and (1.37) are given in the final form after taking into account the metricity constraint (1.38). Here, in a usual way we defined

$$T^{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\partial(\sqrt{g}\mathcal{L}_m)}{\partial g_{\mu\nu}}$$

the metrical energy-momentum tensor of matter, and we denoted

$$N_\alpha{}^{\mu\nu} = \frac{\partial\mathcal{L}_m}{\partial\Gamma^\alpha{}_{\mu\nu}}.$$

Making use of the eq. (1.35), we find

$$N_\alpha{}^{\mu\nu} = \frac{\partial\mathcal{L}_m}{\partial\nabla_\lambda\Phi^A} \frac{\partial\nabla_\lambda\Phi^A}{\partial\Gamma^\alpha{}_{\mu\nu}} = -\frac{\partial\mathcal{L}_m}{\partial\nabla_\nu\Phi^A} \Omega_\alpha{}^{\mu A}{}_B\Phi^B. \quad (1.40)$$

Solution of (1.38) with respect to  $\Gamma^\alpha{}_{\mu\nu}$  yields the Riemann-Cartan connection (1.12). Lowering the index  $\mu$  in (1.37), we obtain the equation

$$-\frac{1}{\kappa} (Q^\nu{}_{\alpha\beta} + 2\delta^\nu_{[\alpha} Q_{\beta]}) + N_{\alpha\beta}{}^\nu - \Lambda^\nu{}_{\beta\alpha} = 0.$$

Let us decompose it into the symmetric and antisymmetric parts. From the symmetric one, we find the Lagrange multipliers explicitly

$$\Lambda^\nu{}_{\alpha\beta} = N_{(\alpha\beta)}{}^\nu,$$

so that (1.36) is recast into

$$G_{(\mu\nu)}(\Gamma) = \kappa \{T_{\mu\nu} - (\nabla_\lambda - 2Q_\lambda) N_{(\mu\nu)}{}^\lambda\}. \quad (1.41)$$

The remaining antisymmetric part of (1.37) reads as follows:

$$Q^\nu{}_{\alpha\beta} + 2\delta^\nu_{[\alpha} Q_{\beta]} = \kappa c S^\nu{}_{\alpha\beta}, \quad (1.42)$$

where

$$cS^\nu{}_{\alpha\beta} = N_{[\alpha\beta]}{}^\nu = -\frac{\partial\mathcal{L}_m}{\partial\nabla_\nu\Phi^A} \Omega_{[\alpha\beta]}{}^A{}_B\Phi^B$$

coincides with the canonical tensor of the spin density arising from the Noether theorem.

For the matter whose Lagrangian does not depend on the connection (for the scalar fields, for example), or for the vacuum gravitational field, the equations (1.42) yield  $Q^\lambda{}_{\mu\nu} = 0$  and then (1.41) reduce to the usual Einstein's equations.

Otherwise, the torsion is non-zero and (1.41) differ from the GR equations. One can show that the term  $(\nabla_\lambda - 2Q_\lambda) N_{(\mu\nu)}{}^\lambda$  leads to the fact that the sources of the gravitational field are the canonical tensor of spin  $S^\lambda{}_{\mu\nu}$ , and the symmetrical part of the canonical tensor of the energy-momentum of matter

$$t^\mu{}_\nu := \delta^\mu_\nu \mathcal{L}_m - \frac{\partial\mathcal{L}_m}{\partial\nabla_\mu\Phi^A} \nabla_\nu\Phi^A.$$

Indeed, let us consider an infinitesimal transformation of local coordinates  $x^\mu \rightarrow \bar{x}^\mu = x^\mu + \xi^\mu \delta\tau$ , where  $\xi^\mu(x^\lambda)$  is an arbitrary smooth vector field, and  $\delta\tau$  is an infinitesimal parameter. The Lie derivative of the Lagrangian density  $\sqrt{g}\mathcal{L}_m(\Phi^A, \nabla_\mu\Phi^A, g_{\mu\nu})$  along the vector field  $\xi^\mu$ , which induces these transformations, is computed as follows:

$$L_\xi(\sqrt{g}\mathcal{L}_m) = \frac{\partial(\sqrt{g}\mathcal{L}_m)}{\partial\Phi^A}L_\xi\Phi^A + \frac{\partial(\sqrt{g}\mathcal{L}_m)}{\partial\nabla_\mu\Phi^A}L_\xi\nabla_\mu\Phi^A + \frac{\partial(\sqrt{g}\mathcal{L}_m)}{\partial g_{\mu\nu}}L_\xi g_{\mu\nu}. \quad (1.43)$$

Using the definition of Lie derivative (which is introduced on the spacetime manifold even in the absence of the metrics and connection [3]), it is straightforward to see that in the Riemann-Cartan space

$$L_\xi\Phi^A = \xi^\lambda\nabla_\lambda\Phi^A + \zeta_\beta^\alpha\Omega_\alpha^{\beta A}{}_B\Phi^B, \quad (1.43a)$$

$$L_\xi\nabla_\mu\Phi^A = \xi^\lambda\nabla_\lambda\nabla_\mu\Phi^A + \zeta_\beta^\alpha\Omega_\alpha^{\beta A}{}_B\nabla_\mu\Phi^B + \zeta_\mu^\nu\nabla_\nu\Phi^A, \quad (1.43b)$$

$$L_\xi g_{\mu\nu} = 2\zeta_{(\mu\nu)}, \quad (1.43c)$$

where  $\zeta_\mu^\nu = \nabla_\mu\xi^\nu + 2Q_\nu^{\mu\lambda}\xi^\lambda$ .

On the other hand, since  $(\sqrt{g}\mathcal{L}_m)$  is a scalar density of the weight +1, we have

$$L_\xi(\sqrt{g}\mathcal{L}_m) = \xi^\lambda\nabla_\lambda(\sqrt{g}\mathcal{L}_m) + \zeta_\mu^\mu(\sqrt{g}\mathcal{L}_m). \quad (1.43d)$$

Substituting (1.43a)-(1.43c) into (1.43) and comparing with (1.43d), we find

$$-t^\mu{}_\nu + T^\mu{}_\nu - (\nabla_\lambda - 2Q_\lambda)N_\nu^{\mu\lambda} + c\frac{\delta S_{ECT}}{\delta\Phi^A}\Omega_\nu^{\mu A}{}_B\Phi^B = 0, \quad (1.44)$$

$$(\nabla_\mu - 2Q_\mu)t^\mu{}_\nu + 2Q^\lambda{}_{\nu\mu}t^\mu{}_\lambda + cS^\mu{}_{\alpha\beta}R^{\alpha\beta}{}_{\mu\nu}(\Gamma) = c\frac{\delta S_{ECT}}{\delta\Phi^A}\nabla_\nu\Phi^A. \quad (1.45)$$

By virtue of the field equations (1.39), we get from (1.44)

$$t_{(\mu\nu)} = T_{\mu\nu} - (\nabla_\lambda - 2Q_\lambda)N_{(\mu\nu)}{}^\lambda, \quad (1.46)$$

$$t_{[\mu\nu]} = c(\nabla_\lambda - 2Q_\lambda)S^\lambda{}_{\mu\nu}. \quad (1.47)$$

Thus we see that eq. (1.46) establishes the relation between the symmetrical part of the canonical and the metrical tensors of energy-momentum of matter. Making use of this relation in (1.41), we find the complete system of the field equations of the Einstein-Cartan theory:

$$G_{\mu\nu}(\Gamma) = \kappa t_{\mu\nu}, \quad (1.48)$$

$$Q^\lambda{}_{\mu\nu} + \delta_\mu^\lambda Q_\nu - \delta_\nu^\lambda Q_\mu = \kappa c S^\lambda{}_{\mu\nu}. \quad (1.49)$$

It is worthwhile to note that to determine the dynamics of the metric it is sufficient to solve only the symmetric part of (1.48) which is equivalent to (1.41). The skew-symmetric part of (1.48) is redundant and is automatically satisfied in view of (1.49) and the contracted Ricci identity.

Summarizing, within the framework of the Einstein-Cartan theory, the matter source of gravitational field  $(g, \Gamma)$  is described by the canonical energy-momentum tensor  $t^\lambda{}_\mu$  and the spin density tensor  $S^\lambda{}_{\mu\nu}$ . The invariance of the theory with respect to the general coordinate transformations and the local Lorentz group leads to the conservation law of energy-momentum (1.45) and of the total angular momentum (1.47).

The important feature of the Einstein-Cartan theory (ECT) is the algebraic nature of the equation (1.49), which will be called a Palatini equation (sometimes it is also called a Cartan equation). As a consequence of such an algebraic coupling, the torsion disappears in the absence of a spinning matter, which confirms Cartan's prediction [19]. Furthermore, the entire Einstein-Cartan theory can be written in terms of the Riemann-Einstein objects. Indeed, using the expression for the connection (1.12), we can explicitly separate the Riemann-Einstein part in the action

$$S_{ECT} = \frac{1}{2\kappa c} \int d^4x \sqrt{g} R(\Gamma) = \frac{1}{2\kappa c} \int d^4x \sqrt{g} (R + 2T^\alpha{}_{\mu[\alpha} T^{\mu\beta}{}_{\beta]}) \quad (1.50)$$

plus the surface integral.

Next, we solve eq. (1.49) expressing the torsion in terms of the spin

$$Q^\lambda{}_{\mu\nu} = \kappa c \left( S^\lambda{}_{\mu\nu} + \frac{1}{2} \delta_\mu^\lambda S_\nu - \frac{1}{2} \delta_\nu^\lambda S_\mu \right),$$

with  $S_\mu = S^\lambda{}_{\mu\lambda}$ . Inserting this into (1.50), we obtain the action

$$S_{ECT} = \int d^4x \sqrt{g} \left( \frac{1}{2\kappa c} R + \frac{1}{c} \mathcal{L}_m^{\text{eff}} \right), \quad (1.51)$$

where the effective matter Lagrangian reads

$$\mathcal{L}_m^{\text{eff}} = \mathcal{L}_m - \kappa c^2 \left[ S^\mu S_\mu + \frac{1}{2} S^{\mu\nu\lambda} (S_{\nu\lambda\mu} + S_{\lambda\mu\nu} + S_{\mu\lambda\nu}) \right].$$

The field equations (1.48) and (1.49) are then recast into

$$G_{\mu\nu} = \kappa T_{\mu\nu}^{\text{eff}}, \quad (1.52)$$

where the effective metrical tensor of the energy-momentum of matter

$$T_{\text{eff}}^{\mu\nu} = \frac{2}{\sqrt{g}} \frac{\delta(\sqrt{g} \mathcal{L}_m^{\text{eff}})}{\delta g_{\mu\nu}}$$

satisfies the covariant conservation law,  $T_{\text{eff}}^{\mu\nu}{}_{;\nu} = 0$ , of GR [188].

Thus, the spin-torsion interaction in the Einstein-Cartan theory predicts the appearance of the (effective) matter Lagrangian of the terms describing the contact spin-spin interaction, which is  $\kappa c^2$  times smaller than the interaction between the energy-momentum and the metric in GR. The change of the matter current  $T_{\mu\nu} \rightarrow T_{\mu\nu}^{\text{eff}}$  affects the spacetime metric via the effective Einstein equation (1.52). In those areas, where the spinning matter is absent, the influence of the spin-spin interaction manifests itself through the matching conditions for the spin discontinuity at the boundary [21].

# 2

## Canonical formalism of gravity theories

### 2.1. Systems with singular Lagrangians

The gravitational field theory is invariant with respect to the group of general coordinate transformations, as a result, the dynamics of this field is essentially hidden due to the arbitrary choice of a coordinate system. The dynamical contents of any theory as such (i.e., the calculation of the number of independent degrees of freedom, their explicit selection from the full set of the field variables, the construction of such dynamical characteristics as the energy, etc.) can be revealed by the methods of the canonical formalism. The solution of the problems listed above is necessary for the canonical quantization of the theory, which leads to the unitary  $S$ -matrix.

Therefore, here and in the next chapter, we will study the specific features of the Hamiltonian gravitational dynamics. We begin with considering of the dynamical consequences of the invariance of the theory with respect to the local group.

*Degeneracy of the dynamical system as a result of invariance with respect to the local group*

Let us find out what are the dynamical consequences of the invariance of an action of the field system  $\Phi^A(x)$  ( $A = 1, \dots, k$ )

$$S[\Phi] = \int d^4x \mathcal{L}[\Phi^A(x), \partial_\mu \Phi^A(x)], \quad (2.1)$$

with respect to transformations of the field variables and of the spacetime coordinates, constituting the local group:

$$\Phi^A(x) \longrightarrow \Phi'^A(x'), \quad x^\mu \longrightarrow x'^\mu(x).$$

If

$$\Delta^f \Phi^A(x) \equiv \Phi'^A(x) - \Phi^A(x), \quad (2.2)$$

$$\Delta^f x^\mu(x) \equiv x'^\mu - x^\mu \quad (2.3)$$

are the infinitesimal transformations of this  $m$ -parameter local group with the local infinitesimal parameters  $f^\alpha(x)$  ( $\alpha = 1, \dots, m$ ), then the condition of the invariance of the action (2.1) reads

$$\int d^4x \left\{ \left( \frac{\partial \mathcal{L}}{\partial \Phi^A} - \partial_\mu \frac{\partial \mathcal{L}}{\partial \partial_\mu \Phi^A} \right) \Delta^f \Phi^A + \partial_\mu \left( \mathcal{L} \Delta^f x^\mu + \frac{\partial \mathcal{L}}{\partial \partial_\mu \Phi^A} \Delta^f \Phi^A \right) \right\} = 0. \quad (2.4)$$

Let us consider such transformation parameters that the variations (2.2) and (2.3) vanish outside some four-dimensional spacetime area  $\Omega$ , as well as its boundary. After the integration in (2.4), we then obtain

$$\int_\Omega d^4x \left( \frac{\partial \mathcal{L}}{\partial \Phi^A} - \partial_\mu \frac{\partial \mathcal{L}}{\partial \partial_\mu \Phi^A} \right) \Delta^f \Phi^A = 0. \quad (2.5)$$

It is convenient to introduce the new condensed notation of the field variables  $\Phi^A(\mathbf{x}) \equiv Q^i(t)$ , where the dependence on time  $t$  is explicitly distinguished, and the generalized index  $i = (A, \mathbf{x})$  labels both the discrete internal components and the spatial coordinates. The summation over  $i$  thus will also imply an integration over  $\mathbf{x} = \{x^1, x^2, x^3\}$ .

We assume that the group transformations are arbitrarily complicated local functions of the fields, of the group parameters  $f^\alpha$ , and of their spacetime derivatives up to the  $N$ -th order. The parameters of the group  $f^\alpha$  will also be labeled by the condensed indices  $\alpha$ , including the spatial coordinates.

Then we can recast (2.2) into

$$\Delta^f Q^i(t) = \sum_{n=0}^N a_\alpha^{(n)i} \partial_0^n f^\alpha(t), \quad \partial_0 \equiv \frac{\partial}{\partial t}, \quad (2.6)$$

where  $a_\alpha^{(n)i}$  are local in time functions of the fields ( $n$  denotes the order of time derivative in the transformation), and summation over  $\alpha$  in field systems also implies integration over  $\mathbf{x}$ .

Substituting (2.6) into (2.5) and integrating by parts, we obtain

$$\int dt f^\alpha(t) \sum_{n=0}^N (-1)^n \partial_0^n \left\{ \left( \frac{\partial \mathcal{L}}{\partial Q^i(t)} - \partial_0 \frac{\partial \mathcal{L}}{\partial \dot{Q}^i(t)} \right) a_\alpha^{(n)i} \right\} = 0,$$

where  $\dot{Q}^i \equiv \partial Q^i / \partial t$ , and  $L = \int d^3x \mathcal{L}$  is system's Lagrange function. Due to arbitrariness of  $f^\alpha(t)$ , this yields the following identity:

$$\sum_{n=0}^N (-1)^n \partial_0^n \left\{ \left( \frac{\partial L}{\partial Q^i(t)} - \partial_0 \frac{\partial L}{\partial \dot{Q}^i(t)} \right) \binom{n}{a}^i_\alpha \right\} = 0. \quad (2.7)$$

We will consider the second-order theories, then the following relation is fulfilled

$$\frac{\partial L}{\partial Q^i} - \partial_0 \frac{\partial L}{\partial \dot{Q}^i} = - \frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} \ddot{Q}^k + \mathcal{F}_i(Q^k, \dot{Q}^k), \quad (2.8)$$

where the functions  $\mathcal{F}(Q^k, \dot{Q}^k)$  do not depend on the second derivatives  $\ddot{Q}^i(t)$ .

Using (2.8) in (2.7), we obtain the identity that is valid for any  $Q^i(t)$ :

$$\binom{N}{a}^i_\alpha \frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} \partial_0^{N+2} Q^k + W_\alpha(Q, \partial_0 Q, \dots, \partial_0^{N+1} Q) = 0, \quad (2.9)$$

where the functions  $W_\alpha$  do not depend on the higher derivatives  $\partial_0^{N+2} Q^i(t)$ . Due to arbitrariness of  $Q^i(t)$ , both terms in (2.9) are identically zero, since the derivatives of the  $(N+2)$ -th order do not depend on the lower derivatives.

Therefore,  $\binom{N}{a}^i_\alpha \frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} = 0$  and the matrix of the second derivatives of the Lagrangian with respect to the velocities is degenerate:

$$\det \frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} = 0, \quad \text{rank} \left( \frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} \right) = k - m. \quad (2.10)$$

The Lagrangians with the property (2.10) are called singular, and the related systems are called degenerate. The equations of motion for such systems

$$\frac{\partial^2 L}{\partial \dot{Q}^i \partial \dot{Q}^k} \ddot{Q}^k - \mathcal{F}_i(Q, \dot{Q}) = 0 \quad (2.11)$$

cannot be solved with respect to the higher derivatives, and  $m$  linear combinations of these equations, containing only the coordinates  $Q^i$  and their velocities  $\dot{Q}^i$ , vanish on the extremals of the degenerate action (2.1):

$$\binom{N}{a}^i_\alpha \mathcal{F}_i(Q, \dot{Q}) = 0. \quad (2.12)$$

The latter means that the initial data for the coordinates and the velocities cannot be fixed independently.

Before we consider the Cauchy problem and analyse how to solve the equation (2.11) of an arbitrary degenerate system, it is instructive to illustrate our derivations by the two examples of systems with the singular Lagrangians: the relativistic particle, and Maxwell's electrodynamics.

The action  $S = m \int dt \sqrt{-\dot{x}^\mu \dot{x}_\mu}$  of a relativistic particle  $\{Q^i(t) \equiv x^\mu(t)\}$  of the mass  $m$  is invariant with respect to the transformations

$$\Delta t = f(t), \quad \Delta x^\mu = \overset{(0)}{a}{}^\mu f, \quad \overset{(0)}{a}{}^\mu = x^\mu(t),$$

and its Lagrangian  $L = m \sqrt{-\dot{x}^\mu \dot{x}_\mu}$  satisfies the relation  $\dot{x}^\mu \partial^2 L / \partial \dot{x}^\mu \partial \dot{x}^\nu = 0$ . Interesting property of the relativistic particle action is the vanishing of its Hamiltonian,  $H = \dot{x}^\mu (\partial L / \partial \dot{x}^\mu) - L \equiv 0$ .

The electromagnetic field action with  $(Q^i \equiv A_\mu(t, \mathbf{x}))$ ,  $S = -\frac{1}{4} \int d^4x F_{\mu\nu} F^{\mu\nu}$ ,  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ , is invariant with respect to the Abelian one-parameter group  $\Delta^f A_\mu = \partial_\mu f(x)$ , so that

$$\overset{(1)}{a}{}_\mu(\mathbf{x}, \mathbf{x}') = \delta_\mu^0 \delta(\mathbf{x} - \mathbf{x}'), \quad (2.13)$$

and the degeneracy condition (2.10) of the system reads

$$\int d^3\mathbf{x}' \overset{(1)}{a}{}_\mu(\mathbf{x}, \mathbf{x}') \frac{\partial^2 L}{\partial \dot{A}_\mu(\mathbf{x}') \partial \dot{A}_\nu(\mathbf{y})} = \frac{\partial^2 L}{\partial \dot{A}_0(\mathbf{x}) \partial \dot{A}_\nu(\mathbf{y})} = 0.$$

### *Canonical formalism for systems with singular Lagrangians*

Thus far, it is shown that the invariance of a dynamical system with respect to a local group leads to its degeneracy. The dynamics of the system and its Cauchy problem are usually studied within the framework of the canonical formalism, so let us consider the specific features of the canonical formalism for systems with singular Lagrangians.

Define the momenta conjugated to the canonical coordinates  $Q^i$ :

$$\mathcal{P}_i = \frac{\partial L}{\partial \dot{Q}^i}. \quad (2.14)$$

Since the matrix of the second derivative of the Lagrangian with respect to the velocities is degenerate (2.10), the equations (2.14) cannot be solved with respect to the velocities in terms of the momenta. Technically, this means that not all  $k$  momenta  $\mathcal{P}_i$  can be taken as independent arguments. Looking at the rank of the matrix  $\partial^2 L / \partial \dot{Q}^i \partial \dot{Q}^k$ , we conclude that there are  $m$  identities connecting momenta and coordinates,

$$\varphi_\mu(Q, \mathcal{P}) = 0. \quad (2.15)$$

These identities are called the primary constraints [40]. Now let us consider the function of the three sets of arguments  $Q^i$ ,  $\dot{Q}^i$  and  $\mathcal{P}_i$ :

$$H_0 = \mathcal{P}_i \dot{Q}^i - L(Q^i, \dot{Q}^i). \quad (2.16)$$

Let us calculate the variation of this quantity with all the arguments treated as independent ones, and then put it in the space of the variables  $(Q, \dot{Q}, \mathcal{P})$  on

the surface defined by the equation (2.14):

$$\delta H_0 \Big|_{\mathcal{P}_i = \partial L / \partial \dot{Q}^i} = \delta \mathcal{P}_i \dot{Q}^i - \frac{\partial L}{\partial Q^i} \delta Q^i.$$

It is obvious that even in spite of the absence of the functional independence of the arguments  $Q$  and  $\mathcal{P}$ , one can express the quantity (2.16) as a function of coordinates and momenta only, which on the surface (2.14) does not depend on the velocities  $\dot{Q}^i$ :

$$H_0(Q, \mathcal{P}) = \left\{ \mathcal{P}_i \dot{Q}^i - L(Q, \dot{Q}) \right\} \Big|_{\mathcal{P}_i = \partial L / \partial \dot{Q}^i}. \quad (2.17)$$

We now develop the first-order variational formalism with the Hamiltonian (2.17). For non-degenerate theories with the independent coordinates and momenta, this formalism is based on the variation with respect to  $Q$  and  $\mathcal{P}$  of the action

$$S[Q, \mathcal{P}] = \int dt \left\{ \mathcal{P}_i \dot{Q}^i - H_0(Q, \mathcal{P}) \right\}.$$

In the theories with singular Lagrangians, it is necessary to study a conditional extremum of the same functional subject to the conditions defined by the constraints (2.15). This is achieved by the inclusion of the constraints with the Lagrange multipliers  $\lambda^\mu$  into the action functional,

$$S[Q, \mathcal{P}, \lambda] = \int dt \left\{ \mathcal{P}_i \dot{Q}^i - H_0(Q, \mathcal{P}) - \lambda^\mu \varphi_\mu(Q, \mathcal{P}) \right\}. \quad (2.18)$$

The equations, obtained by the variation of this action, read as follows:

$$\begin{aligned} \dot{\mathcal{P}}_i &= \{ \mathcal{P}_i, H_0 \} + \lambda^\mu \{ \mathcal{P}_i, \varphi_\mu \}, \\ \dot{Q}^i &= \{ Q^i, H_0 \} + \lambda^\mu \{ Q^i, \varphi_\mu \}, \\ \varphi_\mu &= 0, \end{aligned} \quad (2.19)$$

where  $\{ , \}$  is the Poisson bracket determined by the phase variables  $(Q, \mathcal{P})$ .

However, it is unclear that the total Hamiltonian in the action (2.18) is written correctly, i.e., that all the occurring constraints are taken into account. Indeed, the condition of consistency of the constraints (2.15) with the equations of motion (2.19) means that

$$\dot{\varphi}_\mu = 0, \quad \{ \varphi_\mu, H_0 \} + \lambda^\alpha \{ \varphi_\mu, \varphi_\alpha \} = 0. \quad (2.20)$$

Demanding that this equation is satisfied, we have one the following options:

1.  $\det \{ \varphi_\mu, \varphi_\alpha \} \neq 0$ . Then the equations (2.20) explicitly determine the Lagrange multipliers, and substituting the latter into (2.19) we get the system of equations that have a unique solution. The time dependence of the field variables  $Q^i(t)$  is then uniquely defined.

2. Among the set (2.15), there are such constraints  $\varphi_q$  ( $q = 1, \dots, r \leq m$ ) that  $\{\varphi_q, \varphi_\alpha\} = C_{q\alpha}^\mu \varphi_\mu$ , where  $C_{q\alpha}^\mu$  are some functions of the phase coordinates. Then  $\det \{\varphi_\mu, \varphi_\alpha\}_{\varphi=0} = 0$  and the equations (2.20) yield new  $r$  equations

$$\chi_q = 0, \quad \chi_q(Q, \mathcal{P}) \equiv \{\varphi_q, H_0\},$$

which are called the secondary constraints. Since the phase coordinates must satisfy these new relations, it is necessary to add them to the action with the new Lagrange multipliers. Repeating the process, we again obtain one of the two aforementioned options. Such a sequence of steps can be completed only as follows. We denote the complete set of  $N$  constraints obtained during this process by

$$\Psi_\Phi = 0, \quad \Psi_\Phi = (H_\alpha, \Xi_A), \quad (2.21)$$

where  $R$  constraints  $H_\alpha$  satisfy the relations

$$\{H_\alpha, \Psi_A\} = C_{\alpha A}^{\Phi} \Psi_\Phi, \quad \alpha = 1, \dots, R, \quad (2.22)$$

and the remaining constraints  $\Xi_A$  are numbered by the index  $A = R+1, \dots, N$ . Then

$$\{H_\alpha, H_0\} = C_\alpha^{\Phi} \Psi_\Phi, \quad (2.23)$$

otherwise the consistency conditions of the form (2.20) would yield new constraints, and the procedure would not be completed.

The constraints that satisfy (2.21)-(2.23) are called the constraints of the first kind. They commute in the sense of the Poisson brackets on the surface of constraints in the phase space with each other and with the Hamiltonian  $H_0$ . One can show that the remaining constraints  $\Xi_A$  satisfy

$$\det \{\Xi_A, \Xi_B\} \neq 0. \quad (2.24)$$

These are the constraints of the second kind.

The total canonical action takes the following form:

$$S[Q, \mathcal{P}, \lambda] = \int dt \left\{ \mathcal{P}_i \dot{Q}^i - H_0 - \lambda^\alpha H_\alpha - \lambda^A \Xi_A \right\}. \quad (2.25)$$

The condition (2.24) provides the uniqueness of the solution for  $\lambda^A$  (in terms of the remaining variables  $Q, \mathcal{P}, \lambda^\alpha$ ), while the Lagrange multipliers  $\lambda^\alpha$  corresponding to the constraints of the first kind  $H_\alpha$  remain completely arbitrary. This manifests the ambiguity in the solving of the equations of motion of a degenerate system. The origin of this ambiguity lies in the invariance of the theory with respect to the action of the local group, which leads to system's degeneracy. Later we will demonstrate that variations of the undetermined Lagrange multipliers  $\lambda^\alpha$  correspond to the local group transformations, and that the constraints  $H_\alpha$  are the generators of these transformations in the phase space.

We have described the general algorithm of construction of a complete system of constraints for the case of an arbitrary degenerate theory. Let us now specialize to the theories without constraints of the second kind, and give the form of canonical action (2.25) for this important particular case. We assume that by an appropriate choice of the field variables, we can recast the primary couplings (2.15) into a simple form

$$\varphi_\mu = \mathcal{P}_\mu, \quad \mathcal{P}_\mu = 0. \quad (2.26)$$

This means that the velocities  $\dot{Q}^\mu$  do not appear in the Lagrangian at all. Let us then exclude  $(Q^\mu, \mathcal{P}_\mu)$  from the complete set  $(Q, \mathcal{P}) = (Q^\mu, \mathcal{P}_\mu; q^i, p_i)$ ,  $i = 1, \dots, n$ . Since the Lagrangian does not depend on  $\dot{Q}^\mu$ , the variables  $Q^\mu$  play the role of the Lagrange multipliers,  $Q^\mu = \lambda^\mu$ , and the canonical action is written as<sup>1</sup>

$$S[q, p, \lambda] = \int_{t_0}^{t_1} dt \{p_i \dot{q}^i - H_0(q, p) - \lambda^\mu H_\mu(q, p)\}, \quad (2.27)$$

where for definiteness we introduced the time parameters  $t_0$  and  $t_1$  which determine the initial and the final Cauchy hypersurfaces.

The conservation of the initial constraints (2.26) is a condition of stationarity of the action (2.26) with respect to the variation of  $\lambda^\mu$ , or

$$H_\mu(q, p) = 0. \quad (2.28)$$

Let us consider degenerate systems with the canonical action of the form (2.27), where  $H_\mu(q, p)$  are the constraints of the first kind, that satisfy the so-called involution relations:

$$\{H_\mu, H_\nu\} = U_{\mu\nu}^\alpha H_\alpha, \quad \{H_\mu, H_0\} = V_\mu^\nu H_\nu. \quad (2.29)$$

Here  $U_{\mu\nu}^\alpha$  and  $V_\mu^\nu$  are some quantities depending generally on the phase variables, and the Poisson brackets are taken in terms of the phase variables of the reduced phase space  $(q, p)$ .

The equations of motion, obtained from the action (2.27) for the phase variables  $\Phi(q, p)$ , read as follows:

$$\dot{\Phi} = \{\Phi, H_0\} + \lambda^\mu \{\Phi, H_\mu\}. \quad (2.28a)$$

Varying the Lagrange multipliers arbitrarily, one can obtain different solutions  $\Phi(t)$  even for the same initial conditions. This ambiguity corresponds to an

---

<sup>1</sup>The linear dependence of the complete Hamiltonian  $H = H_0 + \lambda T$  on  $\lambda$  is explained by the relation  $\text{rank } \partial^2 H / (\partial \lambda^\mu \partial \lambda^\nu) = 0$ , the violation of which would mean that it is possible to exclude part of  $\lambda^\mu$  in terms of the other variables, in contradiction with the number  $m$  of the constraints.

uncertainty with which one can fix the state of a physical system, which is invariant with respect to the action of a local group. If  $Q(t) = (\lambda^\mu(t), q^i(t))$  is a solution of the equations of motion of such system, then in view of (2.2), (2.3) and (2.6), the following functions will satisfy the same equations

$$\begin{aligned}\lambda'^\mu(t) &= \lambda^\mu(t) + \Delta^f \lambda^\mu(t), \\ q'^i(t) &= q^i(t) + \Delta^f q^i(t),\end{aligned}\tag{2.29}$$

where the infinitesimal transformations of the group are defined by the formulas (2.6) with some parameters  $f^\alpha$ . Thus, the physical state in the phase space and in the  $\lambda$ -multipliers' space at a moment  $t$  is determined up to a group transformation (2.29). The functions  $Q(t)$ , as such, do not have a direct physical meaning. Only the equivalence classes of the field configurations connected by group transformations (2.2)-(2.3) or the group invariants have physical meaning.

One can verify that the canonical action (2.27) is invariant with respect to transformations of the phase coordinates and  $\lambda$ -multipliers of the following form:

$$\delta^{\mathcal{F}} q^i = \{q^i, H_\mu\} \mathcal{F}^\mu, \tag{2.30}$$

$$\delta^{\mathcal{F}} p_i = \{p_i, H_\mu\} \mathcal{F}^\mu, \tag{2.30a}$$

$$\delta^{\mathcal{F}} \lambda^\mu = \dot{\mathcal{F}}^\mu - U_{\alpha\beta}^\mu \lambda^\alpha \mathcal{F}^\beta - V_\alpha^\mu \mathcal{F}^\alpha, \tag{2.31}$$

with some infinitesimal parameters  $\mathcal{F}^\mu$ . It turns out that there is a one-to-one correspondence between the parameters  $\mathcal{F}^\mu$  and  $f^\alpha$ , so that the transformations (2.30), (2.31) are the realizations of the group transformations (2.29) in the configuration space, under the condition that the transformations (2.30) and (2.31) are calculated on the extremal values of the momenta  $p_i^0 = p_i(q, \dot{q}, \lambda)$ . The extremal values of the momenta are obtained as a result of solving the equations

$$\dot{q}^i = \frac{\partial H_0}{\partial p_i} + \lambda^\mu \frac{\partial H_\mu}{\partial p_i}$$

with respect to  $p_i$  in terms of variables  $(q, \dot{q}, \lambda)$ . Therefore, the constraints of the first kind  $H_\mu(q, p)$  are generators of the phase transformations corresponding to the local group transformations.

Since the dynamics of the physical state is determined not by a particular set of the functions  $(q, p)$ , but by the entire equivalence class of the phase coordinates connected by the transformations (2.30) and (2.30a), we choose the evolution of one representative from this class to describe the dynamics. This is done by means of imposing the additional conditions

$$\chi^\alpha(q, p) = 0, \tag{2.32}$$

which break the invariance with respect to the transformations (2.30), (2.30a). Obviously, the number of gauge conditions should be equal to the number of constraints of the first kind or the number of parameters  $m$  of the local group,

and the requirement that no nonzero  $\mathcal{F}^\beta$  would leave the gauge conditions (2.32) invariant,  $\delta^{\mathcal{F}}\chi^\alpha = \{\chi^\alpha, H_\beta\}\mathcal{F}^\beta = 0$ , with respect to the transformations (2.30)-(2.30a), implies the invertibility of the following *Faddeev-Popov* matrix:

$$\det J_\beta^\alpha \neq 0, \quad J_\beta^\alpha = \{\chi^\alpha, H_\beta\}. \quad (2.33)$$

Following the choice of a gauge, the system becomes non-degenerate and is described by the action

$$S[q, p, \xi] = \int dt (p_i \dot{q}^i - H_0 - \xi_a \psi^a), \quad (2.33a)$$

where the complete set of constraints and the corresponding Lagrange multipliers are defined as

$$\psi^a = (H_\alpha, \chi^\beta), \quad \xi_a = (\lambda^\alpha, \pi_\beta).$$

The system of constraints  $\psi^a$  is of the second kind, because due to (2.33), the matrix

$$Q^{ab} = \{\psi^a, \psi^b\}, \quad \det Q^{ab} = -(\det \{\chi^\alpha, H_\beta\})^2 \quad (2.34)$$

is non-degenerate. The Lagrange multipliers can be found from the condition of conservation of the constraints in time

$$\frac{d\psi^a}{dt} = \frac{\partial\psi^a}{\partial t} + \{\psi^a, H_0\} + \xi_b \{\psi^a, \psi^b\} = 0,$$

which yields

$$\xi_a = \{\psi^b, H_0\} Q_{ba}^{-1} + \frac{\partial\psi^b}{\partial t} Q_{ba}^{-1}, \quad (2.35)$$

where  $Q_{ba}^{-1}$  is the inverse matrix to (2.34). In general case, the additional conditions (2.32) may explicitly depend on time. Hence, there is a  $\partial\psi^a/\partial t$  contribution in (2.35). With an account of (2.35), the equations of motion for the phase variables  $\Phi$  take on the form:

$$\dot{\Phi} = \{\Phi, H_0\}_D - \{\Phi, \psi^a\} Q_{ab}^{-1} \frac{\partial\psi^b}{\partial t}, \quad (2.36)$$

where  $\{, \}_D$  is the so-called Dirac bracket [40] defined as

$$\{A, B\}_D = \{A, B\} - \{A, \psi^a\} Q_{ab}^{-1} \{\psi^b, B\}. \quad (2.37)$$

Since the evolution of the physical system is subject to the set of  $2m$  constraints (2.28) and (2.32), the number of independent (physical) degrees of freedom equals  $(2n - 2m)/2 = n - m$ . These degrees of freedom  $\Phi^* = (q^{*A}, p_A^*)$ ,  $A = 1, \dots, n - m$  parameterize the initial phase coordinates  $\Phi = (q^i, p_i)$  on the subspace of the full set of constraints

$$\Phi = \Phi(\Phi^*) : \quad q^i = q^i(q^*, p^*), \quad p_i = p_i(q^*, p^*), \quad \Psi^a(\Phi(\Phi^*)) \equiv 0, \quad (2.38)$$

so that the following relation holds

$$\int dt (p_i \dot{q}^i - H_0(q, p)) \Big|_{\Phi=\Phi(\Phi^*)} = \int dt (p_A^* \dot{q}^{*A} - H_{\text{phys}}(q^*, p^*)). \quad (2.39)$$

This relation may be interpreted as a definition of the physical Hamiltonian  $H_{\text{phys}}(q^*, p^*)$  and of the canonical action in terms of  $(n - m)$  physical degrees of freedom:

$$S[q^*, p^*] = \int dt (p_A^* \dot{q}^{*A} - H_{\text{phys}}(q^*, p^*)). \quad (2.40)$$

The equations obtained by varying of this action are equivalent to (2.36), since the action (2.40) is the same functional (2.33a), where the equations of the constraints are solved in terms of the smaller number of variables  $(q^*, p^*)$ . Therefore, each gauge choice (2.32) corresponds to a set of  $(n - m)$  physical degrees of freedom, represented by the canonical variables  $(q^*, p^*)$ . One can generate a change of the gauge  $\delta\chi^\alpha(q, p)$  by transformations (2.30), (2.30a) with the specially chosen parameters  $\mathcal{F}^\mu$  (see Chapter 7).

Let us illustrate the above theory on the examples of the relativistic particle and the theory of electromagnetic field. The momenta of the relativistic particle

$$p_\mu = \frac{\partial L}{\partial \dot{x}^\mu} = - \frac{mg_{\mu\nu} \dot{x}^\nu}{\sqrt{-\dot{x}_\alpha \dot{x}^\alpha}}$$

satisfy the constraint

$$H \equiv g^{\mu\nu} p_\mu p_\nu - m^2 = 0, \quad (2.41)$$

which due to the identical vanishing of Hamiltonian  $H_0$  is the constraint of the first kind. Let us choose the following gauge:

$$\chi \equiv x^0 - t = 0. \quad (2.42)$$

The condition (2.33) is fulfilled:  $\{H, \varphi\} = 2p^0 \neq 0$ . Using these quantities in (2.34) and (2.36), we obtain the equations of a straight-line motion of the free relativistic particle

$$\dot{x}^i = \frac{p^i}{p^0}, \quad \dot{p}_i = 0.$$

Taking into account the vanishing of the Hamiltonian  $H_0$ , the action (2.33a) for the relativistic particle can be written in the form

$$S[x^\mu, p_\mu, \lambda, \pi] = \int dt \{p_\mu \dot{x}^\mu - \lambda T - \pi \chi\}.$$

Choosing the spatial coordinates  $x^i = q^*$  and the momenta  $p_i = p^*$  as the physical degrees of freedom, and taking (2.42) into account, we arrive at the canonical action

$$S[x, p] = \int dt \left( p_i \dot{x}^i - \sqrt{m^2 + \mathbf{p}^2} \right)$$

with the Hamiltonian  $H_{\text{phys}} = \sqrt{m^2 + \mathbf{p}^2}$  obtained as a result of solving the constraint (2.41) with respect to the momentum  $p_0 = -H_{\text{phys}}$ .

For the electromagnetic field, the action (2.33a) can be written as

$$S[A_i, p^i, A_0] = \int dt d^3x \left\{ p^i \dot{A}_i - \left( \frac{1}{2} p_i p^i + \frac{1}{4} F_{ik} F^{ik} \right) - (-A_0) \partial_i p^i \right\},$$

where the momentum  $p^i$  conjugated to the vector potential  $A_1$  is equal to

$$p^i = \partial_0 A_i - \partial_i A_0,$$

and  $F_{ik} = \partial_i A_k - \partial_k A_i$ . The role of the phase variables is played by the set  $(A_i, p^i)$ , and the zeroth component of the vector potential  $-A_0 = \lambda$  is an undetermined Lagrange multiplier for the first-kind constraint  $H = \partial_i p^i$  commuting in the sense of the Poisson brackets with the Hamiltonian

$$H_0 = \int d^3x \left( \frac{1}{2} p_i p^i + \frac{1}{4} F_{ik} F^{ik} \right). \quad (2.43)$$

In accordance with the general procedure, we impose the Coulomb gauge

$$\chi = 0, \quad \chi \equiv \partial_i A^i. \quad (2.44)$$

The equation for the Lagrange multiplier  $A_0$  (2.35) reduces to  $\Delta A_0 = 0$  with  $\Delta = \partial_i \partial^i$ , hence we find  $A_0 = 0$  for the zero boundary conditions, and the equations of motion take the following form:  $\dot{A}_i = p^i$  and  $\dot{p}^i = -\Delta A_i$ , equivalent to Maxwell's equations in the gauge (2.44). The dynamically independent components of the vector potential  $q^{*A}$  ( $A = 1, 2$ ) are obtained by decomposing  $A_i(\mathbf{x}, t)$  into the two transversal polarizations  $A_i^{(A)}(\mathbf{x}, t)$  of the electromagnetic field satisfying (2.44).

### *The Hamilton-Jacobi theory of degenerate systems*

Let us consider some solution  $\Phi_0(t)$  (obtained in an arbitrary gauge) of the equations of motion (2.28a) and insert it into the action (2.27). The action (2.27) calculated on its own extremal  $\Phi_0(t)$  becomes the function of initial and final values of the phase coordinates  $q_0 \equiv q(t_0)$  and  $q_1 \equiv q(t_1)$ :

$$S[\Phi, \lambda] \Big|_{\Phi = \Phi_0(t)} = S(q_1, t_1 | q_0, t_0). \quad (2.45)$$

By denoting the current value of time  $t_1 = t$  and the phase coordinates  $q_1 = q$ , we treat the extremal action (2.45) as a function of these variables,  $S(t; q)$ . Using the variation procedure, one can show that this function is a potential of the field of the phase momenta  $p = p(t_1)$ :

$$p^i = \frac{\partial S}{\partial q_i}. \quad (2.43a)$$

Since the extremal momenta and the coordinates at any time satisfy the constraints (2.28), the  $m$  constraint equations are fulfilled for the function of the action  $S(t; q)$ :

$$H_\mu \left( q, \frac{\partial S}{\partial q} \right) = 0. \quad (2.44)$$

Similarly, from the variational procedure it follows that  $S(t; q)$  satisfies the Hamilton-Jacobi equation with the complete Hamiltonian  $H = H_0 + \lambda^\mu H_\mu$ . However, in view of (2.44) this equation reduces to

$$\frac{\partial S}{\partial t} + H_0 \left( q, \frac{\partial S}{\partial q} \right) = 0. \quad (2.45)$$

The equations (2.44) and (2.45) are the basic equations of the Hamilton-Jacobi theory of the systems with constraints. A solution of this system of equations, i.e., its complete integral, contains  $n-m$  parameters  $\alpha^A$  ( $A = 1, \dots, n-m$ ) [a solution of the equation (2.45) depends on  $n$  nontrivial integration constants, and the remaining  $m$  equations (2.44) impose  $m$  relations on them]. Thus, the function  $S$  depends on  $(n-m)$  additional parameters, the number of which is equal to the number of dynamically independent degrees of freedom  $q^*$ :

$$S = S(t; q, \alpha). \quad (2.46)$$

Such a number of parameters labeling the solutions of the system of equations (2.44), (2.45) does not contradict to the fact that the function  $S$  in (2.45) depends on the additional  $(n+1)$  quantities  $(q_0; t_0)$ , since the function (2.45) satisfies the same  $(m+1)$  equations (2.44), (2.45) with respect to the arguments  $(q_0; t_0)$  and the actual number of independent parameters reduces to  $(n+1) - (m+1) = n-m$ .

We will show that the solution of the equations of our system with constraints for some set of gauges  $\chi^\mu(q, p)$  can be found from the known action function  $S(t; q, \alpha)$  by solving the following system of equations:

$$\frac{\partial S}{\partial \alpha^A} = \beta_A, \quad (2.47)$$

$$\chi^\mu \left( q, \frac{\partial S}{\partial q} \right) = 0, \quad (2.48)$$

where  $\beta_A$  are some  $m$  arbitrary parameters.

Let us start with demonstrating that if the gauge  $\chi^\mu$  satisfies (2.33), the system of equations (2.47), (2.48) admits the unique solution  $q^i = q^i(t; \alpha, \beta)$ . Indeed, the solvability condition of this system means the absence of zero eigenvectors of the matrix  $B_{ki}$ ,  $k = (A, \mu)$ ,

$$B_{Ai} = \frac{\partial^2 S}{\partial \alpha^A \partial q^i}, \quad B_{\mu i} = \frac{\partial \chi^\mu}{\partial q^i} + \frac{\partial \chi^\mu}{\partial p^k} \bigg|_{p = \frac{\partial S}{\partial q}} \frac{\partial^2 S}{\partial q^k \partial q^i}.$$

Since the equations (2.44) are fulfilled identically for any  $\alpha^A$  and  $q^i$  if  $S$  is the solution (2.46), then by differentiating (2.44) with respect to  $\alpha^A$  and  $q^i$ , we obtain

$$\left. \frac{\partial H_\mu}{\partial p_i} \right|_{p = \frac{\partial S}{\partial q}} \frac{\partial^2 S}{\partial q^i \partial \alpha^A} = 0, \quad (2.49)$$

$$\left. \frac{\partial H_\mu}{\partial p_k} \right|_{p = \frac{\partial S}{\partial q}} \frac{\partial^2 S}{\partial q^k \partial q^i} = - \left. \frac{\partial H_\mu}{\partial q^i} \right|_{p = \frac{\partial S}{\partial q}}. \quad (2.50)$$

Hence,  $\partial H_\mu / \partial p_i|_{p = \partial S / \partial q}$  are the  $m$  zero vectors of the matrix  $\partial^2 S / \partial q^i \partial \alpha^A$ , and consequently their linear combination with non-zero coefficients  $\gamma^\mu$  is the zero vector of  $B_{ki}$ . However, in view of (2.50), we have

$$\gamma^\alpha \left. \frac{\partial H_\alpha}{\partial p_i} \right|_{p = \frac{\partial S}{\partial q}} B_{\mu i} = \gamma^\alpha \left. \{ \chi^\mu, H_\alpha \} \right|_{p = \frac{\partial S}{\partial q}},$$

whence it follows that the matrix  $B_{ki}$  is non-degenerate because of (2.33).

Now we show that the solution of the system of equations (2.47), (2.48) satisfies (2.28a) with the Lagrange multipliers (2.35).

By differentiating (2.47) with respect to time, we find

$$\frac{\partial}{\partial \alpha^A} \frac{\partial S}{\partial t} + \frac{\partial^2 S}{\partial q^i \partial \alpha^A} \dot{q}^i = 0.$$

On the other hand, differentiation with respect to  $\alpha^A$  of the Hamilton-Jacobi equation (2.45) gives

$$\frac{\partial}{\partial \alpha^A} \frac{\partial S}{\partial t} = - \left. \frac{\partial H_0}{\partial p_i} \right|_{p = \frac{\partial S}{\partial q}} \frac{\partial^2 S}{\partial q^i \partial \alpha^A},$$

which yields

$$\frac{\partial^2 S}{\partial q^i \partial \alpha^A} \left( \dot{q}^i - \left. \frac{\partial H_0}{\partial p_i} \right|_{p = \frac{\partial S}{\partial q}} \right) = 0.$$

Consequently, the expression in the parentheses is the linear combination of the zero vectors of the matrix  $(\partial^2 S / \partial q^i \partial \alpha^A)$  with arbitrary coefficients  $\lambda^\mu$ :

$$\dot{q}^i = \left( \frac{\partial H_0}{\partial p_i} + \lambda^\mu \frac{\partial H_\mu}{\partial p_i} \right) \Big|_{p = \frac{\partial S}{\partial q}}. \quad (2.51)$$

In a similar way, the equation for  $p_i$  is obtained by differentiating of (2.45) with respect to time:

$$\dot{p}_i = \frac{\partial^2 S}{\partial q^i \partial q^k} \dot{q}^k + \frac{\partial}{\partial q^i} \frac{\partial S}{\partial t},$$

from which, using (2.45), (2.50) and (2.51), we obtain

$$\dot{p}_i = - \frac{\partial H_0}{\partial q^i} - \lambda^\mu \frac{\partial H_\mu}{\partial q^i}.$$

Thus, by requiring the preservation of the gauges (2.48) in time, one can derive the expression (2.35) for the Lagrange multipliers.

Thus far, a solution of the system of equations (2.48), (2.47), where  $S$  satisfies the Hamilton-Jacobi equations (2.44), (2.45), fully solves the Cauchy problem in the gauge (2.32). In addition, it is obvious that the  $2m$  variables  $(\alpha^A, \beta_A)$  are in one-to-one correspondence with the initial values of the dynamically independent phase variables  $(q^*, p^*)$ .

## 2.2. Peculiarities of the canonical formalism in curved spacetime

To construct the canonical formalism of a dynamical system, one needs to explicitly single out a coordinate which plays the role of time.

It is easy to choose the time in relativistically invariant theories in the flat spacetime. Identifying time with the coordinate  $x^0$  of any Lorentz coordinate system, so that the tangent vector to the  $x^0$ -axis is the Killing vector of the flat spacetime, one can construct the canonical formalism of the theory.

In the curved spacetime, the situation is much more complicated. In general, there are no time-like Killing vectors, along which the spacetime properties are homogeneous. Hence, in each coordinate system, it is impossible to select a unique time. In addition, the physical theory is invariant with respect to the group of general coordinate transformations (2.2), (2.3)

$$\Delta^f \Phi^A(x) = L_f \Phi^A(x), \quad \Delta^f x^\mu(x) = f^\mu, \quad (2.52)$$

where  $L_f \Phi^A(x)$  is the Lie derivative (1.43a) of the tensor field  $\Phi^A(x)$ , containing the term  $f^\mu \partial_\mu \Phi^A(x)$ :

$$L_f \Phi^A(x) = f^\mu \partial_\mu \Phi^A(x) + (\partial_\mu f^\nu) \Omega_\nu^{\mu A} \Phi^B(x). \quad (2.53)$$

Thus, under any transformation of the form (2.52), the question arises, how to define the time in new coordinate systems when it was already introduced somehow in one of them. In view of the locality of the group (2.52), the time should be defined locally, i.e., its definition may be changed in the local vicinity of a point without changing it in the rest of the spacetime.

Therefore, we obtain a coordinate-free definition of time as a system of the space-like hypersurfaces of the constant parameter

$$t = \tau(x), \quad (2.54)$$

where  $\tau(x)$  is a scalar function of coordinates.

If we also want to define a three-dimensional coordinate chart on each hypersurface, this can be done with the help of the four functions

$$x^\alpha = e^\alpha(\mathbf{x}, t) \quad (2.55)$$

of the three-dimensional coordinates  $\mathbf{x} = \{x^a\}$  and the time  $t$ , so that their substitution in (2.54) yields an identity. Since the canonical formalism on the curved spacetime essentially relies on its (3 + 1)-decomposition (2.55), we will study its geometrical properties now.

## 2.3. Geometry of (3 + 1)-decomposition of spacetime

The parameters  $x^a$  represent the coordinate system on the hypersurface (2.55). Let us define the normal basis on the hypersurface by the triad of the tangent vectors  $e_a^\alpha$  and the normal  $n_\alpha$ , such that

$$e_a^\alpha \equiv \partial_a e^\alpha(\mathbf{x}, t), \quad n_\alpha e_a^\alpha = 0. \quad (2.56)$$

The normal vector  $n_\alpha$  determines the type of a surface:

$$n_\alpha n^\alpha = \sigma, \quad \sigma = \pm 1. \quad (2.57)$$

The value  $\sigma = -1$  for space-like and  $\sigma = +1$  for time-like surfaces.

It is natural to define the three-dimensional metric of the hypersurface

$$g_{ab}(\mathbf{x}) = g_{\alpha\beta}(e(\mathbf{x}, t)) e_a^\alpha e_b^\beta.$$

Then any 4-vector  $\lambda^\alpha$  can be decomposed with respect to the normal basis:

$$\begin{aligned} \lambda^\alpha &= n^\alpha \lambda_\perp + e_a^\alpha \lambda^a, \\ \lambda_\perp &= \sigma n_\alpha \lambda^\alpha, \quad \lambda^a = e_a^\alpha \lambda^\alpha, \end{aligned} \quad (2.58)$$

where

$$e_\alpha^a \equiv g_{\alpha\beta} g^{ab} e_b^\beta.$$

The following identities can be immediately derived:

$$e_\alpha^a e_b^\alpha = \delta_b^a, \quad e_a^\alpha e_\beta^\alpha = \delta_\beta^\alpha - \sigma n^\alpha n_\beta.$$

The three-dimensional Riemannian connection on the hypersurface can be introduced via the three-dimensional covariant derivative of a 3-vector  $\lambda^a$ :

$$\lambda^a{}_{|b} \equiv (\lambda^c e_c^\alpha)_{;\beta} e_\alpha^a e_b^\beta,$$

which yields the three-dimensional connection

$$\gamma^a{}_{bc} = e_{c;b}^\alpha e_\alpha^a. \quad (2.59)$$

Here we use the following notation for the four-dimensional covariant derivatives (more exactly, for their projections):

$$\left. \begin{aligned} \lambda^\alpha{}_{;b} &\equiv e_b^\beta \lambda^\alpha{}_{;\beta}, & \lambda_{a;b} &\equiv e_a^\alpha e_b^\beta \lambda_{\alpha;\beta}, \\ \lambda_{\perp;a} &\equiv \sigma n^\beta e_a^\alpha \lambda_{\beta;\alpha}, & \lambda_{a;\perp} &\equiv \sigma n^\alpha e_a^\beta \lambda_{\beta;\alpha}, \\ \lambda_{\perp;\perp} &\equiv n^\alpha n^\beta \lambda_{\alpha;\beta}. \end{aligned} \right\} \quad (2.60)$$

We define the extrinsic curvature  $K_{ab}$  of the hypersurface as

$$K_{ab} = -n_{a;b}. \quad (2.61)$$

Due to the second equation (2.56),  $K_{ab} = e_{a;b}^\alpha n_\alpha = K_{ba}$ . Comparing (2.61) and (2.59), we see that  $K_{ab}$  and  $\gamma^c{}_{ab}$  are the projections of the quantity  $e_{a;b}^\alpha$  on the normal basis. In view of (2.58), this yields the Gauss-Weingarten formula

$$e_{a;b}^\alpha = \sigma K_{ab} n^\alpha + \gamma^c{}_{ab} e_c^\alpha. \quad (2.62)$$

Let us consider the following vector  $N^\alpha$  and its projections on the normal basis:

$$N^\alpha = \frac{de^\alpha(\mathbf{x}, t)}{dt}, \quad N = \sigma n_\alpha N^\alpha, \quad N^a = e_a^\alpha N^\alpha. \quad (2.63)$$

We are interested in the deformation of the normal basis when it moves in the four-dimensional spacetime along the vector (2.63). Taking into account that

$$\partial_\beta e_a^\alpha \frac{de^\beta(\mathbf{x}, t)}{dt} = \partial_a \frac{\partial e^\alpha(\mathbf{x}, t)}{dt},$$

we obtain for the quantity  $\nabla_N e_a^\alpha \equiv e_{a;\beta}^\alpha N^\beta$  the following expression

$$\nabla_N e_a^\alpha = (N_{|a} + \sigma K_{ab} N^b) n^\alpha + (N_{|a}^b - N K^b{}_a) e_b^\alpha. \quad (2.64)$$

In the derivation we used the decomposition (2.58) for the vector (2.63) and the Gauss-Weingarten formula (2.62).

Taking into consideration (2.56) and (2.57), it is possible to obtain from (2.64) the equation for normal vector deformation

$$\nabla_N n^\alpha = -(\sigma N_{|a} + K_{ab} N^b) e^{a\alpha}. \quad (2.65)$$

Using (2.64), we obtain the relation between the extrinsic curvature of the hypersurface and the velocity of the change of the metric

$$\dot{g}_{ab} = \nabla_N g_{ab} = -2N K_{ab} + 2N_{(a|b)}. \quad (2.66)$$

Let  $\chi^\alpha(x)$  be a vector field in the spacetime with a given hypersurface system (2.55), the projections of which on the normal basis are  $\chi^a$  and  $\varphi_\perp$ . Let us analyze how the covariant derivatives of the vector  $\varphi^\alpha$  are related to the three-dimensional covariant derivatives of its projections. Calculating the covariant derivative of the projection  $\varphi_\perp = \sigma n^\alpha \varphi_\alpha$  along the normal vector component (2.63)  $n^\beta N$ , we have

$$\begin{aligned} \delta_\perp \varphi_\perp &\equiv N n^\beta \nabla_\beta \varphi_\perp, \\ N n^\beta \nabla_\beta (\varphi_\perp) &= -N_{|a} \varphi^a + \sigma \varphi_{\perp; \perp} N, \end{aligned}$$

from which

$$N \varphi_{\perp; \perp} = \sigma \delta_\perp \varphi_\perp + \sigma N_{|a} \varphi^a. \quad (2.67)$$

Doing the same with  $\varphi^a = e_a^\alpha \varphi^\alpha$ , we obtain

$$N\varphi_{a;\perp} = \sigma\delta_\perp\varphi_a + \sigma K^b{}_a\varphi_b N - \varphi_\perp N|_a. \quad (2.68)$$

The meaning of the operation  $\delta_\perp$  is that it describes the increment caused by the displacement which is purely orthogonal to the hypersurface and is equal to  $n^\alpha N$ .

Now let us consider the displacements which are tangent to the hypersurface,  $\delta_{||}\varphi_\perp$  and  $\delta_{||}\varphi_a$ . From the point of view of the dependence of the projections of the fixed vector  $\chi^\alpha$  on the normal basis on the embedding of the hypersurface into the four-dimensional spacetime (2.55),  $\varphi_\perp$  and  $\varphi_a$  are functionals of  $e^\alpha(\mathbf{x})$ . Then

$$\delta_{||}\varphi_a \equiv \int d^3y (N^b(\mathbf{y}) e_b^\alpha(\mathbf{y})) \frac{\delta\varphi_a(\mathbf{x})[e]}{\delta e^\alpha(\mathbf{y})} \quad (2.69)$$

is nothing but the Lie derivative of the vector field  $\varphi_a(\mathbf{x})$  along the vector  $N^\alpha(\mathbf{x})$ :

$$\delta_{||}\varphi_a(\mathbf{x}) = L_{\mathbf{N}}\varphi_a(\mathbf{x}). \quad (2.70)$$

Indeed, the expression (2.69) corresponds to the variation of the form  $\varphi_a(\mathbf{x})$  under the displacement of the three-dimensional coordinate chart  $\mathbf{x} = \{x^\alpha\}$  on the vector  $\mathbf{N} = \{N^\alpha\}$ , from which (2.70) follows.

On the other hand, since  $\varphi_a(\mathbf{x})$  is a four-dimensional scalar, the following equation is fulfilled

$$\delta_{||}\varphi_a = \nabla_N\varphi_a \Big|_{N^\alpha = e_a^\alpha N^\alpha},$$

and hence, taking the equation (2.64) into account, with  $N = 0$ , one can derive

$$\delta_{||}\varphi_a = K_{ab}N^b\varphi_\perp + N^b|_a\varphi_b + \varphi_{a;b}N^b.$$

Comparing with the Lie derivative (2.70),  $L_{\mathbf{N}}\varphi_a = \varphi_{a|b}N^b + \varphi_b N^b|_a$ , in view of arbitrariness of the vector field  $N^b(\mathbf{x})$ , we find

$$\varphi_{a;b} = \varphi_{a|b} - K_{ab}\varphi_\perp. \quad (2.71)$$

A similar analysis for  $\varphi_\perp$  gives

$$\varphi_{\perp;b} = \varphi_{\perp|b} + \sigma K_{ba}\varphi^a. \quad (2.72)$$

The equations (2.67), (2.68), (2.71), and (2.72) are generalized to the projections of the higher rank tensors in an obvious way, for example:

$$\varphi_{ab;c} = \varphi_{ab|c} - K_{ac}\varphi_{\perp b} - K_{bc}\varphi_{a\perp},$$

where

$$\varphi_{a\perp} = \sigma n^\alpha e_a^\beta \varphi_{\beta\alpha}, \quad \varphi_{\perp b} = \sigma n^\alpha e_b^\beta \varphi_{\alpha\beta}.$$

Using the formulas (2.67)-(2.72), one can establish relations between the various projections of the four-dimensional Riemann curvature tensor  ${}^4R^\alpha{}_{\beta\mu\nu}$  and

the components of the extrinsic curvature of the hypersurface. Projecting the identity

$$2\varphi_{a;[\mu\nu]} = {}^4R^\lambda{}_{\alpha\mu\nu} \varphi^\lambda$$

on the normal basis with an account of the equations (2.67), (2.68), (2.71), and (2.72), in view of the arbitrariness of the components  $\varphi_\perp$  and  $\varphi_a$  we obtain the Gauss-Codazzi identities:

$${}^4R_{\perp abc} = -2\sigma K_{a[b|c]}, \quad {}^4R_{abcd} = {}^3R_{abcd} - 2\sigma K_{b[d}K_{c]a}, \quad (2.73)$$

where  ${}^3R_{abcd}$  is the Riemann tensor of the extrinsic curvature of the three-dimensional hypersurface. In a similar way:

$$N {}^4R_{a\perp b\perp} = \delta_\perp K_{ab} + NK_{ad}K_b{}^d - \sigma N_{|ab}. \quad (2.74)$$

In order to rewrite the gravitational field Lagrangian in terms of the quantities of the (3+1)-decomposition of the spacetime, we will use the following relations:

$$\begin{aligned} \delta_\perp(g^{1/2}) &= -Ng^{1/2}K, & K &= K^a{}_a, & g &= \det g_{ab}, \\ \delta_\perp(Kg^{1/2}) &= \nabla_N(Kg^{1/2}) - \delta_{||}(Kg^{1/2}), \\ \delta_{||}(Kg^{1/2}) &= L_N(Kg^{1/2}) = (Kg^{1/2}N^a)_{|a}. \end{aligned}$$

With the help of the equations (2.73) and (2.74), we derive

$$\begin{aligned} Ng^{1/2} {}^4R &= Ng^{1/2} \{ {}^3R - \sigma(K_{ab}K^{ab} - K^2) \} + 2\sigma \nabla_N(Kg^{1/2}) \\ &\quad - 2(\sigma Kg^{1/2}N^a + g^{1/2}N^{|a})_{|a}. \end{aligned} \quad (2.75)$$

## 2.4. Canonical formalism for the fields on the curved spacetime

### *Canonical formalism of the gravitational field*

Let us clarify the physical meaning of the quantities of the (3+1)-decomposition. The system of space-like hypersurfaces in the physical spacetime is characterized by the value of the parameter  $\sigma$  equal to  $-1$ . However, we will keep  $\sigma$  arbitrary in all formulas below in order to demonstrate the role of the spacetime signature.

The choice of a family of hypersurfaces (2.55) and of the normal vector (2.56) must satisfy some requirements. It will be mathematically natural to demand that the orientation of the normal basis

$$e_{(\mu)}^\alpha = (e_\perp^\alpha, e_a^\alpha), \quad e_\perp^\alpha \equiv n^\alpha, \quad (2.76)$$

coincides with the orientation of the local frames of the four-dimensional coordinate system. Since the metric in the normal basis has the form

$$e_{(\mu)}^\alpha g_{\alpha\beta} e_{(\nu)}^\beta = \begin{vmatrix} \sigma & 0 \\ 0 & g_{ab} \end{vmatrix},$$

we find

$$|{}^4g|^{1/2} \det e_{(\mu)}^\alpha = \pm g^{1/2},$$

where the upper sign (+) on the right-hand side is fixed by the aforementioned choice of the orientation of the basis,

$$\begin{aligned} n^\alpha e_a^\beta e_b^\mu e_c^\nu \eta_{\alpha\beta\mu\nu} &= \eta_{abc}, \\ \varepsilon_{\alpha\beta\mu\nu} &= |{}^4g|^{1/2} \eta_{\alpha\beta\mu\nu}, \quad \varepsilon_{abc} = g^{1/2} \eta_{abc}. \end{aligned} \quad (2.77)$$

The parameter  $t$  labels the sequence of the space-like sections of the four-dimensional spacetime in the causal order of increasing of the physical time. This means that the function  $N$  is positive:

$$N = \sigma n_\alpha \frac{de^\alpha}{dt}, \quad N > 0. \quad (2.78)$$

This quantity is called a lapse function. The condition (2.78) fixes the direction of the normal vector towards the increase of the time parameter  $t$ . In particular, this means that  $Ndt$  is the normal interval between the two neighbouring hypersurfaces corresponding to the moments  $t$  and  $t + dt$ .

The shift functions  $N^a$  characterize the displacement  $N^a dt$  between the coordinate lines  $x^a$  on the same two neighbouring hypersurfaces, assuming that the coordinate system on the hypersurface corresponding to  $t + dt$ , is normally projected on the hypersurface corresponding to the moment  $t$ .

An important condition that the curve determined by the equation  $x^a = \text{const}$  is time-like leads to inequality

$$N^2 - N_a N^a > 0 \quad (2.79)$$

(it was taken into account that  $\sigma = -1$  in the physical spacetime). The inequality (2.79) actually means that the coordinate  $t$  is time-like and there may be observers at rest on the three-dimensional coordinate chart on the hypersurface.

Eq. (2.77) yields

$$|{}^4g|^{1/2} d^4x = dt d^3x N g^{1/2}, \quad (2.80)$$

and hence, in view of (2.75), we obtain

$$\begin{aligned} \int_V d^4x |{}^4g|^{1/2} {}^4R &= \int_{t_0}^{t_1} dt d^3x N g^{1/2} \{ {}^3R - \sigma (K_{ab} K^{ab} - K^2) \} \\ &+ 2\sigma \int d^3x g^{1/2} K \Big|_{e_0}^{e_1} - 2 \int_{t_0}^{t_1} dt d\sigma_a \left( \sigma K N^a + N |^a \right). \end{aligned} \quad (2.81)$$

Here  $V$  is a four-dimensional spacetime domain bounded by the ‘‘cylindrical’’ surface, the top and the bottom sides of which are the initial and the final hypersurfaces  $e_0 = e(t_0)$  and  $e_1 = e(t_1)$ , and the lateral surface  $\overline{\partial V}$  is topologically a direct product of the two-dimensional sphere  $\partial e(t)$  of radius  $r$  and the time axis in the interval  $t_0 \leq t \leq t_1$ ,  $\overline{\partial V} = \partial e(t) \times [t_0, t_1]$ , see Fig. 2.1.




Figure 2.1: Four-dimensional spacetime domain  $V$  bounded by “cylindrical” surface  $\partial V = \overline{\partial V} \cup e_0 \cup e_1$ ,  $\overline{\partial V} = \partial e(t) \times [t_0, t_1]$ .

The last term in (2.81) is a surface integral over the “cylindrical” spatial asymptotics bounding the space-time “tube” in the interval  $t_0 \leq t \leq t_1$ , and  $d\sigma_a = n_a d\sigma$  is an element of two-dimensional surface  $\partial e(t)$  with a unit normal  $n_a$ . When deriving the form of the second term in (2.81), we took into account that

$$\nabla_N(Kg^{1/2}) = \frac{d}{dt}(Kg^{1/2}).$$

The action of the gravitational field [14] that leads as a result of the variational procedure to Einstein’s equations of the second order, reads as follows<sup>2</sup>:

$$S = \int_V d^4x |^4g|^{1/2} {}^4R - \int_{\partial V} d\Sigma \sigma(2\tilde{K}). \quad (2.82)$$

The surface integral over the boundary  $\partial V$  is determined by the extrinsic curvature  $\tilde{K}$  given by the formula (2.61), where  $n_\alpha$  is an outer normal to the boundary and  $\sigma$  defines the scalar square of the normal vector (2.57). Its role is to compensate in the volume integral of action (2.82) the terms containing the derivatives of the metric which are normal to the boundary. This allows to correctly formulate the boundary value problem for Einstein’s equations as conditions of the extremum of the action (2.82) (see Chapter 1).

<sup>2</sup>Hawking considered the gravitational field in the Euclidean spacetime where  $\sigma = +1$ . In the physical spacetime, his formula is modified, taking into account the metric signature on the boundary  $\partial V$ .

Substituting (2.81) into (2.82), we obtain the action of the gravitational field in terms of the (3 + 1)-decomposition:

$$S = \int_{t_0}^{t_1} dt \left\{ \int d^3x (Ng^{1/2}\mathcal{L}) - H_0 \right\}, \quad (2.83)$$

$$\mathcal{L} = {}^3R - \sigma(K_{ab}K^{ab} - K^2). \quad (2.84)$$

Here the quantity  $H_0$  is determined by the surface integral of the extrinsic curvature of the boundary  $\tilde{K}$  over this two-dimensional boundary of the three-dimensional space,

$$H_0 = 2 \oint d\tilde{\sigma} g^{1/2} \sigma \tilde{K} + 2 \oint d\sigma_a (N^{|a} + \sigma K N^a), \quad (2.85)$$

and the exact form of the measure  $d\tilde{\sigma}$  is obtained if we represent the surface element  $d\Sigma$  of the ‘‘cylindrical’’ lateral boundary of four-dimensional volume as

$$d\Sigma = dt d\tilde{\sigma}, \quad (2.86)$$

where  $d\tilde{\sigma}$  is some element of the measure on  $\partial e(t)$  which is in general different from  $d\sigma$ .

Let us proceed to the construction of the canonical formalism of the theory described by the action (2.82). Taking into account that according to (2.66)

$$K_{ab} = \frac{1}{2N} (2N_{(a|b)} - \dot{g}_{ab}), \quad (2.87)$$

the action (2.82) is a functional of the set of ten variables  $(g_{ab}, N^a, N)$  which replaces the original set of ten metric components  $g_{\mu\nu}$ . The action (2.82) does not contain time derivatives of the lapse and shift functions, therefore,  $N$  and  $N^a$  are the Lagrange multipliers from the point of view of the theory of systems with singular Lagrangians. In order to verify this, we will bring the action (2.82) to the form (2.27).

For this purpose, we introduce the canonical momenta  $\pi^{ab}$  conjugated to the 3-metric components  $g_{ab}$ :

$$\pi^{ab} = \frac{\partial(Ng^{1/2}\mathcal{L})}{\partial\dot{g}_{ab}} = \sigma G^{abcd} K_{cd}, \quad (2.88)$$

$$G^{abcd} = \frac{1}{2} g^{1/2} (2g^{ab}g^{cd} - g^{ac}g^{bd} - g^{ad}g^{bc}). \quad (2.89)$$

Here  $G^{abcd}$  is the so-called covariant DeWitt supermetric [53]. Its contravariant components are given by the expression

$$G_{abcd} = \frac{1}{2} g^{1/2} (g_{ac}g_{bd} + g_{ad}g_{bc} - g_{ab}g_{cd}), \quad (2.89a)$$

$$G_{abcd}G^{cdef} = \delta_{ab}^{ef}.$$

The gravitational field Hamiltonian has the following form:

$$H = \int d^3x \left( \pi^{ab}(\mathbf{x}) \dot{g}_{ab}(\mathbf{x}) - N g^{1/2} \mathcal{L} \right). \quad (2.90)$$

Since from (2.89) the extrinsic curvature can be expressed in terms of the momenta

$$K_{ab} = \sigma G_{abcd} \pi^{cd}, \quad (2.91)$$

the Hamiltonian (2.90) is recast into

$$H = \int d^3x N^{(\mu)} \mathcal{H}_{(\mu)} + H_0, \quad (2.92)$$

where we introduced the unified notation for the lapse and shift functions

$$N^{(\mu)} = (N^\perp, N^a), \quad N^\perp = N, \quad (2.93)$$

and the quantities  $\mathcal{H}_\mu = (\mathcal{H}_\perp, \mathcal{H}_a)$  are given by the following expressions:

$$\mathcal{H}_\perp = -\sigma G_{abcd} \pi^{ab} \pi^{cd} - g^{1/2} {}^3R, \quad (2.94)$$

$$\mathcal{H}_a = -2g_{ac} \pi^{cd} |_{|d}. \quad (2.95)$$

Therefore, in terms of the canonical variables the action (2.82) reads

$$S_{HE}[g_{ab}, \pi^{ab}, N^{(\mu)}] = \int_{t_0}^{t_1} dt \left\{ \int d^3x \pi^{ab} \dot{g}_{ab} - H_0 - \int d^3x N^{(\mu)} \mathcal{H}_{(\mu)} \right\}, \quad (2.96)$$

which makes it obvious that the functions  $N^{(\mu)}$  are the Lagrange multipliers.

The stationary condition of the action (2.96) with respect to variations of  $N^{(\mu)}$  leads to the equations

$$\mathcal{H}_\perp = 0, \quad (2.97)$$

$$\mathcal{H}_a = 0, \quad (2.98)$$

which are the constraints since they do not contain the time derivatives of the phase variables.

By direct evaluation of the Poisson brackets, one can show that these constraints satisfy the involution relations (2.29):

$$\left. \begin{aligned} \{\mathcal{H}_\perp(\mathbf{x}), \mathcal{H}_\perp(\mathbf{x}')\} &= -\sigma (\mathcal{H}^a(\mathbf{x}) \partial_a \delta(\mathbf{x}, \mathbf{x}') - \mathcal{H}^a(\mathbf{x}') \partial_a \delta(\mathbf{x}', \mathbf{x})), \\ \{\mathcal{H}_a(\mathbf{x}), \mathcal{H}_\perp(\mathbf{x}')\} &= \mathcal{H}_\perp(\mathbf{x}) \partial_a \delta(\mathbf{x}, \mathbf{x}'), \\ \{\mathcal{H}_a(\mathbf{x}), \mathcal{H}_b(\mathbf{x}')\} &= \mathcal{H}_b(\mathbf{x}) \partial_a \delta(\mathbf{x}, \mathbf{x}') - \mathcal{H}_a(\mathbf{x}') \partial_b \delta(\mathbf{x}', \mathbf{x}). \end{aligned} \right\} \quad (2.99)$$

Accordingly, we found that (2.97), (2.98) are the first-kind constraints.

The canonical equations for the phase variables of the gravitational field are obtained by the variation of the action (2.96) with respect to  $g_{ab}$  and  $\pi^{ab}$ . The term  $H_0$  in the total Hamiltonian of the system (2.92) is a surface integral, which does not explicitly contribute to the dynamical equations under the variation. As a result, the equations of motion in terms of the Poisson brackets take the following form:

$$\dot{g}_{ab}(\mathbf{x}) = \int d^3x' \{g_{ab}(\mathbf{x}), \mathcal{H}_{(\mu)}(\mathbf{x}')\} N^{(\mu)}(\mathbf{x}'), \quad (2.100)$$

$$\dot{\pi}_{ab}(\mathbf{x}) = \int d^3x' \{\pi^{ab}(\mathbf{x}), \mathcal{H}_{(\mu)}(\mathbf{x}')\} N^{(\mu)}(\mathbf{x}'). \quad (2.101)$$

Computing the Poisson bracket in (2.100), we obtain the equation (2.66), where the extrinsic curvature  $K_{ab}$  is expressed in terms of the momenta  $\pi^{ab}$  via (2.91). Similarly, one can show that the equation (2.101) is equivalent to the following set of Einstein's equations projected on the normal basis:  $NG_{ab} = 0$ .

The constraints (2.97), (2.98) are also the components of Einstein's equations

$$\mathcal{H} = 2\sigma g^{1/2} G_{\perp\perp}, \quad (2.102)$$

$$\mathcal{H}_a = 2g^{1/2} G_{a\perp}. \quad (2.103)$$

Therefore, the system of equations (2.100), (2.101) is completely equivalent to Einstein's equations  $G_{\mu\nu} = 0$ .

One can read from (2.99) the structural functions of the involution relations (2.29):

$$\begin{aligned} U_{\mathbf{ax}, \perp \mathbf{x}'}^{\perp \mathbf{y}} &= \delta(\mathbf{x}, \mathbf{y}) \partial_a \delta(\mathbf{x}, \mathbf{x}'), \\ U_{\mathbf{ax}, \mathbf{bx}'}^{c\mathbf{y}} &= \partial_a \delta(\mathbf{x}, \mathbf{x}') \delta_b^c \delta(\mathbf{y}, \mathbf{x}) - \partial_b \delta(\mathbf{x}', \mathbf{x}) \delta_a^c \delta(\mathbf{y}, \mathbf{x}'), \\ U_{\mathbf{ax}, \perp \mathbf{x}'}^{b\mathbf{y}} &= U_{\mathbf{ax}, \mathbf{bx}'}^{\perp \mathbf{y}} = 0, \\ U_{\perp \mathbf{x}, \perp \mathbf{x}'}^{a\mathbf{y}} &= -\sigma(g^{ab}(\mathbf{x}) \partial_b \delta(\mathbf{x}, \mathbf{x}') \delta(\mathbf{y}, \mathbf{x}) - g^{ab}(\mathbf{x}') \partial_0 \delta(\mathbf{x}', \mathbf{x}) \delta(\mathbf{y}, \mathbf{x}')). \end{aligned} \quad (2.104)$$

Since the phase variables in the interior points of the spacetime domain  $V$  commute in the sense of the Poisson brackets with the quantities on the boundary  $\partial V$ , the structural functions  $V_{(\nu)}^{(\mu)}$  vanish.

Using the results obtained in (2.30), (2.31), one can see that the canonical action (2.96) is invariant with respect to transformations:

$$\delta^{\mathcal{F}} \pi^{ab} = \int d^3x' \{\pi^{ab}, \mathcal{H}_{(\mu)}(\mathbf{x}')\} \mathcal{F}^{(\mu)}(\mathbf{x}'), \quad (2.105)$$

$$\delta^{\mathcal{F}} g_{ab} = \int d^3x' \{g_{ab}, \mathcal{H}_{(\mu)}(\mathbf{x}')\} \mathcal{F}^{(\mu)}(\mathbf{x}'), \quad (2.106)$$

$$\delta^{\mathcal{F}} N = \dot{\mathcal{F}}_{\perp} - N^a \mathcal{F}_{\perp|a} + N_{|a} \mathcal{F}^a, \quad (2.107)$$

$$\delta^{\mathcal{F}} N^a = \dot{\mathcal{F}}^a - (\mathcal{F}^a{}_{|b} N^b - N^a{}_{|b} \mathcal{F}^b) + \sigma(N \mathcal{F}_{\perp}{}^{|a} - N^a \mathcal{F}_{\perp}), \quad (2.108)$$

with the infinitesimal parameters  $\mathcal{F}^{(\mu)} = (\mathcal{F}^\perp, \mathcal{F}^a)$ ,  $\mathcal{F}^\perp \equiv \mathcal{F}_\perp$ . Let us show that the transformations (2.106)-(2.108), evaluated in the phase space for the extremal values of the momenta (2.88) [i.e., for the momenta which are the solution of the equations (2.100) in terms of  $g_{ab}$  and  $\dot{g}_{ab}$ ], are equivalent to the transformation of the general coordinate group (2.52) with the parameters

$$f^\mu = e^\mu_{(\nu)} \mathcal{F}^{(\nu)}, \quad e^\mu_{(\nu)} \mathcal{F}^{(\nu)} = n^\mu \mathcal{F}_\perp + e^\mu_a \mathcal{F}^a, \quad (2.109)$$

supplemented by the slicing invariance condition (2.55)

$$\Delta^f e^\alpha(\mathbf{x}, t) = 0. \quad (2.110)$$

Indeed, in accordance with (2.52) and (2.110),

$$\delta^{\mathcal{F}} g_{ab} = e^\alpha_a e^\beta_b \Delta^f g_{\alpha\beta} = 2f_{a,b},$$

and hence using (2.71), (2.91) in combination with the fact that  $f_\perp = \mathcal{F}_\perp$  and  $f_a = g_{ab} \mathcal{F}^b$ , we derive (2.106). Similarly, we have

$$\delta^{\mathcal{F}} N = \sigma \Delta^f n_\alpha \dot{e}^\alpha. \quad (2.111)$$

The variation of the normal vector, caused by the variation of the four-dimensional metric, is obtained as a result of the variation of the equations (2.56), (2.57):

$$\Delta^f n_\alpha = -\frac{1}{2} \sigma n_\alpha (n_\mu n_\nu \Delta^f g^{\mu\nu}),$$

from which

$$\Delta^f n_\alpha = \sigma n_\alpha f_{\perp;\perp}. \quad (2.112)$$

Substituting (2.112) into (2.111) and taking into account (2.67), we obtain (2.107), since  $\delta_\perp \mathcal{F}_\perp = \dot{\mathcal{F}}_\perp - \mathcal{F}_{\perp|a} N^a$ . In the same way, the last relation (2.108) is derived.

Thus, we have demonstrated that the transformations (2.106)-(2.108) on the surface of the extremal momenta realize the general coordinate group (2.51) in the configuration space. Since this transformation of coordinates

$$x^\alpha \longrightarrow x^\alpha + f^\alpha(x) \quad (2.113)$$

does not change, in view of (2.110), the form of the functions  $e^\alpha(\mathbf{x}, t)$ , they can be treated as the simultaneous transformation of the coordinate system (2.113) and the change of the hypersurface system

$$e^\alpha(\mathbf{x}, t) \longrightarrow e^\alpha(\mathbf{x}, t) - f^\alpha(x) \Big|_{x=e(\mathbf{x}, t)},$$

so that the total change of the embedding functions  $e^\alpha(\mathbf{x}, t)$  vanishes.

Consequently, the transformations (2.105)-(2.108) realize the group of the hypersurface deformations with the parameters  $(-f^\alpha) = -e^\alpha_{(\mu)} \mathcal{F}^{(\mu)}$ . Eq. (2.109)

shows that  $\mathcal{H}(\mathbf{x}) = \mathcal{H}_\perp(\mathbf{x})$  is the generator of the hypersurface deformations at a point  $\mathbf{x}$  in the direction which is normal to the hypersurface. This generator is called a superhamiltonian. In a similar way, the constraints  $\mathcal{H}_a(\mathbf{x})$ , which are called supermomentum components, are the generators of the tangential deformations or the transformations of the three-dimensional coordinate chart on the hypersurface. Therefore, in particular,

$$\int d^3x' \{g_{ab}(\mathbf{x}), \mathcal{H}_c(\mathbf{x}')\} N^c(\mathbf{x}') = L_N g_{ab}(\mathbf{x}).$$

The Lie derivative of the momentum field is also given by a similar expression.

Let us note that, in contrast to the general coordinate transformations that constitute the group, the transformations realizing them in the phase space do not constitute a group since the structural functions (2.104) are not constant and depend on the phase space variables, i.e., these transformations constitute the canonical “pseudogroup”. The difference of its structure from the group structure is that the coefficients of the transition from the parameters  $f^\mu$  to  $\mathcal{F}^{(\mu)}$  in (2.109) are not constant and depend on the metric. One can show that the structural functions of the involution relations follow from the structural constants of the general coordinate group, and vice versa, the involution relations (2.99) lead to the commutation of the general coordinate transformations using the Lie brackets of the vector fields.

### *Geometrodynamics with the matter sources*

The system of the gravitational and the matter fields is described by the total action

$$S = S_{HE} + S_m. \quad (2.114)$$

As in Chapter 1, we assume that the interaction of the matter field  $\Psi$  with the gravitation is minimal. Therefore, the action of an unspecified matter field  $\Psi$  has the following form:

$$S_m = \int d^4x |{}^4g|^{1/2} \mathcal{L}_m(\Psi, \Psi_{;\mu}). \quad (2.115)$$

The construction of the canonical formalism for the system (2.114) starts with recasting of the matter field action (2.115) into the (3 + 1)-decomposed form. As before, we need to calculate the projections of the tensor field  $\Psi$  on the normal basis. The complete set of these projections is denoted  $\varphi$ : these will be the phase coordinates in the theory. The momenta  $p$  conjugated to them are determined in a standard way,

$$p = N g^{1/2} \frac{\partial \mathcal{L}_m}{\partial \dot{\varphi}},$$

where we took into account the rule (2.80) of transition to the new spacetime coordinates. This procedure recasts the action (2.115) in terms of the canonical

variables into a form which is similar to (2.83):

$$S_m[\varphi, p] = \int dt \int d^3x \left\{ p\dot{\varphi} - N^{(\mu)} \mathcal{H}_{(\mu)}^m \right\}, \quad (2.116)$$

where

$$\mathcal{H}_{(\mu)}^m = \mathcal{H}_{(\mu)}^m(\varphi, p, g_{ab}, K_{ab}) \quad (2.117)$$

are the superhamiltonian and the supermomenta of the matter field on the curved spacetime, which are the functions of the phase variables of the three-dimensional metric and extrinsic curvature  $K_{ab}$ . It is important that  $\mathcal{H}_{(\mu)}^m$  does not depend on the lapse and shift functions, so that the field Hamiltonian in (2.116) turns out to be a linear function of the latter.

The construction of momenta  $p^{ab}$  conjugated to the 3-metric of the gravitational field for the system (2.114) leads to the result

$$p^{ab} = \pi^{ab} + P^{ab}, \quad (2.118)$$

$$P^{ab} = \frac{\partial}{\partial \dot{g}_{ab}} \left( N g^{1/2} \mathcal{L}_m \right), \quad (2.119)$$

where the purely gravitational momentum  $\pi^{ab}$  is determined by (2.88). The quantity  $P^{ab}$  (2.119) is an additional contribution to the gravitational field momentum, due to the presence of time derivatives of the metric in the matter field Lagrangian. It is clear that the derivatives of the metric enter  $\mathcal{L}_m$  in terms of the Riemannian connection in the covariant derivatives  $\Psi_{;\mu}$ . Since the variation of the Christoffel symbols with respect to the metric is expressed in terms of the covariant derivatives of the metric variations  $\delta g_{\mu\nu}$ , one can write the following variational identity:

$$\frac{\partial \mathcal{L}_m}{\partial \Psi_{;\mu}} \delta g(\Psi_{;\mu}) = \frac{1}{2} P^{\mu\nu\sigma} (\delta g_{\mu\nu})_{;\sigma}, \quad (2.120)$$

which also serves as a definition for the tensor  $P^{\mu\nu\sigma}$  (see Chapter 1, Sec. 1.3.). Substituting (2.120) into (2.119) and using (2.68), we arrive at

$$P^{ab} = \frac{1}{2} g^{1/2} P^{ab\perp}. \quad (2.121)$$

For the theories of matter that does not interact with the derivatives of the metric, i.e., when  $\Psi_{;\mu} = \partial_\mu \Psi$  in the action (2.115) (which is the case, for example, for the scalar or the electromagnetic fields) one finds  $P^{\mu\nu\sigma} = 0$ , and the momentum of the gravitational field  $p^{ab}$  identically coincides with the purely gravitational momentum

$$p^{ab} = \pi^{ab}. \quad (2.122)$$

It is convenient to rewrite the superhamiltonian of the matter  $\mathcal{H}_\perp^m$  by isolating the term of a special structure as follows:

$$\mathcal{H}_\perp^m = \mathcal{H}^m + 2K_{ab} P^{ab}. \quad (2.123)$$

Now let us plug into (2.114) the expressions (2.96), (2.116) and (2.123), and in the latter equation we make use of (2.87). Integrating the term  $-2N_{(a|b)}P^{ab}g^{1/2}$  by parts and dropping an inessential surface integral, we obtain the canonical action of the gravitating system:

$$S[g_{ab}, p^{ab}, \varphi, p, N^{(\mu)}] = \int_{t_0}^{t_1} dt \left\{ \int d^3x \left( p^{ab} \dot{g}_{ab} + p \dot{\varphi} - N^{(\mu)} H_{(\mu)} \right) - H_0 \right\}. \quad (2.124)$$

Here the superhamiltonian  $H_{\perp}$  and the supermomenta  $H_a$  of the total system are given by:

$$H_{\perp} = \mathcal{H}^m + \mathcal{H} \Big|_{\pi^{ab} = p^{ab} - P^{ab}}, \quad (2.125)$$

$$H_a = \mathcal{H}_a^m - 2g_{ab} p^{bc} \Big|_c. \quad (2.126)$$

The extrinsic curvature  $K_{ab}$  should be expressed in these formulas in terms of the phase variables from the solution of the equation (2.91), taking the following form:

$$K_{ab} = \sigma G_{abcd} [p^{cd} - P^{cd}(K_{ef})].$$

One can show that the superhamiltonian (2.125) and the supermomenta (2.126) satisfy the involution relations

$$\{H_{(\mu)}, H_{(\nu)}\} = U_{(\mu)(\nu)}^{(\alpha)} H_{(\alpha)}, \quad (2.127)$$

where the Poisson brackets are defined for the complete set of the phase variables  $g_{ab}, \pi^{ab}, \varphi$  and  $p$ , and with the same structural functions as for the pure gravitation<sup>3</sup>. Consequently,  $H_{(\mu)}$  are the first-kind constraints and they are generators of the pseudogroup of the space-like hypersurface deformations, considered in the previous section, that now acts in the total space of the variables  $g_{ab}, p^{ab}, \varphi, p; \Phi = (g_{ab}, p^{ab}, \varphi, p)$ ,

$$\delta^{\mathcal{F}} \Phi = \int d^3x' \{ \Phi, H_{(\mu)}(\mathbf{x}') \} \mathcal{F}^{(\mu)}(\mathbf{x}'). \quad (2.128)$$

Therefore, the gravitating system described by the action (2.124) is degenerate, due to its invariance with respect to the transformations (2.128) realizing the general coordinate transformations in the configuration space.

The first-kind constraints of the theory

$$H_{(\mu)}(g_{ab}, p^{ab}, \varphi, p) = 0$$

---

<sup>3</sup>This conclusion is based on the independence of the total action (2.124) on the choice of a slicing (2.55) in the interior domain of space-time volume.

are equivalent to the four Einstein field equations with the tensor sources

$$\begin{aligned} H_{\perp} &= 2\sigma g^{1/2} \left( G_{\perp\perp} - \frac{1}{2} T_{\perp\perp} \right), \\ H_a &= 2g^{1/2} \left( G_{\perp a} - \frac{1}{2} H_{\perp a} \right), \end{aligned}$$

where  $H_{\perp\perp}$  and  $H_{\perp a}$  are the projections of the energy-momentum tensor  $H_{\mu\nu}$  on normal basis vectors. The equations which are obtained by the variation of (2.124) with respect to  $g_{ab}$  and  $\pi^{ab}$  are equivalent to the remaining six Einstein's equations  $G_{ab} - \frac{1}{2} T_{ab} = 0$ .

Let us note that in a particular case of the interaction without derivatives, when  $P^{ab} = 0$  and the relation (2.122) is valid, the structure of the formulas (2.124)-(2.126) is essentially simplified. The superhamiltonian  $H_{\perp}$  of the total system is decomposed into a sum of the purely gravitational superhamiltonian plus the superhamiltonian of the matter in an external gravitational field.

The non-dynamical character of the torsion field in ECT leads to an insignificant modification of the superhamiltonian  $H_{\perp}$  of the total system, which does not destroy the structure of the general action (2.124) and the involution relations (2.127). Therefore, the canonical formalism of the gravitating fields in the Riemann-Cartan space has the same structure as in the Riemann space.

# 3

## Dynamics of gravity theories of Hilbert-Einstein type

### 3.1. The problem of the “frozen” formalism

The dynamical contents of the field theory is determined by the sector of its physical degrees of freedom cleared of the ambiguity caused by the invariance under the action of the local group.

The gravity theory of the Hilbert-Einstein type in the space of a general affine connection can be effectively formulated in the Riemannian space. The torsion fields in the theories of this type are non-dynamical and they are algebraically excluded in terms of the components of the spin tensor of matter. Hence, they cannot affect the dynamical content of the theory. In this context, without the loss of generality, we will consider the dynamics of Einstein’s GR with arbitrary matter sources.

The canonical action of the gravitating system (2.124) is a particular case of the general class of degenerate systems described by the action (2.27). The lapse and shift functions  $N^{(\mu)}$  are the Lagrange multipliers. Variation of (2.124) with respect to the phase variables  $\Phi$  and the Lagrange multipliers yields the equations

$$\dot{\Phi} = \delta^N \Phi, \tag{3.1}$$

$$H_{(\mu)}(\Phi) = 0, \tag{3.2}$$

where  $\delta^N \Phi$  is the canonical transformation (2.128) defined by the parameters  $\mathcal{F}^{(\mu)} = N^{(\mu)}$ . The Lagrange multipliers are not determined from the equations of motion and remain arbitrary.

As a result, we find ourselves in a strange situation: the right-hand side of the canonical equations (3.1) turns out to be a pure canonical transformation (2.30), (2.30a) that leaves the action invariant, i.e., the dynamics of the phase variables reduces to canonical transformations with arbitrary parameters  $N^{(\mu)}$  realizing the local invariance group in the phase space. It appears that there are no dynamical degrees of freedom in the gravitating system, since its evolution reduces to transformations from the invariance group of the action.

The reason is that in the total Hamiltonian of the system

$$H = H_0 + \int d^3x N^{(\mu)} H_{(\mu)} \quad (3.3)$$

the quantity  $H_0$  does not explicitly contribute to the field equations (3.1), (3.2), since it is the surface integral (2.85). In fact, a contribution does exist, but it is implicit in its nature. The Hamiltonian  $H_0$  ensures the correctness of the variational problem for the action, leading to the proper boundary value problem for the second-order field equations [14].

Such a situation producing an illusion of the absence of dynamics in the gravity theory is called a problem of the “frozen” formalism. In a more explicit form, this problem arises in cosmological closed models, where the surface integral  $H_0$  is absent,  $H_0 \equiv 0$ , and the complete Hamiltonian (3.3) vanishes on the surface of the constraints (3.2) in the phase space. The problem of the “frozen” formalism arises more seriously when we try to construct the quantum gravity theory, since in a naive approach to the quantization the vanishing of the Hamiltonian (3.3) of the system means the absence of evolution of the vector of the quantum field state

$$|\Psi(t)\rangle = \exp(iHt) |\Psi(0)\rangle = |\Psi(0)\rangle.$$

For the development of the canonical formalism of degenerate systems, analyzed in the previous chapter for the general theory, one needs to address the arising problem of the “frozen” formalism.

### 3.2. Arnowitt-Deser-Misner procedure of selection of physical degrees of freedom

Let us assume that the number of pairs of the phase variables of the matter field  $\varphi$  is equal to  $N$ . Thus, the general dimension of the phase space of a gravitating system is equal to  $2n = 12 + 2N$ , since the total number of the gravitational field momenta  $p^{ab}$  plus the components of the 3-metric  $g_{ab}$  is equal to 12. In accordance with the general procedure of the previous chapter, the number of the dynamical degrees of a degenerate system with  $m = 4$  constraints of the first kind (3.2) is equal to  $n - m = 2 + N$ . Therefore, the pure gravitational field accounts for the two dynamical degrees of freedom. Let us find these variables.

Consider the canonical transformation from the initial set of the phase variables  $\Phi = (g_{ab}, p^{ab}, \varphi, p)$  to the new set of  $(12 + 2N)$  variables:

$$(g_{ab}, p^{ab}, \varphi, p) \longrightarrow (X^\mu, \Pi_\mu; g^A, p_A), \quad (3.4)$$

where  $(X^\mu, \Pi_\mu)$  are the four pairs of the canonically conjugated phase variables, and  $(g^A, p_A)$  are the remaining  $(N + 2)$  pairs  $(A = 1, 2, \dots, N + 2)$ .

This transformation may be absolutely arbitrary. The only restriction imposed on it is as follows. Expressing the old phase variables in terms of the new ones<sup>1</sup>

$$\Phi = \Phi [X^\mu, \Pi_\mu; g^A, p_A], \quad (3.5)$$

we obtain the geometrodynamical constraints (2.125), (2.126) in terms of the new phase variables:

$$H_{(\mu)} = H_{(\mu)}(\mathbf{x}) [X^\alpha, \Pi_\alpha; g^A, p_A]. \quad (3.6)$$

One can define the matrix

$$J_{(\nu)}^{(\mu)}(\mathbf{x}, \mathbf{y}) \equiv \frac{\delta H_{(\nu)}(\mathbf{y})}{\delta \Pi_\mu(\mathbf{x})} = \{X^\mu(\mathbf{x}), H_{(\nu)}(\mathbf{y})\}. \quad (3.7)$$

The transformation (3.4) must satisfy the condition of invertibility of the functional matrix (3.7),  $\det \left( J_{(\nu)}^{(\mu)}(\mathbf{x}, \mathbf{y}) \right) \neq 0$ , so that the inverse matrix exists and reads  $J^{-1(\nu)}_{(\lambda)}(\mathbf{y}, \mathbf{z})$ ,

$$\int d^3y J_{(\nu)}^{(\mu)}(\mathbf{x}, \mathbf{y}) J^{-1(\nu)}_{(\lambda)}(\mathbf{y}, \mathbf{z}) = \delta_{(\lambda)}^{(\mu)} \delta(\mathbf{x} - \mathbf{z}),$$

and then one can solve the constraint equations  $H_{(\mu)}(\mathbf{x}) [X^\alpha, \Pi_\alpha; g^A, p_A] = 0$  with respect to  $\Pi_\alpha$ ,

$$\Pi_\mu(\mathbf{x}) = -P_\mu(\mathbf{x}) [X^\alpha; g^A, p_A]. \quad (3.8)$$

Since the transformation (3.4) is canonical, the action (2.124) in terms of new variables takes the following form:

$$S = \int_{t_0}^{t_1} dt \int d^3x \left\{ p_A \dot{g}^A + \Pi_\mu \dot{X}^\mu - H_0 - N^{(\mu)} H_{(\mu)}(\mathbf{x}) [X, \Pi; g^A, p_A] \right\}, \quad (3.9)$$

where the surface integral  $H_0$  must also be written using (3.5) in terms of the new phase variables.

The equations obtained from the variational principle for the action (3.9) are equivalent to the complete set of the canonical equations. Let us write the equation from the variation with respect to  $\Pi_\mu$ :

$$\dot{X}^\mu(\mathbf{x}) = \int d^3y N^{(\nu)}(\mathbf{y}) J_{(\nu)}^{(\mu)}(\mathbf{x}, \mathbf{y}). \quad (3.10)$$

---

<sup>1</sup>Since nothing is assumed about the locality of the transformation (3.4),  $\Phi$  are the functionals of the new variables as the functions of the three-dimensional coordinates  $\mathbf{x}$ .

In accordance with the general procedure of Chapter 2 we introduce the gauge conditions:

$$\chi^\mu(\mathbf{x}) \equiv X^\mu(\mathbf{x}) - f^\mu(t, \mathbf{x}) = 0, \quad (3.11)$$

where  $f^\mu(t, \mathbf{x})$  are some functions of the coordinates and the time parameter  $t$ . Then the parametrization (2.38) of the complete set of the phase variables in terms of the physical degrees of freedom  $(q^*, p^*) = (g^A, p_A)$  via

$$X^\mu = f^\mu(t, \mathbf{x}), \quad \Pi_\mu = -P_\mu[f^\alpha; g^A, p_A], \quad (3.12)$$

guarantees that the complete set of the constraints and gauge conditions (2.39) or (3.11) is satisfied. The action (3.9) in terms of the physical degrees of freedom is obtained after substituting (3.12) into (3.9):

$$S[g^A, p_A] = \int_{t_0}^{t_1} dt \left\{ \int d^3x p_A \dot{g}^A - H_{\text{phys}}(t)[g^A, p_A] \right\}, \quad (3.13)$$

where the total *physical* Hamiltonian is equal to

$$H_{\text{phys}}(t)[g^A, p_A] = H_0[g^A, p_A] + \int d^3x P_\mu(\mathbf{x}) [f^\alpha; g^A, p_A] \dot{f}^\mu(t, \mathbf{x}). \quad (3.14)$$

When varying the action (3.13) with respect to  $(g^A, p_A)$ , we obtain the closed system of equations for the physical degrees of freedom, however, the lapse and shift functions, which are the Lagrange multipliers, remain undetermined. Without the knowledge of them, the dynamics of the gravitating system is not fully determined. If we require the preservation of the gauge conditions (3.11) in time, we obtain the equation (3.10) for  $N^{(\mu)}(\mathbf{x})$ , where the variables  $\dot{X}^\mu(\mathbf{x})$  are replaced with  $\dot{f}^\mu(t, \mathbf{x})$ , and hence

$$N^{(\mu)}(\mathbf{x}) = \int d^3y J^{-1(\mu)}_{(\nu)}(\mathbf{x}, \mathbf{y}) \dot{f}^\nu(t, \mathbf{y}). \quad (3.15)$$

It is easy to see that these equations are equivalent to the equations (2.35) and with  $H_0 = 0$  nonvanishing values of  $N^{(\mu)}(\mathbf{x})$  are entirely due to explicit dependence of the gauge conditions (3.11) on time.

Therefore, the equations (3.13)-(3.15) solve the problem of selection of the dynamically independent degrees of freedom, and one can construct a nontrivial physical Hamiltonian. Let us consider the arbitrariness of the choice of the functions  $f^\mu(t, \mathbf{x})$  in the additional conditions (3.11).

The lapse and shift functions must satisfy the following conditions:

$$N > 0, \quad N^2 - N^\alpha N_\alpha > 0. \quad (3.16)$$

The first condition determines the positive direction of the time counted by the parameter  $t$ , and the second condition means that the coordinate curve  $t$  is time-like, i.e., it lies within the light cone. Strictly speaking, the second condition is not mandatory, however, one should impose this condition if one

requires that the coordinate system  $(t, \mathbf{x})$  could be realized using the physical bodies.

In addition to the arbitrariness related with functions  $f^\mu(t, \mathbf{x})$ , there is a large arbitrariness in the choice of the new set of phase variables (3.4) connected with an old set by a canonical transformation.

Due to the complicated differential character of the gravitational constraints, there are no spatially-local canonical transformations (3.4) leading to the local function for the Hamiltonian (3.14) density [50]. It means that in general case the ADM method produces a non-local action for the physical variables and the non-local expressions for the lapse and shift functions (3.15). As a rule, the functional matrix (3.7) is a differential operator, and hence  $J^{-1}_{(\nu\mathbf{y})}^{(\mu\mathbf{x})}$  is an integral operation requiring to fix the boundary conditions at the spatial boundary.

### 3.3. Asymptotically flat and closed worlds

It was established above that the construction of a nontrivial physical Hamiltonian is performed simultaneously with the selection of the dynamically independent variables of the gravitating system. In addition, the lapse and shift functions are fixed uniquely, i.e., the physical time and coordinates are defined. Without referring to the ADM procedure, it is impossible to construct a regular Hamiltonian, since the naive Hamiltonian (3.3) cannot generate the dynamics of the theory and produces the problem of the “frozen” formalism. That is quite clear, since the basic notions of any theory are not the energy and momentum, but the time and space where the dynamics of the physical degrees of freedom takes place. The energy and momentum are in a certain sense conjugated quantities to the time and coordinates, they are the generators of translations in the spacetime.

To construct the dynamics of the gravitating system, it is necessary to distinguish two important subclasses of problems with essentially different physical properties: the asymptotically flat insular worlds and the closed cosmological systems.

#### *The energy of asymptotically flat worlds*

Asymptotically flat worlds are characterized by the presence of a region in them, where the timelike and spacelike Killing vector fields<sup>2</sup> exist. They cannot be determined globally, since the space on the whole is essentially inhomogeneous, however, they exist asymptotically.

An open asymptotically flat space should be considered as the region of the spacetime with the boundary, all points of which are directed to the spatial infinity. In the asymptotically flat space, the “lateral cylindrical” boundary

---

<sup>2</sup>The worlds, where only timelike asymptotic Killing vector fields exist, can be analyzed similarly to the asymptotically flat worlds.

$\overline{\partial V}$  (see Fig. 2.1) is obtained in the limit of  $r \rightarrow \infty$  for the radius of the two-dimensional sphere representing  $\partial e(t)$ . Besides, the vertical line (parallel to the time axis) generating the cylindrical boundary  $\partial V$  is set orthogonal to the hypersurfaces  $e(t)$ , that is on the whole, the boundary  $\overline{\partial V}$  is orthogonal to  $e(t)$  for any value of  $t$ . Therefore, the normal  $n_\mu$  to  $\overline{\partial V}$  lies in the plane of the hypersurface  $e(t)$  and is fixed by its projections  $n_a$  on  $e(t)$ . Moreover, the vector  $n_a$  coincides with the normal to the two-dimensional sphere  $\partial e(t)$  in the three-dimensional space  $e(t)$ . The element of the measure  $d\tilde{\sigma}$  turns out to be equal to

$$d\tilde{\sigma} = N d\sigma. \quad (3.17)$$

Computing the extrinsic curvature of the boundary  $\partial V$ , one finds the expression for its trace

$$K = -\frac{1}{N} (n^a N)_{|a}. \quad (3.18)$$

Substituting (3.17) and (3.18) into (2.85) and taking into account that the lapse function of the flat space is equal to one, we have

$$H_0 = \int_{\partial e(t)} d\sigma \left\{ -2 [n^a]_{|a} N + 2n_a (p^{ab} N_b - p N^a) \right\}, \quad (3.19)$$

where  $[n^a]_{|a}$  is the difference of the values of  $n^a|_a$  for the curved and the flat spaces.

The boundary conditions for the metric in an asymptotically flat space are fixed as

$$\left. \begin{aligned} g_{ab} &= \delta_{ab} + \frac{M}{8\pi r} n_a n_b + \mathcal{O}(r^{-2}), & n^a &= \frac{x^a}{r}, \\ N &= 1 - \frac{M}{16\pi r} + \mathcal{O}(r^{-2}), \\ N_a &= \mathcal{O}(r^{-2}), \end{aligned} \right\} \quad (3.20)$$

where  $M$  is the total mass of an isolated distribution of matter.

Since  $d\sigma = r^2 d\Omega^2 + \mathcal{O}(r^{-2})$ , where  $d\Omega^2$  is the element of a solid angle, in the limit of  $r \rightarrow \infty$  only the terms of order not smaller than  $\mathcal{O}(r^{-2})$  contribute to  $H_0$ . Therefore, when calculating  $[n^a]_{|a}$ , it is sufficient to restrict oneself just to the first term of the power series of the metric expansion in its deviation from the flat metric  $h_{ab} = g_{ab} - \delta_{ab}$ , which yields

$$-2 [n^a]_{|a} = n^a \left( h_{ab}{}^{|b} - h_b{}^b{}_{|a} \right) + \mathcal{O}(r^{-3}).$$

The momenta of the gravitational field are equal to

$$p^{ab} = \frac{g^{1/2}}{2N} (g^{ab} g^{cd} - g^{ac} g^{bd}) (\dot{g}^{cd} - 2N_{(c|d)}) = \mathcal{O}(r^{-3})$$

and they do not contribute to (3.19).

Thus, the Hamiltonian in asymptotically flat worlds takes the following form:

$$H_0 = \int_{\partial e} d\sigma_a \left( h^{ab} |_{|b} - h_b^{b|a} \right). \quad (3.21)$$

This expression is covariant and does not depend on a choice of the spatial coordinates. On the other hand, in Cartesian coordinates, it reduces to the well-known Arnowitt-Deser-Misner energy [41]

$$E = \int d\sigma_a (\partial_b h^{ab} - \partial^a h_b^b). \quad (3.22a)$$

However, the expression (3.22a) is non-covariant, and for the computation in an arbitrary coordinate system, even an asymptotically Cartesian, it may lead, as shown in [49], to an arbitrary value of the energy  $H_0$ .

The formula (3.21) yields the unambiguous result  $H_0 = M$ , which confirms the identification of the parameter  $M$  in the metric asymptotics (3.20) with the total mass of an isolated distribution of matter. However, one has to remember that the parameter  $M$  in the external Schwarzschild solution is not a pure mass of matter, but it also includes the gravitational mass defect or the own energy of the gravitational field.

Thus, in the framework of the general relativity theory on the basis of a correct gravitational action function (2.82), one can obtain a covariant definition of the energy of asymptotically flat worlds (in contrast to the opinion expressed in [49]). In a similar way, one can covariantize the expressions [43] for the total momentum and the angular momentum of an asymptotically flat gravitating system.

The result (3.22a) for the total energy of the gravitating system in an asymptotically flat world can be derived from the Landau-Lifshitz energy-momentum pseudotensor [27].

When using in the ADM procedure the additional conditions (3.12) with the functions  $f^\mu(t, \mathbf{x})$  that do not depend on time,

$$f^\mu = f^\mu(\mathbf{x}), \quad \dot{f}^\mu(\mathbf{x}) \equiv 0, \quad (3.23)$$

in view of (3.14) one obtains the physical Hamiltonian, which numerically coincides with (3.21), but is a functional of the independent degrees of freedom of the gravitating system

$$H_{\text{phys}}(t) [g^A, p_A] = \oint d\sigma^a \left( \partial^b g_{ab} - \partial_a g_{bb} \right) \Big|_{g_{ab} = g_{ab}[g^A, p_A]}. \quad (3.24)$$

This statement represents the contents of the Regge-Teitelboim theorem [51] about the physical Hamiltonian in the asymptotically flat worlds.

Since the physical Hamiltonian is determined by the spatial asymptotics of the three-dimensional metric satisfying (3.20), we can illustrate the ADM methods on an example of a linearized gravitational field describing small deviations of the metric from the flat one:

$$g_{ab} = \delta_{ab} + h_{ab}, \quad h_{ab} \ll 1.$$

Let us assume that the canonical transformation in (3.4) acts only on the phase variables of gravitational field  $(g_{ab}, p^{ab})$ . We transform them using the decomposition of an arbitrary symmetrical tensor  $f_{ab}$  into the longitudinal, transverse and transverse-traceless parts:

$$f_{ab} = \partial_a f_b + \partial_b f_a + f_{ab}^T + f_{ab}^{TT}. \quad (3.25)$$

Here the transverse part  $f_{ab}^T$  is expressed in terms of the trace  $f^T = f_{aa}^T$  using the projection operator

$$f_{ab}^T = \frac{1}{2} \left[ \delta_{ab} - \frac{\partial_a \partial_b}{\Delta} \right] f^T, \quad (3.26)$$

where  $1/\Delta$  is Green's function of the Laplace operator  $\Delta = \partial_a \partial_a$  in the open flat space. The trace of the transverse part  $f^T$  satisfies the equation

$$\Delta f^T = \Delta f_{aa} - \partial_a \partial_b f_{ab}. \quad (3.27)$$

Using (3.25)-(3.27), we transform the following expression:

$$\int dt \int d^3x p^{ab} \dot{g}_{ab} = \int dt \int d^3x \{ p_{TT}^{ab} \dot{g}_{ab}^{TT} - g_{ab}^T \dot{p}_T^{ab} - 2\partial_b p^{ab} \dot{g}_a \},$$

where  $g_a, g_{ab}^T, g_{ab}^{TT}$  are the components of the decomposition (3.25) of the three-dimensional metric  $g_{ab}$  (a similar notation is used for  $p^{ab}$ ). In the second term of this expression, the integration by parts with respect to time  $t$  was done. Taking into account that according to (3.26)  $g_{ab}^T$  is expressed in terms of  $g^T$ , we obtain

$$\int dt \int d^3x p^{ab} \dot{g}_{ab} = \int dt \int d^3x \left\{ p_{TT}^{ab} \dot{g}_{ab}^{TT} - g^T \frac{1}{2} \dot{p}_T - 2\partial_b p^{ab} \dot{g}_a \right\}, \quad (3.28)$$

from which it is obvious that instead of  $(g_{ab}, p^{ab})$  one can introduce the following canonically conjugated variables:

$$X^a = g_a, \quad \Pi_a = -2\partial_b p^{ab}, \quad (3.29)$$

$$X^0 = -\frac{1}{2} p_T, \quad \Pi_0 = g_T, \quad (3.30)$$

$$g^A = g_{ab}^{TT}, \quad p_A = p_{TT}^{ab}, \quad A = 1, 2. \quad (3.31)$$

Separating the linear part with respect to the fields in the gravitational constraints (2.125) and (2.126), we find

$$\begin{aligned} H_\perp &= \Delta g_{aa} - \partial_a \partial_b g_{ab} + h_\perp(h_{ab}, p^{ab}; \varphi, p), \\ H_a &= -2\partial_b p^{ba} + h_a(h_{ab}, p^{ab}; \varphi, p), \end{aligned}$$

where the terms  $h_\perp$  and  $h_a$  are of higher order with respect to the fields. In terms of the new phase variables (3.29)-(3.31), the constraints are recast into:

$$H_\perp = \Delta \Pi_0 + h_0[X, \Pi; g^A, p_A; \varphi, p], \quad (3.32)$$

$$H_a = \Pi_a + h_a[X, \Pi; g^A, p_A; \varphi, p],$$

where in terms of the new variables,  $t_0$  and  $t_a$  become functionals of the second and higher order in their arguments.

Eq. (3.32) shows that the matrix (3.7) in the linear approximation reads

$$J_{(\nu)}^{(\mu)}(\mathbf{x}, \mathbf{y}) = \begin{bmatrix} \Delta\delta(\mathbf{y} - \mathbf{x}) & 0 \\ 0 & \delta_b^a \delta(\mathbf{y} - \mathbf{x}) \end{bmatrix}. \quad (3.33)$$

Let us impose the additional conditions (3.11) satisfying (3.23):

$$X^0 = 0, \quad X^a = \frac{1}{2} x^a, \quad (3.34)$$

then the Hamiltonian (3.14) takes the form (3.24). We use this form as a volume integral [taking (3.27) into account]

$$H_0 = - \int d^3x \Delta g^T. \quad (3.35)$$

In the linear approximation, it is easy to obtain the dependence of this quantity on the physical degrees of freedom  $(g_{ab}^{TT}, p_{TT}^{ab}, \varphi, p)$ . When solving the constraint (3.32) in this approximation, we have

$$-\Delta g^T(\mathbf{x}) = h_0 [X^\mu, \Pi_\mu; g^A, p_A; \varphi, p] \Big|_{X^0=0, X^a=\frac{1}{2}x^a, \Pi_\mu=0}.$$

Taking into account (2.125) and (2.94), we can find  $t_0$ , from which the physical Hamiltonian of the gravitating system in the linear approximation is equal to

$$H_{\text{phys}}(t) [g_{ab}^{TT}, p_{TT}^{ab}, \varphi, p] = \int d^3x \left\{ (p_{TT}^{ab})^2 + \frac{1}{2} (\partial_a g_{bc}^{TT})^2 + \mathcal{H}_m(\varphi, p) \right\}. \quad (3.36)$$

In this approach, the purely gravitational degrees of freedom are identified with the two transverse-traceless components of the metric and the conjugated momenta. These are the two well-known transverse polarizations of the gravitational wave.

Let us now determine the lapse and shift functions. In view of (3.33) and the conditions (3.23) which are satisfied by the additional conditions (3.34), the equations for  $N^\mu$  take the following form (in the linear approximation):

$$\Delta N(\mathbf{x}) = 0, \quad N^a(\mathbf{x}) = 0. \quad (3.37)$$

The equation (3.37) should be supplemented with the boundary condition, which is induced by the asymptotically flat type of the metric

$$N \Big|_{r \rightarrow \infty} = 1,$$

consequently

$$N(\mathbf{x}) = 1 \quad (3.38)$$

with an accuracy up to the terms linear in the fields.

This result demonstrates how important are the boundary conditions in the cosmological problems with the open models. Indeed, the entire dynamics of the gravitating system and its energy are determined by the spatial asymptotics. From the physical point of view, one can explain this as follows. The time and space coordinates which are realized and measured using the physical instruments, can be uniquely determined only on the spatial asymptotics, since there are time-like and space-like Killing vector fields. Thus, the Hamiltonian (3.21) is a generator of the global translation in time, determined asymptotically.

### *Dynamics of closed cosmological models*

In contrast to the asymptotically flat worlds which were recently comprehensively studied, the closed cosmological models and their dynamics remain a stumbling block in the modern gravity theory.

In closed cosmological systems, the surface integral  $H_0$  is absent, and therefore the physical Hamiltonian is determined only by the second term in (3.14) and it does not vanish only if  $\dot{f}^\mu(t, \mathbf{x}) \neq 0$ . The conditions (3.16) for the lapse and shift functions also require that not all  $\dot{f}^\mu(t, \mathbf{x})$  vanish.

Therefore, in order to have a nontrivial dynamics in the closed cosmological models one needs to introduce the canonical gauges that explicitly depend on time. Such a time-dependence of the gauge generates the dynamics of all the physical variables of the theory. Alternatively, one can introduce the evolution of the closed cosmology by fixing the Lagrange multipliers – the lapse and shift functions, avoiding the use of explicitly time-dependent functions. However, this is at most another way to construct the canonical gauge with a parametric time dependence.

We can demonstrate this by the simplest example of the closed Friedman Universe described by the metric

$$ds^2 = -N^2(t) dt^2 + R^2(t) \gamma_{ik} dx^i dx^k, \quad (3.39)$$

where the lapse function  $N(t)$  and the cosmological radius  $R(t)$  depend only on time, and  $\gamma_{ik}$  is the metric of the three-dimensional hypersphere of a unit radius in some coordinates  $x^i$ . Since the model is homogeneous, its Lagrange function  $L_g$  in terms of the finite number of degrees of freedom is obtained by the integration of (2.84) over the three-dimensional volume of the Universe:

$$L_g = \frac{1}{2l_0} \left\{ -l_0^2 \frac{a\dot{a}}{N} + Na \right\}, \quad (3.40)$$

where  $a$  is the dimensionless radius of the Universe in the units of the Planck length  $l_0 = \sqrt{2G\hbar/3\pi c^3}$ ,

$$R = l_0 a. \quad (3.41)$$

The superhamiltonian corresponding to (3.40) is equal to

$$\int d^3x \mathcal{H}(\mathbf{x}) = -\frac{\varepsilon_0}{2} \left\{ \frac{p^2}{a} + a \right\}, \quad (3.42)$$

where  $p$  is the momentum conjugated to the variable  $a$ :

$$p = \frac{\partial L_g}{\partial \dot{a}} = -\frac{l_0}{N} a \dot{a}, \quad (3.43)$$

and  $\varepsilon_0 = l_0^{-1}$  is the Planck energy ( $\varepsilon_0 = \hbar c/l_0$  in the usual units).

We assume that the Universe is homogeneously filled with a noninteracting dust without pressure, so that the integral over the three-dimensional volume of the Universe from the matter superhamiltonian  $\mathcal{H}_\varphi$  in (2.125) is equal to its conserved mass-energy

$$\int d^3x \mathcal{H}_\varphi(\mathbf{x}) = \varepsilon, \quad \varepsilon = \text{const.}$$

Then the geometrodynamical constraint takes the following form:

$$H_\perp = 0, \quad (3.44)$$

$$H_\perp = \int d^3x H_\perp(\mathbf{x}) = \frac{\varepsilon_0}{2} \left( \frac{p^2}{a} + a \right) + \varepsilon. \quad (3.45)$$

Because of the homogeneity of the Friedman model, other constraints are not important since they are satisfied identically due to the invariance of the system with respect to the spatial translations.

Therefore, a further analysis is equivalent to the ADM procedure with one constraint in which the index  $\mu$  takes only one value.

The equations (3.1) for the complete set of the phase variables  $\Phi = (a, p)$  reduce to

$$\dot{p} = -\frac{N\varepsilon_0}{2} \left( \frac{p^2}{a^2} - 1 \right),$$

and (3.43). The constraint equation (3.44) is the first integral of these equations, which corresponds to the condition of preservation of the constraint in time.

The evolution of the Friedman Universe is usually found by choosing the coordinate condition

$$N = R, \quad (3.46)$$

that brings the metric (3.39) to the conformally static form. Combining (3.43) and (3.46) we have  $p = -\dot{a}$ , as a result, the constraint (3.44) becomes an equation for  $a(t)$ , the solution of which gives

$$a(t) = \frac{\varepsilon}{\varepsilon_0} (1 - \cos t), \quad (3.47)$$

where the moment of time  $t = 0$  corresponds to the cosmological singularity  $a = 0$ .

Let us show that the gauge (3.46), containing the Lagrange multiplier, corresponds to a canonical gauge that explicitly depends on time. For this, we choose the new canonical variables in accordance with the ADM procedure (3.4)  $\Pi_0 = \Pi, X^0 = T$ :

$$a = \sqrt{-2\Pi} \cos T, \quad (3.48)$$

$$p = \sqrt{-2\Pi} \sin T. \quad (3.49)$$

The constraint (3.45) in new variables is recast into the form

$$H_{\perp} = -\frac{\varepsilon_0}{2} \frac{\sqrt{-2\Pi}}{\cos T} + \varepsilon, \quad (3.50)$$

and then a solution (3.8) of this constraint with respect to  $\Pi$  reads

$$\Pi = -P, \quad P \equiv 2 \left( \frac{\varepsilon}{\varepsilon_0} \right)^2 \cos^2 T. \quad (3.51)$$

Imposing now the additional condition (3.11)

$$T = \frac{t - \pi}{2}, \quad (3.52)$$

we find the physical Hamiltonian of the system, see (3.14)

$$H_{\text{phys}}(t) = \frac{\varepsilon}{\varepsilon_0^2} \sin^2(t/2).$$

The dynamically independent degrees of freedom of this Hamiltonian are the variables of the dust-like matter, with the energy  $\varepsilon$  being the function of them. Substituting (3.51) into (3.48), we obtain (3.47) again, that is, the canonical gauge (3.52) which can be rewritten in the original phase variables as

$$a + p \tan(t/2) = 0, \quad (3.53)$$

is equivalent to the coordinate condition (3.46). One can verify this using (3.15):

$$N = \left( \frac{\partial H_{\perp}}{\partial \Pi} \right)^{-1} \dot{T} = l_0 a. \quad (3.54)$$

Therefore, the choice of a one-parameter family of surfaces in the phase space (3.46) or (3.53) determines the family of the space-like hypersurfaces in the four-dimensional spacetime, and the dynamics of the system reduces to the motion along this family with the growth of the time parameter  $t$ . The only difference of fixing the gauge (3.46) from (3.53) is that in (3.46) the family of hypersurfaces is determined not globally, but differentially, by fixing not a one-parameter slicing itself, but rather its derivative with respect to the parameter  $t$ . All this is directly extended to the general case of a non-homogeneous gravitating closed cosmological system, but the choice of a system of space-like hypersurfaces and their coordinate parametrization will be determined by the family of the four canonical gauges  $\chi^{\mu}(t, \Phi) = 0$ , explicitly depending on the time parameter  $t$ .

Moreover, each choice of the gauges, i.e., of the functions  $f^{\mu}(t, \mathbf{x})$  in (3.14) will produce its own Hamiltonian. An ambiguity of determination of the energy in the closed world, in contrast to an asymptotically flat one, is caused by the fact that in closed Universes there are no a preferred observer and a preferred system of the space-like hypersurfaces, which in open systems are associated with the spacetime flat asymptotics.

# 4

## Torsion effects on the structure and evolution of gravitating systems

### 4.1. Matter fields in the Einstein-Cartan theory

As we demonstrated in Sec. 1.3., the spin-connection interaction does not have an explicit dynamical nature in the Einstein-Cartan theory (ECT). Nevertheless, it modifies the structure of the matter sources of the gravitational field, and thereby the spin of matter affects the dynamics of the metric. Furthermore, it is worthwhile to stress that the ETC is not competing with GR since the ( $\Gamma$ - $S$ )-interaction has a microscopic nature and is manifest on a scale of the order of the mean distance between particles. The macroscopic gravitational theory is obtained as a result of an appropriate averaging process. The value of the term quadratic in spin  $T_{\mu\nu}^{\text{eff}}$  becomes comparable with  $T_{\mu\nu}$  at densities  $\rho \geq \rho_{\text{cr}} = \frac{m^2 c^4}{\hbar^2 G}$  of the spinning matter built of particles with the mass  $m$  [7]. For the mass of a nucleon, the critical density  $\rho_{\text{cr}} \approx 10^{57}$  kg/m<sup>3</sup> is much smaller than the Planck density  $\rho_0 \sim 10^{97}$  kg/m<sup>3</sup> at which the quantum-gravitational effects start dominating. Consequently, the torsion can be essential already at the level of the classical theory of the gravitational interactions.

Let us begin our study of the torsion effects in ECT with the analysis of the structure of the matter currents.

#### *Weyssenhoff spinning fluid*

Before we present a consistent variational theory, it is instructive to show that one can include a fluid with spin into the dynamical scheme of ECT on the

basis of covariant conservation laws of the energy-momentum and spin (1.45) and (1.47). Our presentation follows [191].

The ideal fluid with spin (or a spinning fluid) is a continuous medium, each element of which is characterized by the energy, momentum and an intrinsic angular momentum. The phenomenological model of such a medium was developed by Weyssenhoff and Raabe [54] in the flat Minkowski spacetime. Accordingly, the general-relativistic ideal fluid with spin will also be called a Weyssenhoff fluid.

In the phenomenological approach, elements of a medium are described by the (average) 4-velocity  $u^\mu$ , the 4-vector  $P_\mu$  of the energy-momentum density, and by the density of the spin angular momentum  $S_{\mu\nu} = S_{[\mu\nu]}$ . The spatial components of the latter constitute the 3-vector  $\mathbf{S} = \{S_{23}, S_{31}, S_{12}\}$  which in the rest frame of the fluid is equal to the 3-dimensional density of intrinsic angular momentum. The remaining components also constitute the 3-vector  $\mathbf{q} = \{S_{10}, S_{20}, S_{30}\}$ . We assume that this vector vanishes in the rest frame of fluid, i.e.,

$$S_{\mu\nu}u^\nu = 0. \quad (4.1)$$

This assumption is called a Frenkel supplementary condition.

The 4-velocity is a timelike vector which is normalized by  $u_\mu u^\mu = -1$ . The projection of the 4-momentum on the rest frame of the fluid yields the energy density  $\varepsilon = -u^\mu P_\mu$ .

As a first step, we consider the spinning dust on non-interacting elements with the vanishing pressure. The phenomenological postulate for the canonical energy-momentum tensor and the canonical spin tensor of a dust reads as follows:

$$\left. \begin{aligned} t^\lambda{}_\mu &= u^\lambda P_\mu, \\ S^\lambda{}_{\mu\nu} &= u^\lambda S_{\mu\nu}. \end{aligned} \right\} \quad (4.2)$$

Making use of the covariant conservation law of the angular momentum (1.47), we find a relation between the densities of the energy-momentum and spin

$$u_\mu P_\nu - u_\nu P_\mu = 2c(\nabla_\sigma - 2Q_\sigma)(u^\sigma S_{\mu\nu}). \quad (4.3)$$

Contracting this equation with the 4-velocity, we find the 4-vector of momentum explicitly

$$P_\mu = \varepsilon u_\mu + 2u^\lambda c \dot{S}_{\mu\lambda}. \quad (4.4)$$

We introduced a convenient notation for the derivative  $\dot{\varphi}_A := (\nabla_\sigma - 2Q_\sigma)(u^\sigma \varphi_A)$  for an arbitrary tensor density  $\varphi_A$ . Substituting (4.4) back into (4.3), we recast the latter into the equation of motion of spin

$$\dot{S}_{\mu\nu} = u_\mu u^\lambda \dot{S}_{\nu\lambda} - u_\nu u^\lambda \dot{S}_{\mu\lambda}. \quad (4.5)$$

Finally, making use of (4.4) in (4.2), we obtain the canonical energy-momentum tensor of the Weyssenhoff spinning dust

$$t_{\mu\nu} = \varepsilon u_\mu u_\nu + 2u_\mu u^\lambda c \dot{S}_{\nu\lambda}.$$

The first term on the right-hand side represents the usual ideal fluid with the dust equation of state, and the last term is the spin contribution.

The phenomenological theory can be upgraded to a complete dynamical theory of a spinning fluid. In order to construct the Lagrangian, we need to choose the appropriate variables which describe such a continuous medium. In the classical continuum mechanics, the ordinary matter consists of the structureless elements - the material points without physical properties other than mass. In contrast, the spinning fluid represents an example of a matter with microstructure, the elements of which have additional degrees of freedom such as spin.

Following Cosserat brothers [25], to describe the dynamics of a medium with microstructure we attach a triad of vectors with each matter element. The field of such material triads is denoted  $b_i^\mu$ ,  $i = 1, 2, 3$ , and together with the 4-velocity it comprises a local frame  $b_a^\mu$ ,  $a = 0, 1, 2, 3$ ,

$$b_a^\mu = \{b_0^\mu = u^\mu, b_1^\mu, b_2^\mu, b_3^\mu\}$$

defined at all the spacetime points of the domain occupied by the matter. The vectors of the triad are chosen orthogonal to each other and to  $u^\mu$  and have the unit length:

$$g_{\mu\nu} b_a^\mu b_b^\nu = \eta_{ab} = \text{diag}(-1, +1, +1, +1).$$

Such an orthonormal tetrad is called a material frame and it is different from the spacetime tetrad  $h_a^\mu$  which we discussed in Sec. 1.1.. We can choose  $h_a^\mu$  arbitrarily and can always change this tetrad using a local Lorentz rotation. In contrast, the material tetrad  $b_a^\mu$  cannot be chosen arbitrarily, it is rigidly fixed and moves together with the fluid.

We are now in a position to formulate the variational principle for the spinning fluid. We begin by postulating (as in the usual case of an ideal fluid) that the dynamics of the medium is such that the number of particles is constant, and the entropy  $s$  and the identity of particles are conserved along the flow lines of the fluid. Mathematically, this means that we impose the constraints

$$\partial_\mu (\sqrt{g} \rho u^\mu) = 0, \quad u^\mu \partial_\mu s = 0, \quad u^\mu \partial_\mu X = 0.$$

Here  $\rho$  is the particle number density, and the Lin variable  $X$  is introduced to identify matter elements.

We will describe the physical properties of the fluid by the particle density  $\rho$ , the internal energy density  $\varepsilon$ , and the specific spin density  $\mu^{ij} = -\mu^{ji}$  (i.e., the spin density per matter element),  $i, j = 1, 2, 3$ .

With all these prerequisites, the Lagrangian of the spinning fluid reads

$$\begin{aligned} \mathcal{L}_W = & -\varepsilon(\rho, s, \mu^{ij}) + \frac{1}{2} \rho \mu^{ij} g_{\mu\nu} b_i^\mu u^\alpha \nabla_\alpha b_j^\nu \\ & + \rho u^\mu \partial_\mu \lambda_1 + \lambda_2 u^\mu \partial_\mu s + \lambda_3 u^\mu \partial_\mu X + \lambda^{ab} (g_{\mu\nu} b_a^\mu b_b^\nu - \eta_{ab}). \end{aligned}$$

The first line represents the physically essential part related to the kinetic and potential energy of the system, and the second line takes into account all the

constraints which are imposed on the system by means of the Lagrange multipliers  $\lambda_1, \lambda_2, \lambda_3$  and  $\lambda^{ab}$ .

The thermodynamical properties of the spinning fluid are described by the usual Gibbs law corrected by the spin energy contribution:

$$Tds = d\left(\frac{\varepsilon}{\rho}\right) + pd\left(\frac{1}{\rho}\right) - \frac{1}{2}\omega_{ij}d\mu^{ij}.$$

Here  $T$  is the temperature,  $p$  is the pressure, and  $\omega_{ij}$  is the thermodynamical variable conjugated to the specific spin density  $\mu^{ij}$ .

The equations of motion are derived from the variation of the matter action  $\frac{1}{c} \int d^4x \sqrt{g} \mathcal{L}_W$  with respect to the total set of (fundamental and auxiliary) variables  $\Phi^A = \{\rho, s, X, \mu^{ij}, b_a^\mu, \lambda_1, \lambda_2, \lambda_3, \lambda^{ab}\}$ . Variation with respect to the Lagrange  $\lambda$  multipliers, we obtain the set of constraints, whereas the variation with respect to  $\rho, s, X$ , and  $\mu^{ij}$  yields, respectively:

$$\begin{aligned} -\varepsilon - p + \frac{1}{2}\rho\mu^{ij}g_{\mu\nu}b_i^\mu u^\lambda \nabla_\lambda b_j^\nu + \rho u^\mu \partial_\mu \lambda_1 &= 0, \\ \rho T + \dot{\lambda}_2 = 0, \quad \dot{\lambda}_3 = 0, \quad \omega_{ij} &= g_{\mu\nu}u^\alpha b_i^\mu \nabla_\alpha b_j^\nu. \end{aligned}$$

Here we had to make use of the Gibbs law to evaluate the derivatives of the internal energy density  $\varepsilon$  with respect to its arguments.

Finally, from the variation of the material tetrad,  $u^\mu$  and  $b_i^\mu$ , we find

$$\begin{aligned} \frac{1}{2}\rho\mu^{ij}g_{\rho\sigma}b_i^\rho \nabla_\mu b_j^\sigma + \rho\partial_\mu \lambda_1 + \lambda_2\partial_\mu s + \lambda_3\partial_\mu X + 2\lambda^{0a}g_{\mu\nu}b_a^\nu &= 0, \\ g_{\mu\nu}\left(\rho\mu^{ij}u^\lambda \nabla_\lambda b_j^\nu + \frac{1}{2}b_j^\nu u^\lambda \partial_\lambda \mu^{ij} + 2\lambda^{ia}b_a^\nu\right) &= 0. \end{aligned}$$

Contracting these equations with  $u^\mu$  and  $b_k^\mu$ , we find the Lagrange multipliers

$$2\lambda^{00} = \varepsilon + p, \quad 2\lambda^{0i} = \rho\mu^{ij}u_\nu u^\alpha \nabla_\alpha b_j^\nu, \quad 2\lambda^{ik} = -\rho\mu^{(i|j|}b_{\nu}^{k)}u^\alpha \nabla_\alpha b_j^\nu,$$

and obtain the equations of motion of spin:

$$u^\alpha \partial_\alpha \mu^{ij} + \mu^{ik}\omega^j_k + \mu^{kj}\omega^i_k = 0.$$

The Latin 3-dimensional indices that refer to the material tetrad are moved with the help of the Euclidean metric  $\delta^{ij}$  and  $\delta_{ij}$ . We also introduced the inverse material triad by  $b_\mu^i = g_{\mu\nu}\eta^{ij}b_j^\nu = g_{\mu\nu}\delta^{ij}b_j^\nu$ . One can verify the relations

$$b_\mu^j b_i^\mu = \delta_i^j, \quad b_\nu^i b_i^\mu = \delta_\nu^\mu + u^\mu u_\nu.$$

Finally, we apply these results to derive the canonical tensors of energy-momentum and spin. Recalling the definitions of Sec. 1.3., we find from the Weyssenhoff Lagrangian  $\mathcal{L}_W$ :

$$\left. \begin{aligned} t^\lambda_\mu &= p\delta_\mu^\lambda + (p + \varepsilon)u^\lambda u_\mu + 2u^\lambda u^\nu c\dot{S}_{\mu\nu}, \\ S^\lambda_{\mu\nu} &= u^\lambda S_{\mu\nu}, \end{aligned} \right\} \quad (4.6)$$

where we have the spin density explicitly

$$S^{\mu\nu} = \frac{1}{2}\rho\mu^{ij}b_i^\mu b_j^\nu.$$

By construction, the spin tensor satisfies the Frenkel condition (4.1). The canonical sources (4.6) provide a consistent generalization of the phenomenological result (4.2) to the case of fluid with nontrivial pressure  $p$ .

This completes the construction of the variational theory of the Weyssenhoff spinning fluid. It is applicable to any gravitational theory.

Let us now specialize to the Einstein-Cartan gravity with the Lagrangian (1.34) and the field equations (1.48), (1.49). Combining last equation with the Frenkel condition (4.1), we find the torsion  $Q^\lambda{}_{\mu\nu} = \kappa c u^\lambda S_{\mu\nu}$  (the torsion trace vanishes,  $Q_\mu = 0$ , since  $S_\mu = u^\nu S_{\mu\nu} = 0$ ). Plugging this into (1.51), we find the effective Lagrangian of the spinning fluid

$$\begin{aligned} \mathcal{L}_W^{\text{eff}} = & -\varepsilon(\rho, s, \mu^{ij}) + \frac{1}{2}\rho\mu^{ij}g_{\mu\nu}b_i^\mu u^\alpha \overset{\{\}}{\nabla}_\alpha b_j^\nu + \frac{\kappa c^2}{8}\rho^2\mu^{ij}\mu_{ij} \\ & + \rho u^\mu \partial_\mu \lambda_1 + \lambda_2 u^\mu \partial_\mu s + \lambda_3 u^\mu \partial_\mu X + \lambda^{ab}(g_{\mu\nu}b_a^\mu b_b^\nu - \eta_{ab}). \end{aligned}$$

The last term in the first line describes the typical spin-spin contact interaction. One can also write this as

$$\frac{\kappa c^2}{8}\rho^2\mu^{ij}\mu_{ij} = \kappa c^2 S^2, \quad S^2 := \frac{1}{2}S^{\mu\nu}S_{\mu\nu},$$

but this form produces a misleading impression that the spacetime metric tensor is involved. In reality, the effective spin-spin interaction term depends only on the particle density  $\rho$  and on the specific spin density  $\mu^{ij}$ . This is important to keep in one's mind when performing the variation of the effective fluid action with respect to the metric.

Taking into account that the variation of the Christoffel symbols reads

$$\delta\{\overset{\lambda}{\mu\nu}\} = \frac{1}{2}g^{\lambda\alpha}\left(\overset{\{\}}{\nabla}_\mu\delta g_{\nu\alpha} + \overset{\{\}}{\nabla}_\nu\delta g_{\mu\alpha} - \overset{\{\}}{\nabla}_\alpha\delta g_{\mu\nu}\right),$$

it is straightforward to find the effective metrical energy-momentum tensor

$$\left. \begin{aligned} T_{\mu\nu}^{\text{eff}} = & p^{\text{eff}}g_{\mu\nu} + (p^{\text{eff}} + \varepsilon^{\text{eff}})u_\mu u_\nu + 2c(u^\alpha u^\beta - g^{\alpha\beta})\overset{\{\}}{\nabla}_\alpha(u_{(\mu}S_{\nu)\beta}), \\ \varepsilon^{\text{eff}} = & \varepsilon - \kappa c^2 S^2, \quad p^{\text{eff}} = p - \kappa c^2 S^2. \end{aligned} \right\} \quad (4.7)$$

Summarizing, the structure of the sources of the gravitational field in the Riemann-Cartan spacetime with torsion for the Weyssenhoff spinning fluid is given by (4.6). For the case of the Einstein-Cartan theory, the dynamics of the gravitational field is described by Einstein's equation (1.52) with the effective energy-momentum tensor (4.7) that depends on the spin density square and the derivatives of the spin tensor.

The phenomenological way to introduce the spin in the fluid model was discussed in the numerous papers, and a number of variational principle approaches for the spinning fluid were developed, see [12], for example.

### *Spinor fields in ECT*

Of a particular interest is the study of the self-gravitating fields with spin which due to the contact character of spin-spin interaction of the Einstein-Cartan model gives rise to a nonlinear extension of GR expressed by the effective equation (1.52). The spin-spin interaction caused by space-time torsion is manifest as a nonlinearity of these fields, as we demonstrate below.

We consider here the most important physical example of the spinor Dirac field. As the Lagrangian of the Dirac field  $\Psi$  in the space  $U_4$ , we take

$$\mathcal{L}_D(\Psi, \bar{\Psi}, \omega_\mu, g_{\mu\nu}) = \frac{\hbar c}{2} (\bar{\Psi} \gamma^a D_a \Psi - D_a \bar{\Psi} \gamma^a \Psi) + mc^2 \bar{\Psi} \Psi. \quad (4.8)$$

Here  $m$  is the rest mass of the Dirac fermion  $\Psi$ ,  $\gamma^a$  are the flat Dirac matrices in Weinberg's representation [192], and  $D_a = h_a^\mu D_\mu$  with

$$D_\mu \Psi = \partial_\mu \Psi + \omega_\mu \Psi.$$

The expression for the covariant derivatives of spinors is defined by formulas (A2.7)-(A2.8) of Appendix A2.

As independent dynamical variables, one can take  $\{\bar{\Psi}, \Psi, g_{\mu\nu}, \Gamma^\lambda_{\mu\nu}\}$ , on which the metricity condition (1.8) is imposed as a constraint. If the latter is solved explicitly, the connection  $\Gamma^\lambda_{\mu\nu}$  takes the form (1.12), and the spinor connection (A2.6) can be recast into

$$\omega_\mu = \frac{1}{4} \gamma^\nu \overset{\{\}}{\nabla}_\mu \gamma_\nu - \frac{1}{4} \gamma^\nu T^\lambda_{\nu\mu} \gamma_\lambda. \quad (4.9)$$

However, since the tetrad field is fundamentally involved in the definition of the spinor connection and the covariant derivative, it is more convenient to switch to the tetrad formalism and to take the equivalent set of independent dynamical variables  $\{\bar{\Psi}, \Psi, h_\mu^a, \Gamma^a_{b\mu}\}$ . The metricity condition (1.8) in the tetrad formalism reads

$$\nabla_\mu \eta_{ab} = -\Gamma_{ab\mu} - \Gamma_{ba\mu} = 0.$$

Consequently, we conclude that the metric-compatible local Lorentz connection is skew-symmetric:  $\Gamma_{ab\mu} = \Gamma_{[ab]\mu}$ . The same property then shares the tetrad curvature tensor:  $R_{ab\mu\nu} = R_{[ab]\mu\nu}$ .

The field equations of the theory are derived from the variational problem for the total Lagrangian (1.34) in the tetrad disguise

$$S = \int d^4x h \left\{ \frac{1}{2\kappa c} R^{ab}{}_{\mu\nu} h_a^\mu h_b^\nu + \frac{1}{c} \mathcal{L}_D \right\}. \quad (4.10)$$

Here we denoted the determinant of the tetrad  $h := \det h_\mu^a$ ; obviously we have  $h = \sqrt{g}$ . In the tetrad formalism we do not need to add the metricity condition with the Lagrange multipliers, it is sufficient just to take the skew-symmetric local Lorentz connection  $\Gamma^a_{b\mu}$ .

Variation of the action (4.10) with respect to the independent variables yields:

$$\frac{\delta S}{\delta h_a^\mu} = \frac{1}{\kappa c} (R^a{}_\mu - \frac{1}{2} h_\mu^a) - \frac{\mathcal{L}_D}{c} h_\mu^a - \frac{\hbar}{2} (D_\mu \bar{\Psi} \gamma^a \Psi - \bar{\Psi} \gamma^a D_\mu \Psi) = 0, \quad (4.11)$$

$$\frac{\delta S}{\delta \Gamma_{ab}{}^\mu} = -\frac{2}{\kappa c \hbar} \mathcal{D}_\nu (h h_{[a}^\nu h_{b]}^\mu) + \frac{\hbar}{8} \bar{\Psi} \{ \gamma^\mu \gamma_{[a} \gamma_{b]} + \gamma_{[a} \gamma_{b]} \gamma^\mu \} \Psi = 0, \quad (4.12)$$

$$\frac{\delta S}{\delta \bar{\Psi}} = \hbar \gamma^a (D_a - Q_a) \Psi + m c \Psi = 0, \quad (4.13)$$

$$\frac{\delta S}{\delta \Psi} = -\hbar (D_a - Q_a) \bar{\Psi} \gamma^a + m c \bar{\Psi} = 0. \quad (4.14)$$

Here the covariant derivative  $\mathcal{D}_\mu$  acts only on the tetrad indices. For example,  $\mathcal{D}_\mu v_a^\nu = \partial_\mu v_a^\nu - \Gamma_{a\mu}^b v_b^\nu$  for an arbitrary tensor object that carries both (world and tetrad) types of indices. Directly from the definition of the torsion, one can verify the useful geometrical identity

$$\frac{2}{\hbar} \mathcal{D}_\nu (h h_{[a}^\nu h_{b]}^\mu) \equiv Q^\mu{}_{ab} + h_a^\mu Q_b - h_b^\mu Q_a. \quad (4.15)$$

Making use of the definitions of Sec. 1.3., we find the canonical energy-momentum and spin tensors

$$t^\lambda{}_\mu = \mathcal{L}_D \delta_\mu^\lambda + \frac{\hbar c}{2} (D_\mu \bar{\Psi} \gamma^\lambda \Psi - \bar{\Psi} \gamma^\lambda D_\mu \Psi),$$

$$S^\lambda{}_{\mu\nu} = \frac{\hbar}{8} \bar{\Psi} \{ \gamma^\lambda \gamma_{[\mu} \gamma_{\nu]} + \gamma_{[\mu} \gamma_{\nu]} \gamma^\lambda \} \Psi = \frac{i\hbar}{4} \varepsilon_{\mu\nu}{}^{\lambda\sigma} \bar{\Psi} \gamma_\sigma \gamma_5 \Psi.$$

Since the spin tensor of the Dirac field is completely antisymmetric, the torsion trace vanishes  $Q_\mu = 0$  and hence the Dirac equation (4.13) in  $U_4$  is simplified,

$$\hbar \gamma^a D_a \Psi + m c \Psi = 0, \quad (4.13a)$$

and a similar result is found for the conjugated equation (4.14).

With the help of (4.15), the Palatini equation (4.12) determines the torsion

$$Q^\lambda{}_{\mu\nu} = \kappa c S^\lambda{}_{\mu\nu} = \frac{i\kappa\hbar c}{4} \varepsilon_{\mu\nu}{}^{\lambda\sigma} \bar{\Psi} \gamma_\sigma \gamma_5 \Psi. \quad (4.16)$$

This means that in the irreducible decomposition (1.13) only the pseudotrace of torsion is nontrivial,  $\check{Q}_\mu = \frac{i\kappa\hbar c}{4} \bar{\Psi} \gamma_\mu \gamma_5 \Psi$ .

It is straightforward to see that the fermion Lagrangian (4.8) vanishes “on-shell”,  $\mathcal{L}_D = 0$ , when the spinor fields that satisfy the Dirac equation (4.13a). As a result, Einstein’s equation (4.11) reads (converting the tetrad indices into the world ones):

$$G_{\mu\nu}(\Gamma) = \kappa t_{\mu\nu} = \frac{\kappa\hbar c}{2} (D_\nu \bar{\Psi} \gamma_\mu \Psi - \bar{\Psi} \gamma_\mu D_\nu \Psi). \quad (4.17)$$

The system (4.13), (4.13a), (4.16), and (4.17) represents the complete set of the dynamical equations for the Dirac field  $\Psi, \bar{\Psi}$  interacting with the gravitational field  $g_{\mu\nu}, \Gamma^\lambda{}_{\mu\nu}$ .

Using (4.9) and (4.16), we recast the Dirac equation (4.13a) into

$$\gamma^a \overset{\{\}}{D}_a \Psi + l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) \gamma^a \gamma_5 \Psi + \frac{mc}{\hbar} \Psi = 0, \quad (4.18)$$

where  $l_0^2 = \frac{3}{8} \kappa \hbar c$ , i.e.,  $l_0 \approx 4.95 \times 10^{-35}$  m. We thus conclude that the torsion induces a  $\Psi^4$ -nonlinearity in the Dirac equation (4.18).

Combining the general formula for the Riemann-Cartan connection (1.12) with (4.16), we can decompose the Einstein tensor into the Riemannian and post-Riemannian parts:

$$G_{(\mu\nu)}(\Gamma) = G_{\mu\nu} + \left( \frac{\kappa \hbar c}{4} \right)^2 [g_{\mu\nu} (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) + 2 (\bar{\Psi} \gamma_\mu \gamma_5 \Psi) (\bar{\Psi} \gamma_\nu \gamma_5 \Psi)]. \quad (4.19)$$

We need only the symmetric part, since the skew-symmetric part of (4.17) is satisfied due to the conservation law of the total angular momentum. The validity of the latter can be directly verified using the field equation (4.13) and the expressions for the canonical energy-momentum and spin.

As we described in Sec. 1.3., the dynamics of the system is determined by the effective Lagrangian  $\frac{1}{2\kappa c} R + \frac{1}{c} \mathcal{L}_D^{\text{eff}}$ , where the effective Lagrangian for the Dirac field takes the form

$$\mathcal{L}_D^{\text{eff}} = \frac{\hbar c}{2} \left\{ \bar{\Psi} \gamma^a \overset{\{\}}{D}_a \Psi - \overset{\{\}}{D}_a \bar{\Psi} \gamma^a \Psi + l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\} + mc^2 \bar{\Psi} \Psi. \quad (4.20)$$

The gravitational field equation (4.17) then reduces to the Einstein equation (1.52) with the effective energy-momentum tensor

$$T_{\mu\nu}^{\text{eff}} = \frac{\hbar c}{2} \left\{ \overset{\{\}}{D}_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} \overset{\{\}}{D}_{\nu)} \Psi - g_{\mu\nu} l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\}. \quad (4.21)$$

Now let us consider a nonlinear spinor field of the Ivanenko-Heisenberg [56] type as a source of the gravitational field in the framework of the Einstein-Cartan theory. Its generally covariant Lagrangian reads

$$\mathcal{L}_{IH} = \frac{\hbar c}{2} \left\{ \bar{\Psi} \gamma^a D_a \Psi - D_a \bar{\Psi} \gamma^a \Psi \mp \lambda_p^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\}, \quad (4.22)$$

where  $\lambda_p$  is the coupling constant with the dimension of length.

The nonlinear spinor Ivanenko-Heisenberg equation in the Riemann-Cartan spacetime  $U_4$  follows from the Lagrangian (4.22):

$$\gamma^\mu D_\mu \Psi \mp \lambda_p^2 (\Psi^\dagger \gamma_a \gamma_5 \Psi) \gamma^a \gamma_5 \Psi = 0.$$

The canonical tensor of spin coincides with that of the Dirac field, and the torsion is again determined by the Palatini equation (4.16). Substituting the torsion, we recast the nonlinear spinor equation into the form similar to (4.18):

$$\gamma^\mu \overset{\{\}}{D}_\mu \Psi + (l_0^2 \mp \lambda_p^2) (\bar{\Psi} \gamma_a \gamma_5 \Psi) \gamma^a \gamma_5 \Psi = 0. \quad (4.23)$$

If we fix the coupling constants so that  $\lambda_p^2 = l_0^2$  and choose the upper sign in the last term of (4.23), the latter becomes the field equation of a Dirac neutrino in the Riemannian spacetime

$$\gamma^\mu \overset{\{\}}{D}_\mu \Psi = 0. \quad (4.24)$$

The metrical energy-momentum tensor coincides with the symmetric part of the canonical energy-momentum tensor, and for the nonlinear spinor field with the coupling constant  $\lambda_p^2 = l_0^2$  we derive

$$T_{\mu\nu} = t_{(\mu\nu)} = \frac{\hbar c}{2} \left\{ D_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi + g_{\mu\nu} l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\}. \quad (4.25)$$

Decomposing this into the Riemannian and post-Riemannian parts and combining with (4.19), we arrive at the Einstein equation (1.52) with the effective energy-momentum tensor

$$T_{\mu\nu}^{\text{eff}} = \frac{\hbar c}{2} \left\{ \overset{\{\}}{D}_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} \overset{\{\}}{D}_{\nu)} \Psi \right\}. \quad (4.26)$$

The whole dynamics of the coupled spinor and gravitational fields is described by the corresponding effective Lagrangian

$$\mathcal{L}_{IH}^{\text{eff}} = \frac{\hbar c}{2} \left\{ \bar{\Psi} \gamma^a \overset{\{\}}{D}_a \Psi - \overset{\{\}}{D}_a \bar{\Psi} \gamma^a \Psi \right\}.$$

We can formulate the conclusions that follow from our observations in the form of the two equivalence theorems as follows.

The Einstein-Cartan theory of the neutrino (massless Dirac spin- $\frac{1}{2}$  fermion) field in the Riemann-Cartan spacetime

$$\begin{aligned} G_{(\mu\nu)}(\Gamma) &= \frac{\kappa \hbar c}{2} \left\{ D_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi \right\}, \\ \gamma^a D_a \Psi &= 0, \end{aligned}$$

is equivalent to the Einstein theory of the nonlinear spinor (of the Ivanenko-Heisenberg type) field in the Riemannian spacetime:

$$\begin{aligned} G_{(\mu\nu)} &= \frac{\kappa \hbar c}{2} \left\{ \overset{\{\}}{D}_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} \overset{\{\}}{D}_{\nu)} \Psi - g_{\mu\nu} l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\}, \\ \gamma^a \overset{\{\}}{D}_a \Psi + l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) \gamma^a \gamma_5 \Psi + \frac{mc}{\hbar} \Psi &= 0. \end{aligned}$$

In a similar way, under the condition  $\lambda_p^2 = l_0^2$ , the Einstein-Cartan theory of the nonlinear spinor field in the Riemann-Cartan spacetime

$$\begin{aligned} G_{(\mu\nu)}(\Gamma) &= \frac{\kappa \hbar c}{2} \left\{ D_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} D_{\nu)} \Psi + g_{\mu\nu} l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) (\bar{\Psi} \gamma^a \gamma_5 \Psi) \right\}, \\ \gamma^a D_a \Psi - l_0^2 (\bar{\Psi} \gamma_a \gamma_5 \Psi) \gamma^a \gamma_5 \Psi + \frac{mc}{\hbar} \Psi &= 0 \end{aligned}$$

is equivalent to the Einstein theory of the neutrino Dirac spinor field in the Riemannian spacetime:

$$G_{(\mu\nu)} = \frac{\kappa\hbar c}{2} \left\{ \overset{\{\}}{D}_{(\mu} \bar{\Psi} \gamma_{\nu)} \Psi - \bar{\Psi} \gamma_{(\mu} \overset{\{\}}{D}_{\nu)} \Psi \right\},$$

$$\gamma^a \overset{\{\}}{D}_a \Psi = 0,$$

These results actually reveal the geometrical nature of nonlinearity of the Ivanenko-Heisenberg [58] equations<sup>1</sup>. It must be stressed that our conclusions follow from the analysis of the complete system of equations describing the self-gravitating spinor fields.

However, it is worthwhile to mention that the equality of the coupling constants  $l_0$  and  $\lambda_p$  in the Ivanenko-Heisenberg equation is physically unsubstantiated unless it becomes a natural part of a fundamental theory, or the specific physical conditions arise which lead to such an equality. In particular, one can show that the aforementioned equivalence may take place at the initial stages of the cosmological evolution [60] by means of the spontaneous symmetry breaking mechanism of the vacuum state of a self-interacting scalar field with conformal coupling [71, 72] in an external metric of an open Friedman model [61].

Another possibility to adjust the coupling constants  $l_0$  and  $\lambda_p$ , is to extend the Einstein-Cartan model (as it is actually required in the gauge theory for a non-semisimple group) by including into the gravitational Lagrangian, along with the Hilbert term with Einstein's coupling constant  $\kappa$ , the term quadratic in the torsion tensor with an additional dimensionless coupling constant  $\chi$ . A particular example of such an extended theory gives the Lagrangian  $\frac{1}{2\kappa c} (R(\Gamma) + \chi Q_\mu^{\lambda\nu} Q^\mu{}_{\lambda\nu})$ .

### *Vector fields in ECT*

The gravitational interaction of vector fields is an interesting issue in the framework of the Einstein-Cartan theory. If we start with a theory of a massless vector field  $A_\mu$  in the flat spacetime, and formally apply the minimal coupling principle by replacing the ordinary derivatives by the covariant ones, then in  $U_4$  we obtain the Lagrangian

$$\mathcal{L}_A = -\frac{1}{16\pi} F_{\alpha\beta}(\Gamma) F^{\alpha\beta}(\Gamma),$$

where the generalized vector field strength tensor reads

$$F_{\alpha\beta}(\Gamma) = \nabla_\alpha A_\beta - \nabla_\beta A_\alpha = F_{\alpha\beta} + 2Q^\lambda{}_{\alpha\beta} A_\lambda.$$

---

<sup>1</sup>The dynamics of the two-component neutrino in the  $U_4$  Riemann-Cartan spacetime identically coincides with the dynamics of the neutrino in the Riemann spacetime. Within the so-called strong gravitation of A. Salam [59] one has  $l_0 = \lambda_p$ .

Here  $F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$  is the usual Maxwell tensor. The field equations for the coupled gravitational and vector fields are obtained from the action (1.34) by the variation with respect to the fundamental variables  $g_{\mu\nu}$ ,  $\Gamma^\lambda{}_{\mu\nu}$ ,  $A_\mu$ , and the Lagrange multipliers  $\Lambda^{\alpha\beta\gamma}$ . The resulting system of the Einstein-Cartan equations (1.48) and (1.49) describes the dynamics of the gravitational fields ( $g_{\mu\nu}$ ,  $Q^\lambda{}_{\mu\nu}$ ) created by its sources: the canonical tensors of the energy-momentum and the spin of the vector field

$$t_{(\mu\nu)} = \frac{1}{4\pi} \left\{ F_{\mu\alpha}(\Gamma) F_\nu{}^\alpha(\Gamma) - \frac{1}{4} g_{\mu\nu} F_{\alpha\beta}(\Gamma) F^{\alpha\beta}(\Gamma) \right. \\ \left. + (\nabla_\lambda - 2Q_\lambda) A_{(\mu} F_{\nu)}{}^\lambda(\Gamma) \right\}, \\ S^\lambda{}_{\mu\nu} = -\frac{1}{4\pi c} A_{[\mu} F_{\nu]}{}^\lambda(\Gamma).$$

In addition, the dynamics of the vector field  $A_\mu$  is described by the equation

$$(\nabla_\alpha - 2Q_\alpha) F^{\alpha\beta}(\Gamma) = 0. \quad (4.27)$$

Let us solve the Palatini equation (1.49) with respect to the torsion tensor. This is a nontrivial task since the torsion appears on both sides (explicitly on the left-hand side, and implicitly in  $F_{\mu\nu}(\Gamma)$  on the right-hand side). After some algebra we find

$$Q^\lambda{}_{\mu\nu} = -\frac{2G}{c^4} \left\{ A_{[\mu} F_{\nu]}{}^\lambda + \frac{(\frac{G}{c^4} A^\lambda A_{[\mu} - \frac{1}{2} \delta_{[\mu}^\lambda) F_{\nu]\sigma} A^\sigma}{1 - \frac{G}{c^4} A^2}} \right\}. \quad (4.28)$$

Here we denoted  $A^2 = A_\mu A^\mu$ , and let us remind that  $F_{\mu\nu}$  (no dependence on  $\Gamma$ ) is the usual Maxwell tensor. For the trace we have a simple result

$$Q_\mu = -\frac{G}{2c^4} F_{\mu\nu}(\Gamma) A^\nu = -\frac{G}{2c^4} \frac{F_{\mu\nu} A^\nu}{1 - \frac{G}{c^4} A^2}.$$

Note that  $Q^\mu A_\mu = 0$ .

Substituting (4.28) into (4.27), we obtain the generalized ‘‘Maxwell equation’’, the nonlinearity of which is caused by space-time torsion induced by the vector field spin [62]

$$\frac{1}{\sqrt{g}} \partial_\alpha (\sqrt{g} F^{\alpha\beta}(\Gamma)) = \frac{4\pi}{c} J^\beta, \quad (4.29)$$

where the effective 4-current reads

$$J^\beta = \frac{G}{4\pi c^3} F^{\alpha\beta}(\Gamma) F_{\alpha\gamma}(\Gamma) A^\gamma.$$

This current satisfies the continuity relation  $\partial_\beta (\sqrt{g} J^\beta) = 0$  by virtue of the field equation (4.29). Making use of the torsion (4.28), we have the explicit form of the generalized vector field strength

$$F_{\mu\nu}(\Gamma) = F_{\mu\nu} - \frac{\frac{2G}{c^4} A_{[\mu} F_{\nu]}{}^\lambda A^\lambda}{1 - \frac{G}{c^4} A^2}. \quad (4.30)$$

As we explained in Sec. 1.3., the system of the vector field equation (4.27), (4.29) plus the gravitational equation (1.46) can be rewritten as the system of the Einstein equation of GR and of the non-linear vector field equation described by the total Lagrangian  $\frac{1}{2\kappa}R + \mathcal{L}_A^{\text{eff}}$  with the effective Lagrangian

$$\mathcal{L}_A^{\text{eff}} = -\frac{1}{16\pi} \left\{ F_{\mu\nu} F^{\mu\nu} + \frac{2\frac{G}{c^4} F_{\mu\alpha} F^{\mu\beta} A^\alpha A_\beta}{1 - \frac{G}{c^4} A^2} - \frac{2}{\sqrt{g}} \partial_\mu \left( \frac{\sqrt{g} F^{\mu\nu} A_\nu}{1 - \frac{G}{c^4} A^2} \right) \right\}.$$

The torsion was eliminated with the help of (4.28), so that  $R$  is the scalar of the Riemannian curvature, and  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the usual Maxwell tensor. The last term is a total divergence and it does not contribute to the field equations. One can directly verify that the nonlinear vector field equation (4.29) is obtained from the effective Lagrangian above by means of the variation with respect to the vector field  $A_\mu$ , so that  $J^\beta = -c\partial\mathcal{L}_A^{\text{eff}}/\partial A_\beta$ .

When introducing the interaction of a massless vector field with the spacetime torsion, the gauge invariance of the massless vector field theory (4.29) is violated<sup>2</sup>, which essentially changes the dynamical content of the theory. Indeed, the usual Maxwell equations describe the massless spin-1 particle (photon) with the two degrees of freedom. But now the vector field  $A_\mu$  has 3 degrees of freedom and thus cannot be interpreted as the electromagnetic field. Nevertheless, the resulting nonlinear theory still does have the current conservation law, namely  $\partial_\beta(\sqrt{g}J^\beta) = 0$ , i.e., the theory under consideration is partially gauge-invariant in the sense of Glashow and Gell-Mann [64]. Partial gauge-invariant nature of the vector field in the presence of the gravitation is well consistent with the interaction hierarchy, where the weaker interaction violates the symmetries typical for the stronger interaction. In the framework of such a hierarchy, the requirement of the local gauge invariance, according to which each conservation law corresponds to some gauge field, turns out to be valid only approximately, when the weaker interactions are neglected.

Let us study the ‘‘electrostatic’’ solutions of the equation (4.29) in the Minkowski spacetime with  $g_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ . Denoting the Cartesian coordinates  $x^\mu = \{t, x^i\}$ , with  $i = 1, 2, 3$ , we use the standard ansatz for the 4-potential:  $A_\mu = \{\varphi(x^i), 0, 0, 0\}$ . The generalized ‘‘electromagnetic’’ field strength is defined in terms of the tensor  $F_{\alpha\beta}(\Gamma) = (\mathbf{E}, \mathbf{B})$ , which is a covariant (in terms of the spacetime  $U_4$ ) generalization of Maxwell’s tensor  $F_{\alpha\beta}$ . Then (4.29) reduces to the Gauss type equation [62]

$$\nabla \cdot \mathbf{E} = 4\pi\rho, \quad (4.31)$$

where from (4.30) we find

$$\rho = \frac{G}{4\pi c^4} \mathbf{E}^2 \varphi, \quad \mathbf{E} = \frac{-\nabla\varphi}{1 + \frac{G}{c^4} \varphi^2}.$$

---

<sup>2</sup>Something similar happened for the covariant generalization of spinor, neutrino (massless) equations, when their conformal invariance was violated.

The equation (4.31) for the function  $\varphi$  explicitly reads

$$\left(1 + \frac{G}{c^4}\varphi^2\right)\Delta\varphi - \frac{G}{c^4}\varphi(\nabla\varphi)^2 = 0.$$

Although the resulting nonlinear equation looks quite nontrivial, remarkably there exists a substitution  $\varphi = \frac{c^2}{\sqrt{G}}\sinh\chi$ , that yields a linear Laplace equation of Maxwell' electrostatics  $\Delta\chi = 0$ .

It is straightforward to obtain a spherically symmetric solution of (4.31) for  $\varphi(x^i) = \varphi(r)$ , with  $r = \sqrt{x_i x^i}$ :

$$\varphi = \frac{c^2}{\sqrt{G}}\sinh\left(\frac{q\sqrt{G}}{c^2 r}\right).$$

Here  $q$  is an integration constant with a dimension of a charge. This solution obviously has a Coulomb asymptotic behaviour at infinity, for  $r \rightarrow \infty$ . However, the usual Coulomb singularity at  $r = 0$  is absent<sup>3</sup>. As a result, the corresponding “electric” field and the “charge” density are also regularized:

$$\mathbf{E} = \frac{q}{\cosh\left(\frac{q\sqrt{G}}{c^2 r}\right)} \frac{\mathbf{r}}{r^3}, \quad \rho = \frac{\sqrt{G} \sinh\left(\frac{q\sqrt{G}}{c^2 r}\right)}{4\pi c^2 \cosh^2\left(\frac{q\sqrt{G}}{c^2 r}\right)} \frac{q^2}{r^4}.$$

The qualitative behaviour of the solution, see Fig. 4.1, is as follows<sup>4</sup>: Both functions display the Coulomb asymptotics at infinity, whereas they both vanish at the origin,  $\mathbf{E}(0) = 0, \rho(0) = 0$ . The integral over the 3-space of the charge density is finite and easily evaluated:

$$\int_V d^3x \rho = q,$$

so that we can indeed identify the integration constant  $q$  with the total “charge”. The nonsingular generalized “electric” field reaches its largest value at  $r_{E_{max}} = 0.14 \times 10^{-24} q$ , whereas the charge density is maximal at  $r_{\rho_{max}} = 0.07 \cdot 10^{-24} q$ . To make an estimate, for an elementary charge of an electron  $q = 4.8 \cdot 10^{-10}$  (in Gaussian absolute electrostatic units), we find  $r_{E_{max}} = 0.7 \cdot 10^{-34}$  cm and  $r_{\rho_{max}} = 0.35 \cdot 10^{-34}$  cm, respectively.

One can evaluate the energy of the regular “electromagnetic” field configuration corresponding to this solution. It is straightforward to write down the energy density for the spherically symmetric solution making use of the canonical energy-momentum tensor:

$$\mathcal{H} = t_{00} = \frac{1}{8\pi} \mathbf{E}^2 - \varphi \rho + \frac{1}{4\pi} \nabla(\varphi \mathbf{E}).$$

<sup>3</sup>Note, that such form of the potential is obtained by Urbakh within proposed nonlinear electrodynamics [55].

<sup>4</sup>Taking into account the static spherically symmetric gravitational metric field  $g_{\mu\nu}$  does not essentially change the behaviour of  $\mathbf{E}(r)$  and  $\rho(r)$  [62, 67].




Figure 4.1: Behaviour of the static electric field strength (a) and distribution of the charge density (b) (dotted line corresponds to Coulomb's law).

Using the Gauss theorem we see that the last term does not contribute to the total energy since both the potential  $\varphi$  and the “electrostatic” field  $\mathbf{E}$  vanish at the spatial infinity. The first term on the right-hand side is analogous to the energy density of linear (Maxwell's) electromagnetic field, whereas the second may be interpreted as the energy density of the self-interaction due to the nonlinear character of the resulting effective theory. The latter term is singular at the origin and it makes a divergent contribution to the integrated total mass  $\int d^3x \mathcal{H}$ . Subtracting this divergent self-interaction term, we find the total mass of “electrostatic” field configuration:

$$m_{\text{em}} = \frac{W}{c^2} = \frac{1}{8\pi c^2} \int d^3x \mathbf{E}^2 = \frac{q}{\sqrt{G}}.$$

For the electron charge,  $m_{\text{em}} = \sqrt{\alpha \frac{\hbar c}{G}} = 1.86 \times 10^{-9} \text{ kg}$ , which is close to the mass of the classical Markov maximon [66] (with the fine structure constant  $\alpha \approx 1/137$ ).

We conclude our discussion of the vector fields by a brief outline of the massive case. The mass term in the Lagrangian does not change the Palatini equation, and the effective nonlinear field equation of the massive vector field is obtained similarly to (4.29) by eliminating the torsion using (1.49) and (4.28):

$$\frac{1}{\sqrt{g}} \partial_\alpha (\sqrt{g} F^{\alpha\beta}(\Gamma)) - m_A^2 A^\beta = \frac{4\pi}{c} J^\beta. \quad (4.32)$$

Using the static ansatz  $A_\mu = \{\varphi(x^i), 0, 0, 0\}$ , and subsequently making a substitution  $\varphi = \frac{c^2}{\sqrt{G}} \sinh(\eta/2)$ , we recast (4.32) into

$$\Delta\eta = m_A^2 \sinh \eta. \quad (4.32a)$$

Then for the spherically symmetric solution we find  $\eta \sim \frac{\exp(-m_A r)}{r}$ , i.e., the solution asymptotically (with  $r \rightarrow \infty$ ) approaches the Yukawa potential.

Another interesting class of exact solutions can be obtained for the ansatz  $A_\mu = \{0, 0, A_2(x, t), 0\}$  [63]. Plugging this into the nonlinear equation (4.32), and making a substitution  $A_2 = \frac{c^2}{\sqrt{G}} \sin \phi$ , we derive the well-known sin-Gordon equation

$$\square \phi - m_A^2 \sin \phi = 0,$$

where the 2-dimensional d'Alembert operator  $\square = -\frac{1}{c^2} \frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2}$ . This is a completely integrable system which admits soliton solutions.

Therefore, we established the geometrical nature of nonlinearity of sine-Gordon in the same way that the geometrical nature of nonlinearity of spinor fields of Ivanenko-Heisenberg was established.

## 4.2. Conformal invariance and spacetime torsion

In Sec. 1.3., when varying the Lagrangian  $\mathcal{L}_{ECT}$  independently with respect to the metric  $g_{\mu\nu}$  and the linear connection  $\Gamma^\lambda_{\mu\nu}$  without imposing restrictions on these dynamical variables, we obtained the semi-metric theory with the connection (1.32) which is defined by the condition (1.31). This manifests a close relation between the torsion vector  $Q_\mu$  and the nonmetricity vector  $K_\mu$  that appears in the Weyl theory [5]:

$$\nabla_\alpha g_{\mu\nu} = K_\alpha g_{\mu\nu}.$$

Weyl constructed his theory of gravity on the basis of conformal transformations of the interval  $ds^2$ . Weyl's theory is invariant under the transformation of the metric of the following form:

$$g_{\mu\nu} \longrightarrow g'_{\mu\nu} = e^{2\sigma} g_{\mu\nu}. \quad (4.33)$$

These are the so-called Weyl transformations, parametrized by an arbitrary scalar function  $\sigma = \sigma(x^\mu)$ . Despite the mathematical elegance of this theory, a quite serious objection was found that strongly depreciated Weyl's gravity model: the length standards and the clocks do not have an invariant meaning in this theory, and in particular, they "depend on the history". Having failed to overcome this problem, the Weyl theory remained a beautiful geometrical model without direct applications to physics.

However, taking into account the significance of the conformal symmetry in physics, it is interesting to look for a possibility to include Weyl's transformations (4.33) into a gravity theory.<sup>5</sup>

---

<sup>5</sup>This is once again demonstrated in another Dirac's attempt to introduce the dilations by assuming the existence of different types of standards [68].

Here we outline a possible natural solution of the problem of the length standards by constructing the conformally invariant gravity theory of the Hilbert-Einstein type in the Riemann-Cartan spacetime  $U_4$ , [69].

The tetrad formalism proves to be most useful for our purposes, and we take the tetrad and the local Lorentz connection fields,  $h_\mu^a$  and  $\Gamma^a_{b\mu}$ , as the basic independent dynamic variables. The crucial point is as follows: we postulate that the conformal Weyl transformations (4.33) of the metric are induced by

$$\left. \begin{aligned} h_\mu^a &\longrightarrow h'^a_\mu = e^\sigma h_\mu^a, \\ \Gamma^a_{b\mu} &\longrightarrow \Gamma'^a_{b\mu} = \Gamma^a_{b\mu}. \end{aligned} \right\} \quad (4.34)$$

Under these transformations, the tetrad is scaled whereas the Lorentz connection remains invariant, hence the Riemann-Cartan curvature is not changed too:  $R'^a_{b\mu\nu} = R^a_{b\mu\nu}$ . However, the world connection is transformed in accordance with (1.20) as

$$\Gamma^\alpha_{\beta\mu} \longrightarrow \Gamma'^\alpha_{\beta\mu} = \Gamma^\alpha_{\beta\mu} + \delta^\alpha_\beta \partial_\mu \sigma. \quad (4.35)$$

These are the famous Einstein's  $\lambda$ -transformations [70] which leave the curvature (1.1) tensor  $R^\alpha_{\beta\mu\nu}(\Gamma)$  (constructed from the world connection  $\Gamma^\alpha_{\beta\mu}$ ) invariant. It is important that the Riemann-Cartan geometrical structure is preserved by these transformations. This is evident from the fact that the metricity condition (1.8) is not violated:

$$\nabla_\alpha g_{\mu\nu} = 0 \longrightarrow \nabla'_\alpha g'_{\mu\nu} = 0.$$

In accordance with (4.35), the torsion tensor (1.3) is transformed as

$$Q^\alpha_{\mu\nu} \longrightarrow Q'^\alpha_{\mu\nu} = Q^\alpha_{\mu\nu} + \delta^\alpha_{[\mu} \partial_{\nu]} \sigma. \quad (4.36)$$

A remarkable fact is that (4.36) affects the trace of torsion only

$$Q_\mu \longrightarrow Q'_\mu = Q_\mu - \frac{3}{2} \partial_\mu \sigma.$$

The rest of the irreducible parts of the torsion (1.13) are conformally invariant.

Now we will demonstrate that the group of local conformal transformations introduced above (4.34) in the spacetime  $U_4$  provides a natural basis for the description of the scale symmetry of both matter and gravitation.

At first, we consider the massless fermion field described by the Dirac action  $\frac{1}{c} \int d^4x \sqrt{g} \mathcal{L}_D$  with the Lagrangian (4.8) in which we put the rest mass equal zero  $m = 0$ . The fermion action is explicitly invariant with respect to scale transformations (4.34) provided the Dirac field is scaled in accordance with its canonical dimension,

$$\Psi \longrightarrow \Psi' = e^{-\frac{3}{2}\sigma} \Psi.$$

Let us note that the extra terms proportional to  $\partial_\mu \sigma$  occurring in the transformation (4.34) are mutually cancelled, so there is no need to introduce additional

Weyl fields. It is worthwhile to note that such an intrinsic conformal invariance is actually provided by the torsion. Indeed, the variation of (4.8) with respect to  $\bar{\Psi}$  leads to the Dirac equation (4.13) in  $U_4$  (with  $m = 0$ ):  $\gamma^\mu (D_\mu - Q_\mu) \Psi = 0$ . This equation is explicitly conformally invariant due to the fact that the torsion trace  $Q_\mu$  acts as the Weyl field.

Now we will show the possibility to construct a conformally invariant theory of gravity in  $U_4$ . Let us study the Hilbert-Einstein model with the action<sup>6</sup>

$$S_{\text{conf}} = \int d^4x \sqrt{g} \varphi^2 R(\Gamma). \quad (4.37)$$

To provide the conformal invariance, we introduced an additional field  $\varphi$  with the scaling behaviour under (4.34)

$$\varphi \longrightarrow \varphi' = e^{-\sigma} \varphi,$$

in accordance with its canonical dimension. For the curvature scalar we obviously have  $R'(\Gamma) = e^{-2\sigma} R(\Gamma)$ , whereas  $\sqrt{g'} = e^{4\sigma} \sqrt{g}$ . The role of the scalar field  $\varphi$  is (as in the similar cases in the Riemann geometry) to introduce a natural scale in the theory. Comparing (4.37) with the action (1.34) of the Einstein-Cartan model, we notice that the scalar field square  $\varphi^2$  replaces  $\frac{1}{2\kappa c}$  and thus plays the role of a Brans-Dicke-type variable coupling function.

The field equations of the conformally invariant theory of gravity in  $U_4$  can be obtained by variation of (4.37) either with respect to the metric and the torsion  $g_{\mu\nu}, Q^\lambda{}_{\mu\nu}$ , or in the tetrad formalism with respect to the tetrad and the local Lorentz connection  $h^a{}_\mu, \Gamma^a{}_{b\mu}$ . Both methods yield the same result. In the absence of the matter, we obtain the field equations

$$G_{\mu\nu}(\Gamma) = 0, \quad (4.38)$$

$$Q^\lambda{}_{\mu\nu} = \frac{1}{\varphi} \delta^\lambda_{[\nu} \partial_{\mu]} \varphi. \quad (4.39)$$

In addition, we have to vary the action with respect to the scalar field  $\varphi$ . However, the corresponding field equation,  $R(\Gamma) = 0$ , is redundant in view of (4.38). The equation (4.39) shows that the torsion is expressed in terms of its trace  $Q_\mu$ , for which the scalar field  $\varphi$  serves as a ‘‘potential’’. Substituting (4.39) into (4.38), we obtain

$$G_{\mu\nu} = -\frac{6}{\varphi^2} T_{\mu\nu}, \quad (4.40)$$

where

$$T_{\mu\nu} = \partial_\mu \varphi \partial_\nu \varphi - \frac{1}{2} g_{\mu\nu} \partial_\alpha \varphi \partial^\alpha \varphi - \frac{1}{6} \left( \overset{\{\}}{\nabla}_\mu \overset{\{\}}{\nabla}_\nu - g_{\mu\nu} \overset{\{\}}{\square} \right) \varphi^2$$

is the so-called ‘‘improved’’ energy-momentum tensor of Callan-Coleman-Jackiw [71, 72] for the scalar field  $\varphi$ . Here  $\overset{\{\}}{\square} := \overset{\{\}}{\nabla}_\mu \overset{\{\}}{\nabla}^\mu$  is the covariant d’Alembertian

---

<sup>6</sup>The Lagrangian of such a theory may also include all possible terms quadratic in the curvature  $R^\alpha{}_{\beta\mu\nu}(\Gamma) R^\alpha{}_{\beta\mu\nu}(\Gamma), R_{\mu\nu}(\Gamma) R^{\mu\nu}(\Gamma), R^2(\Gamma)$ , etc., which are all conformally invariant.

with respect to the Christoffel connection of the Riemannian space. The contraction of (4.40) yields the explicitly conformally invariant equation

$$\square \varphi - \frac{R}{6} \varphi = 0. \quad (4.41)$$

Thus, we once again observe how the torsion provides the intrinsic conformal invariance of the gravitational theory. The “wrong” sign of the energy-momentum tensor  $T_{\mu\nu}$  of the scalar field in (4.40) manifests a non-physical nature of  $\varphi$  which is actually the Goldstone type field [73, 74]. Using the conformal freedom, we can choose the gauge  $\varphi = \text{const}$  (which means the choice of a certain length scale), and thus we arrive at the standard ECT.

Summarizing, we have demonstrated a natural way to introduce the Weyl transformations (4.33) and to construct the conformally invariant theory of gravity of the Hilbert-Einstein type in the Riemann-Cartan spacetime with torsion. Distinctive features of such an approach are the postulate of the tetrad scaling (4.34) as a source of conformal transformations (4.33) and the special role of the torsion trace  $Q_\mu$  as an effective Weyl field.

One of the most important advantages is the absence of the problem of the length scales, since  $D_\mu \eta_{ab} = 0$  and, correspondingly,  $\nabla_\alpha g_{\mu\nu} = 0$  (which are both explicitly conformally invariant relations). Consequently, length standards are conserved and there is no need in Dirac’s propositions of the two types of clocks [68], etc.

The analysis of the possible relation of the torsion with conformal transformations clarifies the source of the vacuum production process of scalar particles described by the Klein-Gordon equation with the conformal coupling in the homogeneous and isotropic spacetimes (see Sec. 4.4.). Indeed, we find  $Q_\mu = 0$  due to the Palatini equation (4.16) which results in the violation of the conformal invariance of the Dirac equation.

### 4.3. Pre-Friedman stage of Universe’s evolution and spacetime torsion

In this section, we turn to the study of the cosmological effects of ECT.

The theory of an expanding Universe – the Friedman cosmology based on Einstein’s equations (with the initial data) – is the most spectacular achievement of GR. Numerous important physical effects predicted on its basis were verified experimentally [27], among them the relict microwave radiation, the cosmological red shift, etc. However, the problem of an initial state of the Universe remains unsolved within this framework. The presence of singularities in the general solution of Einstein’s equations [86] is one of the central problems of the theory of gravitational interactions. One can expect that a further development of the fundamental principles of Einstein’s GR, that would allow to take into account the effects of the gravitational and matter fields which

are insignificant under the normal conditions (quantum and/or microstructural properties, e.g.), may provide possible ways to solve these problems [87].

In connection with this, one of the most important achievements of the Einstein-Cartan theory is the prediction of the avoidance (for an appropriate choice of the torsion tensor and of its spinning sources) of singularities in the cosmological models of the Universe. Such an opportunity is opened up when the torsion violates the Penrose-Hawking energy condition [86], which in the ECT takes the following form [7, 88]:

$$T_{\mu\nu}^{\text{eff}} t^\mu t^\nu \geq \frac{1}{2} T_{\mu\nu}^{\text{eff}} g^{\mu\nu}, \quad (4.42)$$

where  $t^\mu$  is an arbitrary time-like vector.

Such a violation is not unexpected. The Einstein-Cartan theory, as we will demonstrate in Chapters 5, 6, represents a special model of the gauge theory of gravity for the Poincaré group  $\mathcal{P}_{10}$  (a semidirect product of the Lorentz group  $SO(1, 3)$  times the translation group  $T_4$ ). The non-compactness of the Lorentz group results in the indefiniteness of the energy sign, which leads to a possibility of violation of the condition (4.42).

It is important to note that in the regions occupied by the spinning matter, even in the case of chaotically oriented spins such that the mean value  $\langle S^\lambda{}_{\mu\nu} \rangle = 0$  vanishes, the macroscopic spin effects are different from zero. This is due to the fact that in the equation (1.52) that has a microscopic meaning before the averaging procedure, there are terms quadratic in spin. Let us model the source of the gravitational field by the ideal spinning Weyssenhoff fluid (see Sec. 4.1.), consisting of fermions with the particle density  $\rho$ , then  $S^2 = \frac{1}{2} S_{\mu\nu} S^{\mu\nu} = \frac{\hbar^2}{8} \rho^2$ . Averaging yields  $\langle S^2 \rangle = \langle \frac{1}{2} S_{\mu\nu} S^{\mu\nu} \rangle = \hbar^2 \langle \rho^2 \rangle / 8$ , where the mean particle density  $\langle \rho \rangle$  is related to the energy density  $\varepsilon$  via  $\varepsilon = A_k \langle \rho \rangle^{1+k}$  for the equation of state  $p = k\varepsilon$  with  $0 \leq k \leq 1$ , and  $A_k$  is some constant depending on the equation of state [120]. Then

$$\langle S^2 \rangle = \frac{\hbar^2}{8A_k^{\frac{2}{1+k}}} \varepsilon^{\frac{2}{1+k}}. \quad (4.43)$$

If  $\langle S \rangle = 0$ , one can understand  $\langle S^2 \rangle$  as a squared dispersion of spin density distribution about the zero mean value,

$$\langle (\Delta S)^2 \rangle = \langle (\langle S \rangle - S)^2 \rangle = \langle \langle S \rangle \langle S \rangle - 2S \langle S \rangle + S^2 \rangle = \langle S^2 \rangle.$$

The existence of a nonzero  $\langle S^2 \rangle$  here is similar to the existence of a nonzero average kinetic molecular energy  $\langle p^2/2m \rangle \neq 0$ , when the average momentum vanishes,  $\langle p \rangle = 0$ , for the classical gas in equilibrium.

Let us consider the homogeneous isotropic cosmological model with the metric

$$ds^2 = -c^2 dt^2 + R^2(t) \left[ \frac{dr^2}{1 - \mathcal{K}r^2} + r^2 d\Omega^2 \right], \quad \mathcal{K} = \pm 1, 0, \quad (4.44)$$

with the Weyssenhoff spinning fluid as a matter source. The parameter  $\mathcal{K} = +1$  ( $\mathcal{K} = -1$ ) corresponds to the closed (open) Universe, and  $\mathcal{K} = 0$  to the quasi-Euclidean world [120]. Einstein's equations (1.52) with the effective energy-momentum tensor (4.7) reduce to

$$\frac{\dot{R}^2(t)}{R^2(t)} + \frac{\mathcal{K}c^2}{R^2(t)} = \frac{8\pi G}{3c^2} \left( \varepsilon - \frac{8\pi G}{c^2} S^2 \right), \quad (4.45a)$$

$$-\frac{2\ddot{R}(t)}{R(t)} - \frac{\dot{R}^2(t)}{R^2(t)} - \frac{\mathcal{K}c^2}{R^2(t)} = \frac{8\pi G}{c^2} \left( p - \frac{8\pi G}{c^2} S^2 \right). \quad (4.45b)$$

The covariant conservation law of the effective energy-momentum tensor  $T_{\text{eff}}^{\mu\nu}{}_{;\mu} = 0$  yields

$$\frac{\dot{\varepsilon}}{\varepsilon + p} = 3 \frac{\dot{R}}{R}. \quad (4.46a)$$

Therefore, for the case of the dust ( $p = 0$ ), we have<sup>7</sup>

$$\varepsilon = \frac{\varepsilon_0}{R^3}, \quad S^2 = \frac{S_0^2}{R^6}, \quad (4.46b)$$

where  $\varepsilon_0$  and  $S_0$  are the integration constants which describe the energy density and the particle density of the matter in the Universe at the moment of time when  $R = 1$ . When obtaining (4.46b), we took (4.43) into account.

Substituting (4.46a) into the Friedman equation (4.45a), we find

$$\dot{R}^2 = -\mathcal{K}c^2 + \frac{8\pi G\varepsilon_0}{3c^2 R} - \frac{64\pi^2 G^2 S_0^2}{3c^4 R^4}. \quad (4.47)$$

As compared to GR, we have an additional term proportional to  $R^{-4}$  which arises due to the ( $\Gamma$ - $S$ )-interaction. It can be interpreted as a centrifugal potential energy, and it is easy to see that this term leads to the absence of a cosmological singularity. In particular, for the quasi-Euclidean Universe with  $\mathcal{K} = 0$ , the explicit solution of (4.47) reads

$$R(t) = \sqrt[3]{R_{\text{min}}^3 + 6\pi G\varepsilon_0 t^2/c^2}, \quad R_{\text{min}}^3 = \frac{8\pi G S_0^2}{c^2 \varepsilon_0}. \quad (4.48)$$

To make an estimate, if we assume that the Universe is filled with  $N \approx 10^{80}$  nucleons (hence the total mass of matter  $\approx 10^{53}$  kg), we find for the minimal value of the cosmological scale factor  $R_{\text{min}} \approx 10^{-28}$ . The corresponding minimal radius of the Universe  $\approx 10^{-2}$  m, and the maximal density  $\approx 10^{57}$  kg/m<sup>3</sup> [97].

Another estimate can be obtained if we replace the nucleon as an elementary unit of matter with a hypothetical ‘‘fundamental atom’’ – the Planckon [121]. The latter is a black hole with the mass of  $m_{\text{pl}} \approx 10^{-8}$  kg and the spin  $\frac{1}{2}\hbar$ .

---

<sup>7</sup>On the right-hand side of the equations, the averaged quantities are understood.

Since the Compton wave length of particles  $\lambda = \hbar/mc$  cannot be greater than the gravitational radius of a configuration, the Planckeon is a smallest possible black hole [122]. Then, assuming that at an initial moment of time the Universe (as the interior of a collapsing star) consisted of completely spin-polarized Fermi fluid formed by  $10^{61}$  Planckeoens (to keep the total mass  $\approx 10^{53}$  kg), we find for the minimal radius of the Universe  $\approx 10^{-15}$  m.

Since the spin is a vector and thus it obviously defines a preferred direction in space, thereby breaking the spatial isotropy, it is perhaps more realistic to consider anisotropic spatially homogeneous models. Let us study the Bianchi I model with the metric

$$ds^2 = -c^2 dt^2 + a^2(t)(dx^2 + dy^2) + b^2(t)dz^2,$$

where the two scale factors  $a = a(t)$  and  $b = b(t)$  describe a possible difference of the cosmological dynamics along a preferred axis ( $z$ -coordinate) and in the orthogonal plane.

The effective Einstein equations for this metric read

$$\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{b}}{ab} = \kappa\varepsilon^{\text{eff}}, \quad -\frac{\ddot{a}}{a} - \frac{\ddot{b}}{b} - \frac{\dot{a}\dot{b}}{ab} = \kappa p^{\text{eff}}, \quad \frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} - \frac{\ddot{b}}{b} - \frac{\dot{a}\dot{b}}{ab} = 0.$$

Introducing the new variables

$$R^3 := a^2 b, \quad \sigma := \frac{\dot{a}}{a} - \frac{\dot{b}}{b},$$

we recast this system into

$$\begin{aligned} 3\frac{\dot{R}^2(t)}{R^2(t)} &= \frac{8\pi G}{c^2} \left( \varepsilon - \frac{8\pi G}{c^2} S^2 \right) + \frac{1}{3}\sigma^2, \\ -\frac{2\ddot{R}(t)}{R(t)} - \frac{\dot{R}^2(t)}{R^2(t)} - \frac{\mathcal{K}c^2}{R^2(t)} &= \frac{8\pi G}{c^2} \left( p - \frac{8\pi G}{c^2} S^2 \right) + \frac{1}{3}\sigma^2, \\ \dot{\sigma} + 3\frac{\dot{R}(t)}{R(t)}\sigma &= 0. \end{aligned}$$

The last equation is easily integrated and yields the anisotropy expansion function (the difference of the Hubble functions):

$$\sigma = \frac{\sigma_0}{R^3}.$$

The equation of motion of spin (4.5) is also straightforwardly integrated to give

$$S = \frac{S_0}{a^2 b} = \frac{S_0}{R^3}.$$

Substituting all this into the Friedman equation, for the Weyssenhoff dust ( $p = 0$ ) as a source of gravitational field, we find a modified version of (4.47)

$$\dot{R}^2 = \frac{8\pi G\varepsilon_0}{3c^2 R} - \frac{\mu_0}{R^4},$$

where we introduced a combination of the integration constants

$$\mu_0 = \frac{64\pi^2 G^2 S_0^2}{3c^4} - \frac{\sigma_0^2}{9}.$$

The qualitative behaviour of a cosmological solution depends crucially on the balance between the spin and anisotropy: in simple terms, on the value of the constant  $\mu_0$ . For  $\mu_0 = 0$  one finds the usual singular Friedman solution for the dust. For  $\mu_0 < 0$ , we also have a singular solution. However when  $\mu_0 > 0$ , the solution does not have cosmological singularities [123]:

$$a = \sqrt[3]{\frac{3c^2\mu_0}{8\pi G\varepsilon_0} + \frac{6\pi G\varepsilon_0}{c^2} t^2} \times \exp\left(\frac{2\sigma_0}{9\sqrt{\mu_0}} \arctan \frac{4\pi G\varepsilon_0 t}{c^2\sqrt{\mu_0}}\right),$$

$$b = \sqrt[3]{\frac{3c^2\mu_0}{8\pi G\varepsilon_0} + \frac{6\pi G\varepsilon_0}{c^2} t^2} \times \exp\left(-\frac{4\sigma_0}{9\sqrt{\mu_0}} \arctan \frac{4\pi G\varepsilon_0 t}{c^2\sqrt{\mu_0}}\right).$$

Besides the elimination of singularities in the cosmological solutions, the spin terms produce another interesting effect by stabilizing the matter distribution, so that for the closed Universes (only in this case) there exist static solutions of the Friedman equations (4.45). For the dust, such a Universe has the size  $\approx \lambda_f$  of a Compton wave length of a fermion with the mass  $m_f$ , and the matter density is about  $\rho_{\text{cr}} = m_f^2 c^4 / G \hbar^2$  ( $\approx 10^{57}$  kg/m<sup>3</sup> for a nucleon), [125]. Such a static solution

$$ds^2 = -c^2 dt^2 + R_0^2 \left[ \frac{dr^2}{1-r^2} + r^2 d\Omega^2 \right] \quad (4.49)$$

coincides with the metric of Einstein's Universe [120]. However, in contrast to GR, here the matter density does not vanish, and the cosmological  $\Lambda$ -term is absent.

The dynamics of the local small perturbations is an important issue in cosmology. Within the framework of GR, this problem was solved for the first time by E.M. Lifshitz [124]. One can show that on the background of the static solution (4.49), the density perturbation  $\delta\varepsilon$  develops in the two exponential modes

$$\delta\varepsilon = \frac{c^2 \omega^2}{120G} \left[ C_1 \exp\left(\frac{\sqrt{5}}{3}\omega t\right) + C_2 \exp\left(-\frac{\sqrt{5}}{3}\omega t\right) \right],$$

where  $\omega = 2\pi c/\lambda$ , and  $\lambda$  is a perturbation wave length;  $C_1$  and  $C_2$  are the integration constants. The development of the exponentially growing and decaying modes is typical for any exact static solutions.

Another important type of perturbations is the occurrence of a global homogeneous isotropic motion of matter, that does not violate the spatial symmetry of the model.

The static solution (4.49) obtained in the framework of the Einstein-Cartan theory differs from the Einstein solution of GR with the  $\Lambda$ -term in that it turns out to be stable with respect to this type of perturbations for the equation




Figure 4.2: Cosmological scenario.

of state  $p = k\varepsilon$ , with  $0 \leq k < 1$ . Indeed, small perturbations  $\delta R$  satisfy the equation

$$\delta\ddot{R} + \omega^2(k)\delta R = 0, \quad \omega^2(k) = \frac{4\pi G\varepsilon_0}{c^2}(-3k^2 + 2k + 1), \quad (4.50)$$

where  $\varepsilon_0$  is the constant energy density of the solution, and  $\omega^2(k) > 0$  for  $0 \leq k < 1$ .

These properties allow one to view such a static solution as the initial and the final stages of the observable non-stationary picture of the Universe [125]. In such a model, the evolution of the Universe consists of the following stages (Fig. 4.2): the time interval  $t_0t_1$  describes the initial static state, when small perturbations are quickly damped; at the moment  $t_1'$  a large fluctuation occurs which is sufficient to form for the Universe observed today; the interval  $t_1't_2$  covers the initial stage of expansion with the dominating torsion effects; the next interval  $t_2t_3$  is a stage of the “standard” cosmology when the torsion effects are negligible and the solution is close to the Friedman one; finally, during the interval  $t_3t$  the Universe returns back to the static state. If the energy of the initial perturbation does not manage to be spent on dissipative phenomena during the expansion and contraction phases, the Universe contracts to a smaller size than the original one  $R_0$  and returns to the initial scale after several oscillations (the branch *I*), otherwise the damping is smooth (the branch *II*). No matter how large the initial perturbation is, the expansion necessarily is followed by a contraction (the closed Universe), and the contraction does not reach a singularity due to the short-range spin term ( $\sim S^2$ ) in the Friedman equations.

The issue of a transition from the static stage ( $t_0t_1$ ) to the dynamical one ( $t_1't_2$ ) is not completely clear in this model. In order to answer this question, we consider in detail how the small homogeneous isotropic perturbations behave. Up to now, we neglected the dissipative phenomena (i.e., the increase of the entropy) and, as a result, we obtained the small harmonic oscillations (4.50) of the scale factor around its equilibrium value  $R_0$ . However, it is easy to see that the second law of thermodynamics forbids the exact repetition of the previous

cycles. Moreover, the increase of the entropy  $s$ ,  $ds/dt > 0$  (both at the expansion stage, and at the contraction stage) turns out to be a natural mechanism that drives the initially static Universe into an observable strongly non-stationary mode. Indeed, the condition that the scale factor reaches an extremal value  $R_{\text{extr}}$  is given by the Friedman equation (4.45a):

$$R_{\text{extr}}^2 \left\{ \varepsilon(R_{\text{extr}}, s) - \frac{8\pi G}{c^2} S^2(R_{\text{extr}}) \right\} = \frac{3c^4 \mathcal{K}}{8\pi G}. \quad (4.51)$$

Evaluating the logarithmic derivative of (4.51), after some calculations, we find

$$\frac{dR_{\text{min}}^{\text{max}}}{ds} = \pm \frac{4\pi G}{3c^2} \frac{R_{\text{min}}^{\text{max}}}{\left| \ddot{R}(R_{\text{min}}^{\text{max}}) \right|} \frac{\partial \varepsilon}{\partial s}, \quad \frac{\partial \varepsilon}{\partial s} > 0. \quad (4.52)$$

The equation (4.52) demonstrates that the maximum of the scale factor  $R_{\text{max}}$  in the repeating cycles increases due to the irreversible phenomena such as friction (for example, the viscosity of matter). This ultimately results in the natural transition from the static pre-Friedman stage ( $t_0 t_1$ ) to the dynamical Friedman stage ( $t'_1 t_2$ ).

Summarizing the analysis of the influence of spin on the cosmological evolution of the Universe in the framework of the Einstein-Cartan theory, we can make the following conclusion.

Already the simplest model of a spinning matter in the form of the Weyssenhoff fluid demonstrates that the torsion brings in a new essential bit to the understanding of the spacetime structure of the initial state of the Universe. However, as was noted above, the Weyssenhoff fluid represents a quasi-classical limit of a real quantum matter with spin. Thus, the consistent analysis of cosmology in the Einstein-Cartan model should be performed also for the Dirac spinor matter. Although the *classical* spinor fields do not eliminate<sup>8</sup> singularities [126], the quantum treatment of spinors [127] essentially confirms the correctness of the results obtained for the Weyssenhoff fluid.

#### 4.4. Production of scalar particles by cosmological torsion field

One of the most important issues of the modern relativistic astrophysics is the problem of the gravitational collapse, i.e., the study of the evolution of massive gravitational systems (both, isolated objects and the Universe on the whole) when all the internal resources of energy are exhausted. The analysis of such a process in the classical approximation is based on Einstein's equations (the classical gravitation and the classical matter). The classical picture of the

---

<sup>8</sup>Due to space-like nature of the pseudovector of the spin,  $(\bar{\Psi}\gamma^\mu\gamma_5\Psi)^2 \geq 0$ .

gravitational collapse is based of these equations, for which all the key evolution scenarios are established and classified: the equilibrium configurations, the cases when the contraction stops under the gravitational radius, and actually the case of a collapse when the matter is uncontrollably contracting under the influence of gravitational forces. During the contraction, the matter can become extremely dense (reaching the nuclear density  $10^{17}$  kg/m<sup>3</sup> and greater). Strictly speaking, at this stage it is necessary to use the Einstein equation with the source on the right-hand described by the averaged energy-momentum tensor of the quantum matter<sup>9</sup>. At the final stages of the gravitational collapse at the density greater than  $10^{99}$  kg/m<sup>3</sup> and the distance smaller than  $10^{-35}$  m, it is necessary to replace the classical equations by their quantum counterparts.

However, due to the great complexity of the problem of construction of the full quantum version of Einstein's equations (see Chapter 7), it is reasonable to consider the semi-consistent problem as a first approximation, when the gravity is treated classically and the matter - quantum-mechanically. The expediency of such approximation is obvious since already in this picture one can expect new features in the evolution of the gravitating configurations as compared to existing processes at a classical level.

The external classical non-stationary gravitational field may lead to the instability of vacuum of the quantized physical fields, i.e., to the production of the respective particles. As we show here, the torsion (within the framework of ECT) brings the new features to the particle generation process.

The classical dynamics of a massless scalar field  $\varphi$  will be described by the Klein-Gordon equation with the conformal coupling (4.41), with the Riemann-Cartan scalar curvature  $R(\Gamma)$  in place of  $R$ . The action for the scalar field has the following form:

$$S = \frac{1}{c} \int d^4x \sqrt{g} \mathcal{L}_\varphi, \quad \mathcal{L}_\varphi = -\frac{\hbar c}{2} \left( g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi + \frac{R(\Gamma)}{6} \varphi^2 \right). \quad (4.53)$$

We rewrite the metric (4.44) of the quasi-Euclidean (with  $\mathcal{K} = 0$ ) homogeneous isotropic Universe as

$$ds^2 = a^2(-d\eta^2 + \gamma_{ik} dx^i dx^k), \quad (4.54)$$

introducing the conformal time  $cdt/R(t) = d\eta$  and denoting the new scale factor  $a(\eta) = R(t(\eta))$ . The three-dimensional Euclidean metric  $\gamma_{ik}$  is not necessarily written in Cartesian coordinates. Furthermore, we assume that the evolution of the Universe is described by the cosmological solution (4.48).

We model the source of the external torsion field as a massless fermionic fluid described by the Dirac equation (4.18) (with  $m_\Psi = 0$ ) with the spin directed

---

<sup>9</sup>The averaged energy-momentum tensor of matter is usually understood as the result of the quantum averaging over the pure states, for example, over the space of occupation numbers. But because of the fact that the system during the evolution can come to a thermodynamic equilibrium, it seems to be necessary to make also a statistical averaging over the equilibrium distribution [75] (see Sec. 7.2.).

along the axis  $z$ . The torsion is then represented by the pseudotraces that has the form  $\tilde{Q}_\mu = (0, 0, 0, Q)$ , where  $Q = \kappa c \frac{\hbar}{2} \rho$  and  $\rho$  is the particle density of fermions. The conservation law of the number of particles in the metric (4.54) yields  $Qa^2 = \frac{\hbar}{2} \kappa c \rho_0$  with the integration constant  $\rho_0$  that gives the fermion particle density at the moment when  $a = 1$ .

Introducing a new variable for the scalar field  $\chi = a\varphi$ , we get the Lagrange function

$$L_\chi = \frac{\hbar c}{2} \int d^3x \sqrt{\gamma} \left\{ \dot{\chi}^2 + \chi \Delta \chi - \frac{Q_0^2}{a^4} \chi^2 \right\}. \quad (4.55)$$

Here the dot denotes the derivative with respect to the conformal time  $\eta$ , the 3-metric  $\gamma_{ik}$  determines the integration measure  $\sqrt{\gamma} = (\det \gamma_{ik})^{1/2}$  and the covariant Laplacian  $\Delta = \gamma^{ik} \overset{\{\}}{\nabla}_i \overset{\{\}}{\nabla}_k$ , and  $Q_0 = \kappa c \frac{\hbar}{2} \rho_0$  is the parameter with the dimension of 1/length which is proportional to the particle density of fermions that create the spacetime torsion.

Let us decompose the redefined scalar field

$$\chi(\mathbf{x}, \eta) = \sum_n \chi_n(\eta) Z_n(\mathbf{x}) \quad (4.56)$$

with respect to the complete orthonormal system of eigenfunctions  $Z_n(\mathbf{x})$

$$\left. \begin{aligned} \Delta Z_n(\mathbf{x}) &= -\omega_n^2 Z_n(\mathbf{x}), \\ \int d^3x \sqrt{\gamma} Z_n(\mathbf{x}) Z_m(\mathbf{x}) &= \delta_{nm}, \end{aligned} \right\} \quad (4.57)$$

of the 3-dimensional Laplace operator, where the collective index  $n$  labels the whole set of eigenvalues  $\omega_n^2$ . For the quasi-Euclidean model, the spectrum of the Laplacian is continuous, so the sum over the eigenmodes should be actually understood as an integral with an appropriate choice of the measure.

Substituting (4.56) and using (4.57), we rewrite the Lagrange function (4.55) in terms of an infinite set of discrete degrees of freedom of the scalar field  $\chi_n(\eta)$ :

$$L_\chi = \frac{\hbar c}{2} \sum_n \left\{ \dot{\chi}_n^2 - \omega_n^2 \chi_n^2 - \frac{Q_0^2}{a^4} \chi_n^2 \right\}.$$

The corresponding equation of motion

$$\ddot{\chi}_n(\eta) + \Omega_n^2(\eta) \chi_n(\eta) = 0, \quad \Omega_n^2(\eta) = \omega_n^2 + \frac{Q_0^2}{a^4(\eta)} \quad (4.58)$$

represents a discrete non-Riemannian version of (4.41).

Introducing as usual the canonical momenta<sup>10</sup>,  $p_n = \partial L_\chi / \partial (c\dot{\chi}_n) = \hbar \dot{\chi}_n$  we construct the Hamilton function

$$H_\chi = \sum_n p_n c \dot{\chi}_n - L_\chi = \frac{\hbar c}{2} \sum_n \left\{ (p_n / \hbar)^2 + \omega_n^2 \chi_n^2 + \frac{Q_0^2}{a^4} \chi_n^2 \right\}.$$

---

<sup>10</sup>In  $\dot{\chi}_n = d\chi_n/d\eta$  the variable  $\eta$  has the dimension of length, so we need a factor  $c$  to obtain the velocity  $c\dot{\chi}_n$  with the correct dimension.

The same Hamiltonian can be alternatively derived from the metrical energy-momentum tensor [71, 72]

$$T_{\mu\nu} = \hbar c \left\{ \partial_\mu \varphi \partial_\nu \varphi - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \partial_\alpha \varphi \partial_\beta \varphi + \frac{1}{6} \left( G_{\mu\nu} \varphi^2 - \overset{\{\}}{\nabla}_\mu \overset{\{\}}{\nabla}_\nu \varphi^2 + g_{\mu\nu} \overset{\{\}}{\square} \varphi^2 \right) + \overset{\check{}}{Q}_\mu \overset{\check{}}{Q}_\nu \varphi^2 - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \overset{\check{}}{Q}_\alpha \overset{\check{}}{Q}_\beta \varphi^2 \right\}. \quad (4.59)$$

The Hamilton function is then recovered as the integral

$$H_\chi = a^2 \int d^3x \sqrt{\gamma} T_{00}. \quad (4.60)$$

The scalar field is quantized in a standard way by treating  $\chi_n(\eta)$  and the canonically conjugated momenta  $p_n(\eta)$  as operators satisfying the equal-time commutation relations

$$[\chi_m(\eta), \chi_n(\eta)] = 0, \quad [p_m(\eta), p_n(\eta)] = 0, \quad [\chi_n(\eta), p_m(\eta)] = i\hbar \delta_{nm}. \quad (4.61)$$

To introduce the particle interpretation, we decompose the operator  $\chi_n(\eta)$  into the two complex-conjugate solutions  $(u_n, u_n^*)$  of the equations (4.58):

$$\chi_n(\eta) = a_n u_n(\eta) + a_n^* u_n^*(\eta). \quad (4.62)$$

The basis functions  $(u_n, u_n^*)$  are chosen so that they satisfy the normalization conditions

$$u_n \dot{u}_n^* - \dot{u}_n u_n^* = i. \quad (4.63)$$

The creation  $a_n^*$  and annihilation  $a_n$  operators satisfy

$$[a_n, a_m] = 0, \quad [a_n^*, a_m^*] = 0, \quad [a_n, a_m^*] = \delta_{nm}.$$

These standard commutation relations follow from (4.61)-(4.63).

A detailed analysis of the particle interpretation of the system can be found in [61]. Inserting (4.62) into the Hamilton function, we find

$$H_\chi = \frac{\hbar c}{2} \sum_n \Omega_n(\eta) \{ E_n(a_n a_n^* + a_n^* a_n) + F_n(a_n)^2 + F_n^*(a_n^*)^2 \}, \quad (4.64)$$

where we denoted the dimensionless functions

$$E_n(\eta) = \frac{|\dot{u}_n|^2}{\Omega_n(\eta)} + \Omega_n(\eta) |u_n|^2, \quad F_n(\eta) = \frac{(\dot{u}_n)^2}{\Omega_n(\eta)} + \Omega_n(\eta) (u_n)^2. \quad (4.65)$$

Note that  $(E_n)^2 - |F_n|^2 = 1$  in view of (4.63).

We choose the initial conditions  $\{u_n, u_n^*, \dot{u}_n, \dot{u}_n^*\}_{|\eta=\eta_0}$  at an arbitrary moment of time  $\eta = \eta_0$  so that to satisfy (4.63) and to make the Hamilton function (4.64) diagonal by setting  $E_n(\eta_0) = 1$  and  $F_n(\eta_0) = 0$ :

$$u_n(\eta_0) = u_n^*(\eta_0) = \frac{1}{\sqrt{2\Omega_n(\eta_0)}}, \quad \dot{u}_n(\eta_0) = -\dot{u}_n^*(\eta_0) = -i\sqrt{\frac{\Omega_n(\eta_0)}{2}}.$$

The operators  $a_n$  and  $a_n^*$  are the creation and annihilation operators of scalar particles at the initial moment  $\eta_0$ . This interpretation is also preserved for any  $\eta$  if  $\Omega_n$  is constant, then the Hamiltonian remains diagonal at all times. However, for the time-dependent  $\Omega_n(\eta)$  the Hamilton function (4.64) ceases to be diagonal in terms of  $a_n$  and  $a_n^*$  at any time  $\eta > \eta_0$ . This means that the physical vacuum is unstable and hence the notion of a particle is not constant but depends on time, which is manifest in the production of scalar particles.

In GR this effect is absent in the conformally-flat space-time (4.54) for the model under consideration due to the conformal invariance of the scalar field theory. In the ECT, the particle creation effect is nontrivial which is explained by the fact that the fermion torsion sources violate the conformal invariance of the field  $\varphi$ , as we noted in Sec. 4.2.

Thus, at an initial time  $\eta = \eta_0$ , we have the particle interpretation of  $\varphi$ . The field energy in  $n$ -mode is  $\varepsilon_n = \frac{1}{2}\hbar c \Omega_n(\eta_0) (a_n a_n^* + a_n^* a_n)$ . In order to find the particle interpretation for an arbitrary time, we use the diagonalization method of the Hamilton function (4.64). In accordance with this method, it is necessary to introduce the new (time-dependent) creation and annihilation operators  $A_n^*(\eta)$  and  $A_n(\eta)$  by

$$a_n = \alpha_n^*(\eta) A_n(\eta) + \beta_n(\eta) A_n^*(\eta), \quad a_n^* = \alpha_n(\eta) A_n^*(\eta) + \beta_n^*(\eta) A_n(\eta), \quad (4.66a)$$

where the coefficients  $\alpha_n(\eta)$  and  $\beta_n(\eta)$  satisfy the conditions

$$|\alpha_n(\eta)|^2 - |\beta_n(\eta)|^2 = 1, \quad F_n \beta_n = (1 - E_n) \alpha_n. \quad (4.66b)$$

In terms of these operators, the Hamilton function (4.64) has diagonal form

$$H_\chi = \frac{\hbar c}{2} \sum_n \Omega_n(\eta) \{A_n^*(\eta) A_n(\eta) + A_n(\eta) A_n^*(\eta)\}.$$

At any given time, the field particles are determined by the time-dependent creation  $A_n^*(\eta)$  and annihilation  $A_n(\eta)$  operators. By construction, we have  $\alpha_n(\eta_0) = 1, \beta_n(\eta_0) = 0$  and hence the operators coincide  $A_n(\eta_0) = a_n$ , and  $A_n^*(\eta_0) = a_n^*$  at the initial moment of time  $\eta_0$ .

The instantaneous physical vacuum  $|0_\eta\rangle$  is defined as the quantum state that satisfies  $A_n(\eta)|0_\eta\rangle = 0$ . For an arbitrary operator  $\Phi$  we introduce the normal ordering

$$\mathcal{N}_\eta(\Phi) = \Phi - \langle 0_\eta | \Phi | 0_\eta \rangle.$$

This obviously depends on time: one subtracts the vacuum average with respect to the instantaneous vacuum, thus removing the (usually diverging) contribution of the vacuum excitations a  $\eta$ . In order to find the physical effects due to the particle production, one needs to compute vacuum averages with respect to the initial vacuum:

$$\langle 0_{\eta_0} | \mathcal{N}_\eta(\Phi) | 0_{\eta_0} \rangle = \langle 0_{\eta_0} | \Phi | 0_{\eta_0} \rangle - \langle 0_\eta | \Phi | 0_\eta \rangle.$$

Particularly important are the operators of number of particles  $A_n^*(\eta)A_n(\eta)$  (in the  $n$ -th mode) and the operators of the energy-momentum  $T_i^k(x)$ . The latter is given explicitly in (4.59). Direct computation of the vacuum averages of the normal ordered operators yields:

$$\left. \begin{aligned} n(\eta) &= \langle 0_{\eta_0} | \mathcal{N}_\eta(A_n(\eta)A_n^*(\eta)) | 0_{\eta_0} \rangle = \sum_n |\beta_n|^2, \\ \mathcal{E}(\eta) &= -\langle 0_{\eta_0} | \mathcal{N}_\eta(T_0^0) | 0_{\eta_0} \rangle = \frac{\hbar c}{a^4(\eta)} \sum_n \Omega_n(\eta) |\beta_n|^2, \\ \mathcal{P}_1(\eta) &= \langle 0_{\eta_0} | \mathcal{N}_\eta(T_1^1) | 0_{\eta_0} \rangle = \frac{1}{3} \{ \mathcal{E}(\eta) - \gamma(\eta) \}, \\ \mathcal{P}_2(\eta) &= \langle 0_{\eta_0} | \mathcal{N}_\eta(T_2^2) | 0_{\eta_0} \rangle = \frac{1}{3} \{ \mathcal{E}(\eta) - \gamma(\eta) \}, \\ \mathcal{P}_3(\eta) &= \langle 0_{\eta_0} | \mathcal{N}_\eta(T_3^3) | 0_{\eta_0} \rangle = \frac{1}{3} \{ \mathcal{E}(\eta) + 2\gamma(\eta) \}, \end{aligned} \right\} \quad (4.67)$$

where

$$\gamma(\eta) = \frac{\hbar c Q_0^2}{a^8(\eta)} \sum_n \left\{ |u_n|^2 - \frac{1}{2\Omega_n(\eta)} \right\}.$$

One can verify that for the off-diagonal components of the energy-momentum tensor, the vacuum averages vanish,  $\langle 0_{\eta_0} | \mathcal{N}_\eta(T_i^k) | 0_{\eta_0} \rangle = 0$  with  $i \neq k$ . The total number of particles  $n(\eta)$  created from vacuum by the torsion field, as well as the energy density  $\mathcal{E}(\eta)$  and the anisotropic pressure  $\mathcal{P}_1(\eta), \mathcal{P}_2(\eta), \mathcal{P}_3(\eta)$  of the created matter are determined by the solutions  $u_n(\eta)$  of the evolution equation (4.58). From (4.67), one can obtain the equation of state for the created matter

$$\mathcal{E} = \mathcal{P}_1 + \mathcal{P}_2 + \mathcal{P}_3.$$

It is impossible to solve the oscillator equation (4.58) exactly for the time-dependent frequency  $\Omega_n(\eta)$  with an arbitrary cosmological evolution  $a = a(\eta)$ . So, in order to obtain numerical estimates for  $(n, \mathcal{E}, \mathcal{P}_i)$ , one needs a perturbation scheme to construct approximate solutions of the equation (4.58) with the given initial conditions. As a first step, we recast the differential equation (4.58) into the equivalent integral equation of Volterra type:

$$u_n(\eta) = \frac{e^{-i\Omega_n(\eta_0)(\eta-\eta_0)}}{\sqrt{2\Omega_n(\eta_0)}} + \int_{\eta_0}^{\eta} d\xi G(\eta - \xi) \Delta Q^2(\xi) u_n(\xi),$$

where  $\Delta Q^2(\eta) = Q^2(\eta_0) - Q^2(\eta) = Q_0^2 \{ 1/a^4(\eta_0) - 1/a^4(\eta) \}$ , and  $G(\eta - \xi)$  is the Green function that satisfies

$$\ddot{G}(\eta - \xi) + \Omega_n^2(\eta_0)G(\eta - \xi) = \delta(\eta - \xi).$$

The resulting integral equation is then solved iteratively, and the solution reads

$$\left. \begin{aligned} u_n(\eta) &= u_n^{(0)}(\eta) + u_n^{(1)}(\eta) + \dots + u_n^{(m)}(\eta) + \dots, \\ u_n^{(0)}(\eta) &= \frac{e^{-i\Omega_n(\eta_0)(\eta-\eta_0)}}{\sqrt{2\Omega_n(\eta_0)}}, \\ u_n^{(m)}(\eta) &= \int_{\eta_0}^{\eta} d\xi G(\eta-\xi) \Delta Q^2(\xi) u_n^{(m-1)}(\xi). \end{aligned} \right\} \quad (4.68)$$

We take  $G(\eta-\xi) = \frac{\sin\{\Omega_n(\eta_0)(\eta-\xi)\}}{\Omega_n(\eta_0)}$  as the explicit form of the Green function.

We choose the initial time  $\eta_0$  as the moment when the scale factor  $a(\eta)$  takes the minimal value  $a(\eta_0) = a_{\min} = R_{\min}$  (see Sec. 4.3.). Limiting ourselves to the first two terms in (4.68), from (4.67), (4.68) we can obtain the numerical estimates for the particle number, the energy and the pressure of the created matter. In the lowest order we find

$$\left. \begin{aligned} n(\eta) &= \sum_n \frac{(\Omega_n(\eta) - \Omega_n(\eta_0))^2}{2\Omega_n(\eta)\Omega_n(\eta_0)}, \\ \mathcal{E}(\eta) &= \frac{\hbar c}{a^4(\eta)} \sum_n \frac{(\Omega_n(\eta) - \Omega_n(\eta_0))^2}{2\Omega_n(\eta_0)}, \\ \gamma(\eta) &= \frac{\hbar c Q_0^2}{a^8(\eta)} \sum_n \frac{\Omega_n(\eta) - \Omega_n(\eta_0)}{2\Omega_n(\eta)\Omega_n(\eta_0)}. \end{aligned} \right\} \quad (4.69)$$

The details of the computations can be found in [78]. Evaluation of the sums in (4.69) (more exactly, of the integrals) is a nontrivial task. Here we summarize the qualitative results.

1. The maximal number of created particles corresponds to the scale factor  $a_0 \approx \sqrt{2}a_{\min}$  and the density of created particles is equal to  $\approx Q_0^3/a_{\min}^3$ , that is directly related to the density of fermions generating the spacetime torsion.

2. At the large distances (far from the Planck region  $a > a_0$ ), the intensity of the scalar particle production by the torsion field drops. Conversely, due to the high power of the inverse scale factor in the expressions for  $n(\eta)$ ,  $\mathcal{E}(\eta)$ , at the small distances the process of scalar particle production by the torsion field dominates over the similar processes of the matter particle production in GR.

3. The account of the back reaction of the created particles on the classical evolution of the isotropic Universe should lead to its anisotropization in the early epochs due to an extra contribution  $\gamma(\eta)$  making the pressure anisotropic. The estimate of the maximal anisotropization degree yields the times close to  $a_0$ . At the large distances (late epochs with  $a \rightarrow \infty$ ),  $\gamma(\eta)$  quickly decreases to zero. As a result, during Universe's evolution, the isotropy is recovered.

# 5

## Kinematics of gauge theories of gravity

### 5.1. Special aspects of gauge approach in gravitation

Mathematical beauty of the classical GR as a dynamical theory of a three-dimensional spatial geometry, as well as its agreement with the present-day experimental data in cosmology, astrophysics and planetary astronomy, allow one to view GR as a fair description of the macroscopic gravitational phenomena. However, the existence of essential difficulties in it (such as the quantization problem [92] and the singularity problem [86]) at small distances apparently manifests its limited validity in the microworld.

Recently numerous attempts were made to formulate the microscopic gravity theory within the framework of a consistent gauge-theoretic approach. This was essentially motivated by the successful construction of renormalizable unified gauge models of the weak and electromagnetic, as well as of the strong interactions. On the other hand, construction of the microscopic gravity theory appears to be a necessary step to establish the geometrical structure of supergravity [93] as the theory of the local supersymmetry.

The gravity theory, understood in a broad sense as the geometrodynamics<sup>1</sup> of the spacetime, is much richer and more complex in the differential-geometrical aspects than the gauge theories of internal symmetries. As a result, there exists a variety of approaches and interpretations of the gravitational field as a gauge one, and there is no clear understanding of the answer to the key question: is gravity a gauge theory and in which sense? This question includes the three

---

<sup>1</sup>That is the dynamics of the geometrical structure of spacetime.

aspects [97]: 1) What is the gauge group of the theory; what is the role of the general coordinate transformations? 2) What are the relevant gauge fields; what is their relation with the geometrical objects on the spacetime manifold; in particular, what is the status of the metric? 3) What is theory's dynamics; is it determined by the requirements of the gauge invariance?

In view of these issues, it is reasonable to consider in detail the difficulties arising in the development of a consistent gauge approach to the gravity theory.

To begin with, we have to clarify the terminology, because different researchers understand the meaning of expressions “gauge field” and “gauge theory” in different ways. We define the gauge theory as a physical theory in which the fundamental dynamical variable—the gauge field—is a connection on a principal bundle. Among such theories, we distinguish the consistent models that do not contain any additional postulates of the non-gauge type from the inconsistent theories containing such assumptions<sup>2</sup>. Besides that, certain theories that are not covered by this definition, still have some features of the gauge theories, such as the presence of arbitrary functions in the description of the fields and the existence of constraints; we call them pseudo-gauge theories. There are many examples: the theory of massless fields of arbitrary spin; generally covariant models of material fields in a curved space; the simple supergravity [93]; the general relativity theory [35]. Quantization of the gauge and pseudo-gauge theories takes into account their common feature of the invariance with respect to an infinite group of local transformations, and this similarity brings in a terminological confusion, which results in the unfortunate statements, such as “GR is the gauge theory of the group of general coordinate transformations”.

The issue of finding the gauge group for the gravity theory is one of the most complicated ones, since the gravity is related to spacetime symmetries, in contrast to the gauge fields of “internal” symmetries. However, being well-defined as global symmetries in the flat Minkowski space, the “localized” spacetime symmetries become general coordinate transformations, losing their original features. Moreover, there is a difficulty of interpretation of the “local” translations which are usually identified with the infinitesimal general coordinate transformations [12, 13]. The corresponding “gauge fields”—the tetrads—do not have a standard (Yang-Mills) transformation law, since they are not connections.

As it is clear now, all these and related inconsistencies arise from the confusion of the gauge structure with the coordinate invariance. However, they should be carefully distinguished, because the group of general coordinate transformations does not have a direct relation to the gauge theory of gravity [96]. If one views the general coordinate transformations simply as the maps between charts covering the spacetime manifold, they obviously do not carry any dynamical meaning. The invariance with respect to such passive transformations is a *common property* of all covariant physical theories, not only of gravity. More-

---

<sup>2</sup>An example of the latter is an ad hoc postulate of the non-dynamical vanishing of the torsion of a linear connection (see below).

over, using modern coordinate-free methods for the description of the physical systems (the exterior calculus), one can provide the general covariance without any appeal to the gravity theory. There is also another point of view on general coordinate transformations as the group of local diffeomorphisms of the spacetime manifold. These are active transformations due to the motion of spacetime points. But also in this case, the study of the invariance properties of physical systems is not related to gravity since the corresponding mathematical formalism of the Lie derivatives along vector fields, defined by diffeomorphisms, does not demand the existence of a geometrical structure on the manifold.

Below we show that it more natural to describe the action of the spacetime symmetries not on the base –the spacetime itself– but rather in bundle of affine or linear frames over it. Then the gravitational field turns out to be a connection in this principal fiber bundle. There is a natural homomorphism from the bundle of frames to the tangent bundle, using which the gauge gravitational field induces geometrical structures on the spacetime: the connection and the metric. The issue of the status of the metric in this approach has an unexpected dual solution. Namely, the metric structure is explained with the help of a spontaneous symmetry breaking mechanism when the reduction of the inhomogeneous general affine group  $GA(4, R)$  to linear group  $GL(4, R)$  allows to interpret the tetrad field as a nonlinear realization of the connection, and on the other hand, the reduction of  $GL(4, R)$  to the Lorentz group adds a Goldstone contribution to the metric. Roughly speaking, the metric turns out to be some hybrid of the two popular interpretations – the translational gauge field (in the nonlinear realization) and the Goldstone field taking values in the homogeneous space  $GL(4, R)/SO(3, 1)$ .

In the consistent gauge approach to the gravity theory, the largest difficulties arise when constructing its dynamics. This is explained by an essentially non unique definition of an action by the gauge invariance. In chapter 6, we will consider some models, for which the study of the physical properties can lead to the final establishment of the structure of the gauge gravity theory.

## 5.2. Bundle of frames and generalized affine connection

We construct the consistent gauge scheme for the gravitational field using the mathematical theory of the generalized affine and linear connections in the bundles of frames over the spacetime  $M_4$ . We understand the latter as the four-dimensional smooth ( $C^\infty$ ) manifold. When discussing the geometrical aspects of the bundle spaces, we use the local coordinates writing the basic objects in components and defining their transformation laws. Certainly, we then lose the advantages of coordinate-free methods that make a stress on the global invariant meaning of various operations, however, the component formalism is more familiar to physicists. A brief summary of the basic mathematical definitions (manifold, forms, bundles, connection, etc.) is given in Appendix A1.

### *Bundle of linear frames*

A linear frame at the point  $x \in M_4$  is the basis  $\{e_a\} = \{e_0, e_1, e_2, e_3\}$  of the tangent space  $T_x M$ . The set of frames  $G_x$  at this point is isomorphic to the general linear group  $GL(4, R)$ , since any frame is obtained by a linear transformation from the standard basis  $E_a = \{E_1 = (1, 0, 0, 0), E_2 = (0, 1, 0, 0), E_3 = (0, 0, 1, 0), E_4 = (0, 0, 0, 1)\}$  in  $R^4 \cong T_x M$ :  $e_a = L^b{}_a E_b$  with the  $4 \times 4$  matrix  $L \in GL(4, R)$ . The union  $\bigcup_{x \in M_4} G_x = L(M)$ , provided with a smooth

structure, is a  $(4 + 4^2)$ -dimensional manifold, where the natural projection  $\pi : L(M) \rightarrow M$  maps the linear frame  $e_a$  at the point  $x$  to  $x$ . The manifold  $L(M)$  is called the bundle of linear frames. This is the principal bundle with the structural group  $GL(4, R)$ ; the right action  $GL(4, R)$  on  $L(M)$  is evident:  $e_a \in G_x \rightarrow e'_b = e_a L^a{}_b \in G_x$  for  $L \in GL(4, R)$ . The tangent bundle  $T(M)$  may be viewed as an associated with  $L(M)$  bundle with the standard fiber  $R^4$ .

The connection in  $L(M)$  is called the linear connection in  $M_4$ . It introduces the parallel transport in  $L(M)$  and induces the parallel transport of the tangent vectors in the associated  $T(M)$  and defines the covariant differentiation of the vector fields. Choosing the local coordinates  $\{x^\mu\}$  on  $M_4$  and a cross-section  $\sigma : M \rightarrow L(M)$ , we find the 1-form on the base  $\sigma^* \omega = \omega^b{}_{a\mu} E^a{}_b dx^\mu$  which is uniquely determined by the linear connection form  $\omega$  on  $L(M)$  with the values in the Lie algebra  $gl(4, R)$ . Here  $E^a{}_b$  is the natural basis of the Lie algebra  $gl(4, R)$ , i.e., the  $4 \times 4$  matrix with elements at the intersection of  $a$ -th column and  $b$ -th line equal to 1, and with the remaining elements equal zero. The coefficients  $\omega^b{}_{a\mu}$  of this form are called the coefficients of the local linear connection. They define the coordinate notation of the covariant derivative of the tangent vector  $\nabla_\mu v^a = \partial_\mu v^a + \omega^a{}_{b\mu} v^b$ , where the vector  $v = v^a e_a$  is decomposed with respect to the basis  $e_a(x)$  that is defined by the cross-section  $\sigma$ . When the cross-section is changed,  $\sigma \rightarrow \sigma'$ , the connection coefficients are transformed as follows:

$$\omega^a{}_{b\mu} \longrightarrow \omega'^a{}_{b\mu} = L^a{}_c \omega^c{}_{d\mu} L^{-1d}{}_b + L^a{}_c \partial_\mu L^{-1c}{}_b, \quad (5.1)$$

where the matrix of the linear transformation is determined from

$$e'_a(x) = e_b(x) L^b{}_a(x).$$

In the bundle of linear frames  $L(M)$ , one can introduce (absolutely independently from the connection) another basic object – the canonical 1-form  $\theta$ . This  $R^4$ -valued form is defined on  $L(M)$  by the equation

$$d\pi(X) = \theta^a(X) e_a, \quad (5.2)$$

where  $X \in T_u(L(M))$ ,  $u = (x, e_a)$ . The canonical 1-form is horizontal, i.e., it vanishes on vertical (tangent to the fiber) vectors

$$\theta^a(X) = 0 \quad \Longleftrightarrow \quad d\pi(X) = 0.$$

This property is obvious. In addition,  $\theta$  is equivariant, i.e., for any  $L \in GL(4, R)$

$$R_L^* \theta^a \equiv \theta^a \circ dR_L = L^{-1a}{}_b \theta^b.$$

Indeed, let  $u = (x, e_a) \in L(M)$ , and  $d\pi(X) = \theta^a(X)e_a$ . Then the right action  $R_L u = (x, e_a L^a_b)$ , and its differential  $dR_L(X) \in T_{R_L u} L(M)$ , as a result,

$$(R_L^* \theta^a) e_a = d\pi(dR_L X) = d\pi(X) = (\theta^a(X) L^{-1c}_a) (L^b_c e_b).$$

In the local notation, the canonical form determines the coframe 1-form  $h^a : T_x(M) \rightarrow R^4$ , i.e., the basis of the cotangent space  $T_a^* M$  dual to  $e_a$ . If the cross-section is chosen  $\sigma : M \rightarrow L(M)$  [i.e. the field of frames  $e_a(x)$ ], then we have  $h^a(x) = (\sigma^* \theta^a)(x)$ , where  $h^a(e_b) = \delta^a_b$ .

With the connection  $\omega$  and the canonical form  $\theta$  defined on the bundle of linear frames, one can introduce a parallelization of  $L(M)$ . Namely, one can: 1) map each element of the Lie algebra  $gl(4, R)$  to a fundamental<sup>3</sup> vertical vector field; in particular, map the basis of the Lie algebra  $gl(4, R)$ :  $E^a_b \rightarrow E^{*a}_b$ , so that  $\omega(E^{*a}_b) = E^a_b$ , and 2) map each vector from  $R^4$  to a standard horizontal vector field, in particular map the basis  $E_a \rightarrow E_a^*$ , so that  $\theta(E_a^*) = E_a$ . Then  $4 + 4^2$  vector fields  $E_a^*$ ,  $E^{*a}_b$  constitute at each point  $u \in L(M)$  the basis of the tangent space  $T_u(L(M))$ .

The linear connection in  $L(M)$  is characterized by the curvature and the torsion. Let us define the curvature 2-form  $R$  of the linear connection  $\omega$  as covariant exterior differential<sup>4</sup>

$$R = D\omega = d\omega(h). \quad (5.3)$$

The torsion 2-form  $Q$  of the linear connection  $\omega$  is defined as the covariant exterior differential of the canonical form  $\theta$ :

$$Q = D\theta = d\theta(h). \quad (5.4)$$

The 2-forms  $R$  and  $Q$  satisfy the structure equations

$$\left. \begin{aligned} Q^a &= d\theta^a + \omega^a_b \wedge \theta^b, \\ R^a_b &= d\omega^a_b + \omega^a_c \wedge \omega^c_b, \end{aligned} \right\} \quad (5.5)$$

where we made the expansion with respect to the natural bases in  $R^4$  and  $gl(4, R)$ :  $Q = Q^a E_a$ ,  $\theta = \theta^a E_a$ ,  $R = R^a_b E^b_a$ .

Introducing the local coordinates  $x^\mu$  on the manifold  $M_4$ , we naturally define the corresponding cross-section in  $L(M)$  by attaching the coordinate basis  $\{\partial_\mu\}$  at  $x \in M_4$ . Then the local coordinates in  $L(M)$  are  $(x^\mu, h^a_\mu)$ , where  $e_a = h^a_\mu \partial_\mu$ , and it is obvious that  $\det h^a_\mu \neq 0$ . Determining the inverse matrix  $h^a_\mu$  so that  $h^a_\mu h^b_\mu = \delta^a_b$ , we obtain from the definitions (5.2)-(5.4) the local expressions for the canonical 1-form

$$\theta^a = h^a_\mu dx^\mu \quad (5.6)$$

<sup>3</sup>For definitions, see Appendix A3 and references therein.

<sup>4</sup>In (5.3) and (5.4),  $h$  denotes the projection to the horizontal subspace of  $TL(M)$ .

and for the connection 1-form

$$\omega^a{}_b = h_\alpha^a \Gamma^\alpha{}_{\beta\nu} h_b^\beta dx^\nu + h_\mu^a dh_b^\mu. \quad (5.7)$$

Then using the structure equations (5.5), we arrive at the ordinary tensor expressions for the torsion (1.3) and the curvature (1.1), constructed from the components of the linear connection  $\Gamma^\alpha{}_{\beta\mu}$ .

### *Bundle of affine frames*

The tangent space  $T_x(M) \cong R^4$  may be considered as the affine space  $A^4$ . In this case, we call it the tangent affine space and denote  $A_x(M)$ . Its basis, which is called the affine frame, includes the point  $z \in A_x(M)$  and linear frame  $e_a$ . The set of affine frames at the point  $x \in M_4$  is isomorphic to general affine group  $GA(4, R)$ , since an arbitrary affine frame  $(z, e_a)$  is obtained using the affine transformation  $A \in GA(4, R)$  from the natural frame  $(0, E_a)$  in  $A^4$ . The set of all affine frames on  $M$  is called the bundle of affine frames  $A(M)$ . The projection  $\pi : A(M) \rightarrow M$  maps a frame at the point  $x$  to this point  $x \in M_4$ . The bundle  $A(M)$  is the principal bundle over  $M$  with the structural group  $GA(4, R)$  with an obvious definition of the group action.

A natural realization of affine transformations in  $A_x(M)$  is obtained, if we view the affine space  $A^4$  as a hypersurface in  $R^5$  with a fixed fifth coordinate.

Then an arbitrary element of  $A^4$  is represented as a column  $\begin{pmatrix} z \\ 1 \end{pmatrix}$ , and the action

of the affine group  $GA(4, R)$  in the affine space  $A^4$  is described by the  $5 \times 5$

matrix of the form  $g = \begin{pmatrix} L & b \\ 0 & 1 \end{pmatrix}$ , where  $L \in GL(4, R)$  and  $b \in R^4$ :

$$\begin{pmatrix} z \\ 1 \end{pmatrix} \longrightarrow \begin{pmatrix} z' \\ 1 \end{pmatrix} = \begin{pmatrix} L & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ 1 \end{pmatrix} = \begin{pmatrix} Lz + b \\ 1 \end{pmatrix}.$$

The bundle of affine frames  $A(M)$  and the bundle of linear frames  $L(M)$  are closely related to each other and to the associated tangent bundles. In particular, the affine tangent bundle represents an important example of a soldered bundle, the properties of which are reviewed below.

We say that the bundle  $E(M, F, G, P)$  associated with the principal bundle  $P(M, G)$  is soldered to the base  $M$ , if [3]:

1) the fiber  $F$  of the bundle  $G$  is a homogeneous factor-space  $F = G/H$ , where  $H \subset G$  is a stationary subgroup (or stabilizer), and the dimension of  $F$  is equal to dimension of the base  $M$ ;

2) there is a global cross-section  $\sigma : M \rightarrow E$ , and there is a natural isomorphism between the tangent bundle  $T(M)$  and the space of all vectors tangent to the fibers  $F$  at the points of the cross-section  $\sigma(M)$ . In other words, the "soldering" of the bundle  $E$  to the base  $M$  means that the fibers  $F$  are tangent

to the manifold  $M$  (in the sense of coincidence of the tangent spaces) at the soldering point set by the cross-section  $\sigma$ . Furthermore, the bundle  $\tilde{T}(M)$  of the tangent spaces  $T_{\sigma(M)}F$  may be considered as a vector bundle associated with the principal bundle  $\tilde{P}(M, H)$ , obtained as a submanifold in  $P(M, G)$  when homeomorphisms  $F \rightarrow F_x$  are such that the “center” of  $F$  is mapped into  $\sigma(M)$  – the soldering point.

Let us consider the most important case when the fiber  $F$  of the soldered bundle  $E$  is a weakly reductive homogeneous space. This means that the Lie algebra  $\mathcal{G}$  of the structural group  $G$  is decomposed into the sum of the subalgebra  $\mathcal{H}$  (corresponding to the subgroup  $H$ ) and the vector space  $V$

$$\mathcal{G} = \mathcal{H} \oplus V \quad (5.8)$$

so that  $[\mathcal{H}, \mathcal{H}] \subset \mathcal{H}$ ,  $[\mathcal{H}, V] \subseteq V$ , where  $[\ , \ ]$  is the commutator in the Lie algebra. The tangent space  $T_{\sigma(M)}F$  may be identified with  $V$ .

A  $V$ -valued 1-form  $\tilde{\theta}$  on the principal bundle  $\tilde{P}(M, H)$  is called a soldering form if it satisfies the following conditions:

1)  $\tilde{\theta}(\tilde{X}) = 0 \iff d\tilde{\pi}(\tilde{X}) = 0$ , where  $\tilde{X} \in T\tilde{P}$ , i.e., it vanishes on vertical vectors;

2) under the right action  $R_h^*\tilde{\theta} \equiv \tilde{\theta} \circ dR_h = h^{-1}\tilde{\theta}h$ , where  $h \in H$ .

One can show [3] that the existence of the soldering 1-form  $\tilde{\theta}$  is necessary and sufficient condition for the bundle  $E$  to be soldered to  $M$ .

Now we consider in details the relation of the bundles of affine and linear frames. It is obvious that  $L(M)$  may be realized as a natural subbundle  $A(M)$ . The corresponding embedding  $\gamma : L(M) \rightarrow A(M)$  is defined by the map

$$\varphi : A(M) \longrightarrow R^4, \quad (5.9)$$

such that

$$R_{g^{-1}}^* \circ \varphi^a = A^a{}_b \varphi^b + b^a, \quad g = \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} \in GA(4, R).$$

Then the subbundle  $L(M)$  is determined as a kernel

$$L(M) = \{u \in A(M) \mid \varphi(u) = 0\}.$$

The canonical projection  $\beta : A(M) \rightarrow L(M)$  can be defined as

$$\beta(u) = R_{\Phi(u)} \circ (u), \quad \Phi(u) = \begin{pmatrix} \mathbf{1} & \varphi(u) \\ 0 & 1 \end{pmatrix}, \quad u \in A(M),$$

where  $\mathbf{1}$  is the unit in  $GL(4, R)$ . It is obvious that  $\beta \circ \gamma = id_{L(M)}$ . In local coordinates, the embedding of  $L(M)$  into  $A(M)$  maps the linear frame  $(x, e_a)$  to the affine frame  $(x, \mathcal{O}_x, e_a)$ , where  $\mathcal{O}_x$  is the origin of the affine tangent

space  $A_x$ . Conversely, the projection  $\beta : (x, z, e_a) \rightarrow (x, e_a)$ . Using the above-mentioned realization of the general affine transformations, and consequently, of the affine frames as a  $5 \times 5$  matrix of the following form:

$$u = \begin{pmatrix} L & z \\ 0 & 1 \end{pmatrix}, \quad L \in GL(4, R), \quad z \in R^4,$$

we get explicitly the mapping  $\varphi: A(M) \rightarrow R^4$ . For  $(x, u) \in A_x$ , it is defined as

$$\varphi(u) = -L^{-1}z.$$

The matrix inverse to  $g = \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix}$  reads  $g^{-1} = \begin{pmatrix} A^{-1} & -A^{-1}b \\ 0 & 1 \end{pmatrix}$ , and it is easy to show that under the right group action

$$R_{g^{-1}}u = ug^{-1} = \begin{pmatrix} L & z \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A^{-1} & -A^{-1}b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} LA^{-1} & -LA^{-1}b + z \\ 0 & 1 \end{pmatrix}$$

we have the correct transformation property

$$\begin{aligned} R_{g^{-1}}^* \circ \varphi(u) &\equiv \varphi(R_{g^{-1}}u) = -(LA^{-1})^{-1}(z - LA^{-1}b) \\ &= -AL^{-1}z + b = A\varphi(u) + b. \end{aligned}$$

Defined as the kernel  $\varphi(u) = 0$ , the subbundle  $L(M)$  is formed by the set of matrices  $\begin{pmatrix} L & 0 \\ 0 & 1 \end{pmatrix} \in GL(4, R)$ , which is indeed the bundle of linear frames. The canonical projection is also quite transparent in the matrix representation:

$$\begin{aligned} \beta(u) &= R_{\Phi(u)} \circ u = \begin{pmatrix} L & z \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \mathbf{1} & \varphi(u) \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} L & L\varphi(u) + z \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} L & 0 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

The connection in the bundle of affine frames  $A(M)$  is called a generalized affine connection in  $M_4$ . The generalized connection 1-form on  $A(M)$  is denoted as  $\Omega$ . Using the above-mentioned reduction of  $A(M)$  to  $L(M)$ , one can reduce the study of the generalized affine connection to the investigation of the linear connection and some  $R^4$ -valued 1-form on the bundle of linear frames  $L(M)$ .

Since the connection 1-form in the principal bundle is uniquely restored in terms of the set of the local forms on the base (satisfying certain compatibility condition in the intersections of local maps) [3], we will consider everything in the local coordinates. Given a cross-section  $\sigma: M \rightarrow A(M)$ , we define the

field of affine frames  $u(x) = \begin{pmatrix} L(x) & z(x) \\ 0 & 1 \end{pmatrix}$  on  $M_4$ , and the generalized affine connection 1-form  $\Omega$  induces the form  $\Omega_\sigma = \sigma^*\Omega$  on the base space. Since the Lie algebra of the general affine group is decomposed into the (semidirect) sum

$$ga(4, R) = gl(4, R) + R^4,$$

the local connection form  $\Omega_\sigma$  also can be decomposed into the sum  $\Omega_\sigma = \tilde{\omega} + \tilde{\varphi}$ . It is more convenient to use the matrix representation for the Lie algebra

$$ga(4, R) = \left\{ \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix}, \quad a \in gl(4, R), \quad b \in R^4 \right\}$$

and write the local connection form as the  $5 \times 5$  matrix

$$\Omega_\sigma = \begin{pmatrix} \tilde{\omega} & \tilde{\varphi} \\ 0 & 0 \end{pmatrix}. \quad (5.10)$$

Now we will show that this generalized affine connection determines the linear connection and some tensor form in  $M_4$ . For that purpose, we consider the transformation of the connection (5.10) under the change of the cross-section  $\sigma \rightarrow \sigma'$ . Then we have

$$u'(x) = R_{g(x)}u(x) = u(x) \begin{pmatrix} A(x) & b(x) \\ 0 & 1 \end{pmatrix}$$

and  $\Omega_{\sigma'} = g\Omega_\sigma g^{-1} + gdg^{-1}$ . Explicitly,

$$\begin{aligned} \begin{pmatrix} \tilde{\omega}' & \tilde{\varphi}' \\ 0 & 0 \end{pmatrix} &= \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tilde{\omega} & \tilde{\varphi} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix}^{-1} + \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} d \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} A\tilde{\omega}A^{-1} + AdA^{-1} & A\tilde{\varphi} - db - (A\tilde{\omega}A^{-1} + AdA^{-1})b \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Upon restriction on  $L(M)$ , the form  $\tilde{\omega}$  is a 1-form of linear connection on  $M_4$ ,  $\gamma^*\tilde{\omega} = \omega$ , with the correct transformation law (5.1). The 1-form  $\tilde{\varphi}$  is not a tensor form, because under translations ( $b \neq 0$ ), it transforms inhomogeneously. However, due to the reduction of  $A(M)$  to  $L(M)$ , one can define with its help a tensor form. For this purpose, we pull back the 0-form  $\varphi$ , that introduces the reduction map (5.9), to the base and compute its covariant exterior differential  $D\varphi = \tilde{\theta}$ . The result is the  $R^4$ -valued 1-form. In the matrix representation,

$$\begin{pmatrix} \tilde{\theta} \\ 0 \end{pmatrix} = D \begin{pmatrix} \varphi \\ 1 \end{pmatrix} = d \begin{pmatrix} \varphi \\ 1 \end{pmatrix} + \begin{pmatrix} \tilde{\omega} & \tilde{\varphi} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \varphi \\ 1 \end{pmatrix} = \begin{pmatrix} d\varphi + \tilde{\omega}\varphi + \tilde{\varphi} \\ 0 \end{pmatrix}.$$

Therefore, the reduction map  $\varphi$  determines a tensor 1-form for each form  $\tilde{\varphi}$

$$\tilde{\theta} = \tilde{\varphi} + d\varphi + \tilde{\omega}\tilde{\varphi}. \quad (5.11)$$

Indeed, under the change of the cross-section

$$\begin{aligned} \tilde{\varphi}' &= A\tilde{\varphi} - db - (A\tilde{\omega}A^{-1} + AdA^{-1})b, \\ \tilde{\omega}' &= A\tilde{\omega}A^{-1} + AdA^{-1}, \\ \varphi' &= -L'^{-1}z' = A\varphi + b, \end{aligned}$$

from which we find

$$\begin{aligned} \tilde{\theta}' &= \tilde{\varphi}' + d\varphi' + \tilde{\omega}'\varphi' = A\tilde{\varphi} - db - (A\tilde{\omega}A^{-1} + AdA^{-1})b \\ &\quad + d(A\varphi + b) + (A\tilde{\omega}A^{-1} + AdA^{-1})(A\varphi + b) = A\tilde{\theta}. \end{aligned} \quad (5.12)$$

The converse is also true. Indeed, given a local linear connection 1-form  $\omega$  and a tensor form  $\varphi_1$  with the transformation law  $\varphi'_1 = A\varphi_1$ ,  $A \in GL(4, R)$ , this pair uniquely determines the generalized affine connection in  $M_4$ :

$$\Omega_\sigma = \begin{pmatrix} \omega & \varphi_1 - d\varphi - \omega\varphi \\ 0 & 0 \end{pmatrix}. \quad (5.13)$$

Obtained in local coordinates, these conclusions are valid also in the bundle (without introducing the cross-sections). Namely, there is a one-to-one correspondence between the set of generalized affine connections and the set of pairs (linear connection + tensor 1-form) [3].

Recall now that another important structure –the canonical form  $\theta$ – is introduced in  $L(M)$  independently of the linear connection  $\omega$ . In accordance with (5.13), the pair  $(\omega, \theta)$  determines a special case of the generalized affine connection  $\Omega$ , which is then called the affine connection. However, the tensor form  $\tilde{\theta}$  defined by the generalized affine connection using (5.11) does not coincide with the canonical form. Moreover, the set of generalized affine connections may be divided into two essentially different classes. The first class are connections  $\Omega$  that define non-degenerate tensor 1-forms  $\tilde{\theta}$ , for which  $\tilde{\theta}(X) = 0$  means that the vector  $X$  is vertical, i.e., tangent to a fiber in  $L(M)$ . The second class encompasses the connections  $\Omega$  that define degenerate  $\tilde{\theta}$ . In the local coordinates  $\tilde{\theta} = \tilde{\theta}_\mu^\alpha E_\alpha dx^\mu$ , and the degeneracy means that  $\det \tilde{\theta}_\mu^\alpha = 0$ . One can show that  $\tilde{\theta}$  of the first class is a soldering form for the bundle of tangent affine spaces.

The affine tangent bundle  $E(M, A_4, GA(4, R), A(M))$  is associated with the principal bundle  $A(M)$  of affine frames. The typical fibre in it is the affine space  $F = R^4 = GA(4, R)/GL(4, R)$  with the dimension  $4 = \dim M$ , and this is a weakly reductive homogeneous space, since  $\mathcal{G} = ga(4, R)$ ,  $\mathcal{H} = gl(4, R)$ ,  $V = R^4$  and  $[\mathcal{H}, V] \subseteq V$ ,  $[V, V] \subseteq \mathcal{H}$ . The soldering is performed by the choice of a vector field of the “origin” of tangent affine spaces. The fact that the non-degenerate 1-form  $\tilde{\theta}$  is a soldering form, is almost trivial. Indeed, if  $X \in TL(M)$  is vertical,  $d\pi(X) = 0$ , then  $\tilde{\theta}(X) = D\varphi(X) = d\varphi(hX) \equiv 0$ ; the converse is true due to the non-degeneracy of  $\tilde{\theta}$ . The second property is also fulfilled:

$$R_h^* \begin{pmatrix} 0 & \tilde{\theta} \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & A^{-1}\tilde{\theta} \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} A^{-1} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & \tilde{\theta} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix},$$

with

$$h = \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix} \in GL(4, R).$$

The vector field  $\varphi^a(x) = -L^{-1}a_b(x)z^b(x)$  defines the origin of tangent affine spaces for the chosen cross-section  $\sigma$ :

$$M \longrightarrow A(M), \quad u = u(x).$$

### 5.3. Spontaneous symmetry breaking and non-linear realizations

Now we consider the general formalism of nonlinear group realizations as the most appropriate method of description of the spontaneous symmetry breaking.

At the core the method of nonlinear realizations is the notion of auxiliary fields which are identified with the Goldstone fields [115]. The latter appear in a physical system when the initial symmetry group  $G$  of Lagrangian of the theory is spontaneously broken to some subgroup  $H \subset G$ , with respect to which the physical vacuum is invariant. In this case, the ground state becomes degenerate, and the “manifold of vacua” is isomorphic to homogeneous factor space  $G/H$ . The parameters of  $G/H$  corresponding to the generators of the broken symmetry can be identified with the Goldstone fields which are considered as vector fields over  $M$  with the values in  $G/H$ . From the geometrical point of view, the Goldstone fields are thus the cross-sections of the vector bundle with the fiber  $G/H$  that is associated with some principal bundle  $P(M, G)$ . This refers to the local or the gauge symmetries of a physical system. According to the reduction theorem [3], the existence of a cross-section of an associated  $(G/H)$ -bundle is necessary and sufficient for the reduction of the principal  $P(M, G)$  bundle to the subbundle  $\tilde{P}(M, H)$ . Therefore, in the geometrical approach to the theory of gauge fields, one can speak of a spontaneous symmetry breaking  $G \rightarrow H$ , when there is a reduction of the corresponding principal bundles. Here we consider this geometrical picture, that should be distinguished from the spontaneous symmetry breaking in the sense of a non-invariance of the quantum vacuum state, since the “vacuum” and everything else in this theory are the purely classical (geometrical) notions.

A nonlinear realization of the group  $G$  in the theory with the broken  $G \rightarrow H$  symmetry is constructed from the linear realization, and the key link in this construction are the Goldstone or auxiliary fields. Let the matter field  $\Psi(x)$  be described as a cross-section of a vector bundle  $E(M, V^m, G, P)$  associated with the principal bundle  $P(M, G)$ , where the fiber  $V^m$  is the vector space of a representation  $\rho$  of the group  $G$  in  $GL(m, R)$ .

The field  $\Psi$  is transformed with respect to the linear (irreducible) representation  $\rho$  of the group  $G$ :

$$\Psi \longrightarrow \rho(g)\Psi, \quad g \in G. \quad (5.14)$$

The connection in the principal bundle  $P(M, G)$  introduces the parallel transport in  $E$ . Accordingly, choosing some cross-section, the covariant derivative of the field  $\Psi$  is defined, which has the following form in the local coordinates:

$$D\Psi = d\Psi + \rho'(\omega)\Psi, \quad (5.15)$$

where  $\omega$  is a  $\mathcal{G}$ -valued connection 1-form, and  $\rho'$  is an appropriate representation of the Lie algebra  $\mathcal{G}$ .

When the symmetry  $G$  is spontaneously broken to the group  $H$ , there is a reduction  $P(M, G) \rightarrow \tilde{P}(M, H)$ , and hence there exists a cross-section of an associated  $G/H$ -bundle – the Goldstone field:

$$\xi(x) \in G/H. \quad (5.16)$$

Using the auxiliary fields (introducing the “origin” in the fibers of an associated bundle), one can construct the nonlinear realization of the group  $G$ . For this purpose, we consider nonlinear fields defined by

$$\Psi_H(x) \equiv \rho(\xi^{-1}(x))\Psi(x). \quad (5.17)$$

We see that a nonlinear realization  $G$  is defined, if under the action of elements of the group  $g \in G$ , the pair  $(\xi, \Psi_H)$  is transformed according to

$$\xi \xrightarrow{g} \xi', \quad (5.18a)$$

$$\Psi_H \xrightarrow{g} \rho(h')\Psi_H, \quad (5.18b)$$

where  $\xi' \in G/H$ ,  $h' \in H$ , and  $\xi' = \xi'(g, \xi)$ ,  $h' = h'(g, \xi)$  are the nonlinear functions of their arguments that are uniquely determined from

$$g\xi = g' = \xi'h', \quad (5.19)$$

since any element of the group  $G$  is uniquely represented as a product of the element from the factor space  $G/H$  times an element of the subgroup  $H$ . It is obvious that under the action of the subgroup, the representation (5.18) is linear. Indeed, let  $g = h \in H$ , then  $\xi \xrightarrow{h} \xi' = Ad_h\xi = h\xi h^{-1}$ ,  $\Psi_H \xrightarrow{h} \rho(h)\Psi_H$ .

The notion of a nonlinear gauge field is the crucial one in the theory with the spontaneously broken gauge symmetry. The nonlinear gauge field defines the covariant derivative of  $\Psi_H$  and it is constructed on the basis of the linear representation (5.14), (5.15) with the help of the transition formula (5.17). Let  $\omega$  be the original  $\mathcal{G}$ -valued connection 1-form on  $M$ . Under the change of the cross-section, it transforms as

$$\omega \longrightarrow \omega' = g\omega g^{-1} + dg g^{-1}, \quad g(x) \in G.$$

In the method of nonlinear realizations, we define a nonlinear gauge field as

$$A = \xi^{-1}\omega\xi + \xi^{-1}d\xi, \quad (5.20)$$

where  $\xi(x)$  is an auxiliary field (5.16). Under the action of  $g$ , the connection  $A$  is transformed nonlinearly,

$$A \xrightarrow{g} A' = h'Ah'^{-1} + h'dh'^{-1}, \quad (5.21)$$

where  $h' = h'(\xi, g)$  is determined from (5.19). Therefore,

$$D^A\Psi_H = d\Psi_H + \rho'(A)\Psi_H$$

is called the covariant derivative of  $\Psi_H$  in the nonlinear realization of  $G$ .

Usually, the homogeneous space  $G/H$  is weakly reductive. The Lie algebra of the gauge group  $G$  can be decomposed into the sum  $\mathcal{G} = \mathcal{H} \oplus V$ , where  $\mathcal{H}$  is the Lie algebra of the subgroup  $H$ , [116]. Let us consider the corresponding decomposition  $A = \Gamma + B$  of the nonlinear gauge field  $A$  into the  $\mathcal{H}$ -valued 1-form  $\Gamma$  and the  $V$ -valued form of  $B$ . From the definition (5.20), the following transformation law under the action of  $g \in G$  is obvious:

$$\Gamma \longrightarrow \Gamma' = h'\Gamma h'^{-1} + h'dh'^{-1}, \quad (5.22)$$

$$B \longrightarrow B' = h'Bh'^{-1}. \quad (5.23)$$

We thus find that  $\Gamma$  transforms as the usual connection (associated with the connection in reduced principal bundle  $\bar{P}(M, H)$ ), although with the nonlinearly realized  $h' = h'(\xi, g) \in H$ . However, the  $V$ -component of the nonlinear connection  $B$  transforms homogeneously. Taking this fact into account, one can treat the method of nonlinear realizations as an alternative to the Higgs mechanism, since due to the homogeneous law (5.23), the nonlinear gauge field allows for a non-zero mass term in the Lagrangian.

In conclusion, let us consider the important example of a nonlinear realization that clearly demonstrates the above-mentioned method: the spontaneous symmetry breaking in the Higgs mechanism.

Take the Lagrangian for the multiplet of scalar (Higgs) fields  $\varphi$ ,

$$L = -\frac{1}{2}g^{\mu\nu}(D_\mu\varphi)(D_\nu\varphi) + V(\varphi), \quad (5.24)$$

where  $D_\mu\varphi = \partial_\mu\varphi + \rho'(\omega_\mu)\varphi$ . This Lagrangian is invariant with respect to the linear gauge transformations,  $\varphi \rightarrow \rho(g)\varphi$ , of the group  $G \ni g$ . When the symmetry is spontaneously broken  $G \rightarrow H$ , we proceed to the nonlinear realization according to (5.17) by defining the Goldstone fields  $\xi(x)$ . We introduce the nonlinear matter field  $\varphi_H = \rho(\xi^{-1})\varphi$ , in terms of which (5.24) is recast into

$$L = -\frac{1}{2}g^{\mu\nu}(D_\mu^A\varphi_H)(D_\nu^A\varphi_H) + V(\varphi_H). \quad (5.25)$$

In view of the spontaneous symmetry breaking, the vacuum average value  $\eta = \langle\varphi_H\rangle = \text{const} \neq 0$ , and we make the usual shift to the physical fields  $\chi = \varphi_H - \eta$ . Furthermore, we have  $D^A\varphi = D^\Gamma\chi + \rho'(B)\eta$ , since by assumption,

the “vacuum” is invariant with respect to the subgroup  $H \subset G$ , therefore its generators annihilate  $\eta$ , whereas the Higgs field itself transforms as a nonlinear multiplet  $\chi \xrightarrow{g} \rho(h')\chi$  under the action of  $G$ . Hence, the Lagrangian (5.25) manifests the Higgs phenomena in the nonlinear realization: the Goldstone degrees of freedom  $\xi$  were absorbed, so that the nonlinear field  $B_\mu$  became massive:

$$g^{\mu\nu}(D_\mu^A \varphi_H)(D_\nu^A \varphi_H) = g^{\mu\nu}(D_\mu^\Gamma \chi)(D_\nu^\Gamma \chi) + 2\rho'(B^\mu)\eta D_\mu \chi + \eta^2 B^\mu B_\mu.$$

The total Lagrangian (5.25) is still invariant with respect to the group  $G$ , which is now realized nonlinearly, and this symmetry is “hidden”. The remaining symmetry with respect to the subgroup  $H$  is explicit and realized linearly.

## 5.4. Kinematics of the gauge gravity theory

The kinematics of the gauge theory is usually understood as the identification of the basic symmetry group and the description of the corresponding dynamical variables – the gauge fields, as well as the investigation of their connection with the geometrical structure of spacetime, and the related issues.

### *Nonlinear gauge theory of Poincaré group*

The problem of a choice of the basic symmetry group in the gauge gravity theory is not purely theoretical, and it can be ultimately settled by experiment only. In a similar way, when constructing the unified theories of strong, weak and electromagnetic interactions, one can discuss only to a limited extent such questions as the number of quarks, the basic group  $SU(N)$ ,  $N = ?$ , etc; all this belongs to the field of experiment, so at best, one can specify only some *minimal model* that consistently describes the experimental data available at the current technological level.

The Poincaré group is important in the relativistic particle theory, and one can assume that this spacetime symmetry is directly relevant to the formation of the geometrical spacetime structure. Thus, the minimal model of the gauge gravity theory should be most probably based on the Poincaré group.

We construct the gauge theory of the Poincaré group  $\mathcal{P}_{10}$  similarly to the gauge theories of internal symmetries as a theory of connection in a corresponding principal bundle. However, the specific feature of the gravity theory (namely, the relation between the gauge transformations and the symmetries of spacetime) calls forth the attempts to explain the spacetime geometry directly in terms of the gauge fields. This means that, in contrast to the theories of internal symmetries, where the spacetime exists as an external background unrelated to the gauge fields, in gravity theory the geometrical spacetime structure (metric and connection, see Chapter 1) is fully determined by the corresponding gauge theory potentials. Accordingly, the principal bundle cannot be arbitrary in this case, but should be constructed as a natural spacetime object.

As the principal  $\mathcal{P}_{10}$ -bundle of the gauge theory of Poincaré group, we consider the bundle of affine orthonormal frames  $O(M)$  as a natural subbundle of the bundle of affine frames  $A(M)$ . The gauge potential is then obtained as a restriction of the generalized affine connection to  $O(M)$ .

The central issue is to explain how the gauge gravity field (namely, the  $\mathcal{P}_{10}$ -connection) determines the geometrical structure of  $M_4$ , i.e., the metric and connection on the tangent bundle  $TM$ . One conclusion looks fairly obvious: the rotational (corresponding to the Lorentz subgroup  $L_6$ ) part  $\tilde{\omega}$  of the generalized affine connection  $\Omega$ , recall (5.10), may be identified with the Lorentz connection (when the cross-section  $\sigma : M \rightarrow O(M)$  is chosen):  $\Gamma^b_a \equiv \tilde{\omega}^b_a|_{\mathcal{P}_{10}}$ .

The situation is more complex with the translational part  $\tilde{\varphi}$  (corresponding to the subgroup of translations  $T_4 = R^4$ ) of  $\Omega$  in (5.10). It is clear that the 1-form  $\tilde{\varphi}$  cannot be identified with any tensor object like the tetrad or the metric field, as one could expect, taking into account that one of the sources of the gravitational field –the energy-momentum tensor– is obviously related to translations. However, we have shown that the reduction of  $A(M)$  to the bundle of linear frames  $L(M)$ , defined by the 0-form  $\varphi$  (5.9), uniquely maps  $\tilde{\varphi}$  into the tensor 1-form of  $\tilde{\theta}$  (5.11). Accordingly, we can try to explain the spacetime metric structure in terms of the field  $\tilde{\theta}$ .

The analysis of the generalized affine connection (see Sec. 5.2.), demonstrates that the gauge  $\mathcal{P}_{10}$ -theory is essentially wider than its usual interpretation as a dynamical theory of the tetrad  $h^a_\mu$  and the Lorentz connection  $\Gamma^a_{b\mu}$  fields. Indeed, the  $\mathcal{P}_{10}$ -theory of the second class, see Sec. 5.2., does not define the metric structure on  $M_4$  at all, since in this case the form  $\tilde{\theta}$  is degenerate ( $\det \tilde{\theta}^a_\mu = 0$ ) and cannot be identified with the 1-form of coframe. More exactly, we can say that in this case the spacetime has the ultralocal degenerate Carroll geometry [117], where the metric is degenerate,  $\det g_{\mu\nu} = 0$ , everywhere in  $M_4$ . As it is shown in [118], such a state could be realized in the singularity, and within the perturbation theory developed in [118, 119] this corresponds to the zero metric signature with  $\sigma = 0$  in (2.94), and (7.51). Physically, such region is characterized by distances much smaller than the Planck length  $l_0 = 10^{-35}$  m.

In contrast to this unusual situation, the gauge  $\mathcal{P}_{10}$ -theory of the first class introduces the metric on  $TM$  in a consistent way. However, in this case the resulting structure is much wider than the Riemann-Cartan structure. Indeed, we have shown in Sec. 5.2. that the 1-form  $\tilde{\theta}$  is the generalized soldering form of the affine tangent bundle, but it *does not coincide* with the canonical form  $\theta$  (the coframe 1-form  $\theta = h$ ), as it was noticed in [103]. The difference of  $\tilde{\theta}$  from  $\theta$  is in arbitrary general linear transformation  $\tilde{\theta}^a = L^a_b \theta^b$ ,  $L \in GL(4, R)$ .

With an account of the explicit covariance with respect to the local Lorentz transformations (we consider the principal bundle of orthonormal frames), it reduces to the difference on the Goldstone transformation from the factor space  $GL(4, R)/SO(3, 1)$ . One can use this arbitrariness in the definition of tetrads to construct the conformal gravity theory of new type in the Riemann-Cartan spacetime, as we discussed in Sec. 4.1., see [69].

In the rest of the section, we present the rigorous foundation for these conclusions in the framework of the nonlinear realization of the gauge  $\mathcal{P}_{10}$ -theory.

Construction of the nonlinear gauge theory of the Poincaré group consists (schematically) of the two stages. The first one is formulated [99]-[105] as the soldering procedure, when the bundle of affine frames is reduced to the bundle of linear frames. In physical terms, this means the spontaneous breaking of the translational symmetry, which reflects the fact that the physical matter fields depend only on the points of the base  $M_4$ , but not on the points of the fiber [99, 100]. The (implicit) second stage is very important: the principal  $\mathcal{P}_{10}$ -bundle of the gauge gravity theory is itself a subbundle in  $A(M)$ , in other words, it is obtained as the corresponding restriction or reduction of the structural group  $G(4, R) \rightarrow \mathcal{P}_{10}$ . In accordance with this, we notice that an arbitrary element  $g \in GA(4, R)$  is uniquely represented in the  $5 \times 5$  matrix disguise as

$$g = \begin{pmatrix} A & b \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{1} & \xi \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix} h, \quad \text{where} \quad h = \begin{pmatrix} L & 0 \\ 0 & 1 \end{pmatrix}.$$

Here, the first factor is the Goldstone translational field – the element from  $GA(4, R)/GL(4, R)$ , and the second factor  $u \in GL(4, R)/SO(3, 1)$  is the Goldstone field responsible for the reduction of  $A(M)$  to  $AO(M)$ ; finally, the matrix  $h \in SO(3, 1)$  represents the exact (unbroken) Lorentz symmetry.

Now we proceed from the linear gauge field  $\Omega$ , the generalized affine connection given in (5.10), to the nonlinear gauge field via (5.20) with the help of the auxiliary fields

$$\begin{pmatrix} 1 & \xi \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} u & \xi \\ 0 & 1 \end{pmatrix}.$$

A straightforward computation yields

$$\begin{aligned} A &= \begin{pmatrix} u & \xi \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} \tilde{\omega} & \tilde{\varphi} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u & \xi \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} u & \xi \\ 0 & 1 \end{pmatrix}^{-1} d \begin{pmatrix} u & \xi \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} u^{-1}\tilde{\omega}u + u^{-1}du & u^{-1}(\tilde{\varphi} + d\xi + \tilde{\omega}\xi) \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

As a result, we finally find that the rotational part of the gauge gravitational field  $\Gamma := u^{-1}\omega u + u^{-1}du$  is realized linearly with respect to the Lorentz group  $L_6$  and defines the local Lorentz connection on  $M_4$ , whereas the tetrad field (the canonical form or the 1-form of coframe)  $\theta := u^{-1}\tilde{\theta} = u^{-1}(\tilde{\varphi} + d\xi + \tilde{\omega}\xi)$  is interpreted as the nonlinear gauge field with an additional contribution of the Goldstone fields  $u \in GL(4, R)/SO(3, 1)$ , transforming homogeneously under the action of translations.

In this way, the Riemann-Cartan (see Chapter 1) geometry  $U_4$  naturally arises on the spacetime manifold  $M_4$ , where the metric (tetrads) and the connection

are expressed in terms of the basic dynamical variables of the gauge  $\mathcal{P}_{10}$ -theory of gravity – the generalized affine connection  $\Omega$ , or the equivalent set of the nonlinear gauge fields  $\Gamma$  and  $\theta$  and the Goldstone variables  $u$ ,  $\xi$ .

### *Nonlinear gauge theory of de Sitter group*

The Poincaré group  $\mathcal{P}_{10}$  is not semi-simple. In Chapter 6, we will demonstrate that this fact has the far-reaching consequences when constructing the dynamics of the gauge gravity theory, and actually yields unsatisfactory results. In an effort to improve the situation, we now consider the gauge gravity theory for the nearest semi-simple extension of  $\mathcal{P}_{10}$  – the de Sitter group  $S_{10}$ , which reduces to the Poincaré group via the Wigner-Inonu “contraction”.

Technically, it will be more convenient to work in a vector bundle, and not in the principal one, although one should remember that the gauge field is originally defined as a connection in the principal bundle, and then it uniquely introduces the connection in the associated bundle, which for this reason is also called the gauge field.

De Sitter group  $S_{10} = SO(1, 4)$  is defined as a group of motion of the de Sitter space  $\Sigma^4$  which is a 4-dimensional pseudo-Riemannian space of constant negative curvature. The latter is usually represented as a hypersurface in  $R^{1,4}$ , the 5-dimensional pseudo-Euclidean space with the metric  $\eta_{AB} = \text{diag}(-1, 1, 1, 1, 1)$  (the indices  $A, B, C, D \dots = 0, 1, 2, 3, 4$  label 5-dimensional components). The de Sitter space is defined by the equation

$$\eta_{AB}x^A x^B = \ell^2, \quad (5.26)$$

where  $\{x^A\}$  are the global Cartesian coordinates in  $R^{1,4}$ , and  $\ell > 0$  is the curvature radius of the de Sitter space. The equivalent form of (5.26) reads

$$\eta_{ab}x^a x^b + (x^4)^2 = \ell^2,$$

in terms of the 4-dimensional Minkowski metric  $\eta_{ab} = \text{diag}(-1, 1, 1, 1)$ ; the lower case indices  $a, b, c, \dots = 0, 1, 2, 3$ .

The group  $SO(1, 4)$  is ten-parametric, and its generators  $M_{AB} = -M_{BA}$  satisfy commutation relations

$$[M_{AB}, M_{CD}] = \eta_{AD}M_{BC} - \eta_{BD}M_{AC} - \eta_{AC}M_{BD} + \eta_{BC}M_{AD}. \quad (5.27)$$

It is convenient to decompose the generators  $M_{AB}$  into two groups:  $J_{ab} = M_{ab}$  and  $P_a = \ell^{-1}M_{4a}$ . Then we rewrite (5.27) as

$$\left. \begin{aligned} [J_{ab}, J_{cd}] &= \eta_{ad}J_{bc} - \eta_{bd}J_{ac} - \eta_{ac}J_{bd} + \eta_{bc}J_{ad}, \\ [P_a, P_b] &= -\frac{1}{\ell^2}J_{ab}, \quad [P_a, J_{bc}] = \eta_{ab}P_c - \eta_{ac}P_b, \end{aligned} \right\} \quad (5.28)$$

which shows that the Lie algebra  $\mathcal{G} = so(1, 4)$  is decomposed into the sum  $\mathcal{G} = \mathcal{H} \oplus V$ , where  $\mathcal{H}$  is the subalgebra formed by six generators  $J_{ab} = -J_{ba}$ ,

isomorphic to the algebra  $so(1, 3)$  of the Lorentz group;  $V$  is the 4-dimensional vector space spanned by  $P_a$ .

The de Sitter space  $\Sigma^4$  is homogeneous, since  $SO(1, 4)$  acts transitively on  $\Sigma^4$ . Denoting  $\mathbf{0} = \{x^a = 0, a = 0, 1, 2, 3\}$ , we choose the point  $\begin{pmatrix} \mathbf{0} \\ l \end{pmatrix} \in \Sigma^4$  as the center, its stabilizer is the Lorentz group  $SO(1, 3) = L_6$  formed by the  $5 \times 5$  matrices  $S^A{}_B = \begin{pmatrix} L^a{}_b & 0 \\ 0 & 1 \end{pmatrix}$ , with  $L^a{}_b \in SO(1, 3)$ . Accordingly,  $\Sigma^4 = S_{10}/L_6$ .

From (5.28) we see that this is the weakly reductive space, since  $[\mathcal{H}, V] \subset V$ .

Let  $E(M, R^{1,4}, S_{10}, P)$  be a vector bundle associated with the principal bundle  $P(M, S_{10})$ . The action of the structural group  $S_{10}$  in  $E$  we realize by the  $5 \times 5$  matrices with the following parametrization distinguishing the subgroup  $L_6$  (we omit the indices to make the formulas more compact):

$$S = S_T S_L, \quad S \in SO(1, 4). \quad (5.29)$$

Here  $S_L = \begin{pmatrix} L & 0 \\ 0 & 1 \end{pmatrix}$ ,  $L \in SO(1, 3)$  describes the purely Lorentz rotations, and

$S_T$  is the matrix of the de Sitter boost mapping the vector  $\begin{pmatrix} \mathbf{0} \\ l \end{pmatrix} \in \Sigma^4$  into

an arbitrary point of the de Sitter space  $V^A = \begin{pmatrix} V^a \\ V^4 \end{pmatrix} \in \Sigma^4$ , with  $V^A = \ell t^A$ ,

$\eta_{AB} V^A V^B = \ell^2$ . Explicitly:

$$S_T = \begin{pmatrix} \delta_b^a - t^a t_b (1 + t^4)^{-1} & t^a \\ -t_b & t^4 \end{pmatrix}. \quad (5.30)$$

The finite rotation  $S_T$  can be obtained from the infinitesimal one by the exponential map  $(S_T)^A{}_B = \exp \begin{pmatrix} 0 & \chi^a \\ -\chi_b & 0 \end{pmatrix}$ , where  $\chi^a$  (with  $\chi_b = \eta_{ab} \chi^a$ ) are the four transformation parameters which determine  $t^A$  in (5.30) by the formulas

$$t^a = (\sin \chi) \chi^{-1} \chi^a, \quad t^4 = \cos \chi, \quad \chi^2 = \eta_{ab} \chi^a \chi^b.$$

The linear gauge field for the de Sitter group is identified with the connection in  $E$ . As an element of the Lie algebra of the de Sitter group  $SO(1, 4)$ , the connection 1-form has the following structure:

$$\Omega^A{}_B = \begin{pmatrix} \tilde{\omega}^a{}_b & \tilde{\theta}^a \\ -\tilde{\theta}_b & 0 \end{pmatrix}, \quad \tilde{\omega}_{ab} = -\tilde{\omega}_{ba}, \quad \tilde{\theta}_b = \eta_{ab} \tilde{\theta}^a. \quad (5.31)$$

However, this decomposition is invariant only with respect to the Lorentz group, whereas under the action of the full  $S_{10}$  transformations (5.29), the fields  $\tilde{\omega}$  and  $\tilde{\theta}$  are mixed. Therefore, one cannot identify  $\tilde{\omega}$  with the connection in the tangent bundle, and  $\tilde{\theta}$  with the fundamental form. To obtain the geometrical structure on the base  $M_4$ , it is necessary to take into account that the  $S_{10}$ -theory is not an “internal” one, and to perform a soldering of the bundle to the spacetime manifold. For this purpose, as we observed from the example of the Poincaré group above, we have to choose the cross-section of the  $(G/H)$ -bundle, i.e., the Goldstone field, and proceed to the nonlinear realization of the group  $G$ . In this case,  $G = SO(1,4)$ ,  $H = SO(1,3)$ ,  $G/H = \Sigma^4$  and the Goldstone field is actually the 5-vector  $t^A = \begin{pmatrix} t^a \\ t^4 \end{pmatrix}$ , instead of which however we will (more rigorously) consider the de Sitter boost (5.30).

Following the general framework of Sec. 5.3., we construct the nonlinear gauge field using  $\xi = S_T$ :

$$A = \xi^{-1}\Omega\xi + \xi^{-1}d\xi = \begin{pmatrix} \Gamma^a_b & \theta^a \\ -\theta_b & 0 \end{pmatrix}, \quad (5.32)$$

where  $\theta_b = \eta_{ab}\theta^a$ , and we find explicitly

$$\Gamma^a_b = \tilde{\omega}^a_b + \frac{t^a Dt_b - t_b Dt^a}{1 + t^4}, \quad (5.33)$$

$$\theta^a = t^4 \tilde{\theta}^a + Dt^a - \frac{t^a(dt^4 - \tilde{\theta}_b t^b)}{1 + t^4}. \quad (5.34)$$

Here we denoted  $Dt^a = dt^a + \tilde{\omega}^a_b t^b$ .

Under the action of the de Sitter group  $S_{10}$ , which is now realized nonlinearly, the gauge fields are transformed as

$$\Gamma'^a_b = L'^a_c \Gamma^c_d L'^{-1d}_b + L'^a_c dL'^{-1c}_b, \quad \theta'^a = L'^a_b \theta^b,$$

where  $S'_L = \begin{pmatrix} L'^a_b & 0 \\ 0 & 1 \end{pmatrix}$ , and the nonlinear function  $L' = L'(t^A, S)$  is determined from  $S\xi = SS_T = S'_T S'_L$ .

Thus, we can finally identify the 1-form  $\Gamma^a_b$  with the local Lorentz connection, and the 1-form  $\theta^a$  with the canonical form. As a result, the Riemann-Cartan geometry naturally arises on  $M_4$ . Note that, in general, the field  $\theta^a$  depends on the additional Goldstone variables, corresponding to the embedding of the de Sitter group  $S_{10}$  into the general linear group  $GL(5, R)$ , similarly to the  $\mathcal{P}_{10}$ -theory. However, we will not distinguish them explicitly, since we are more interested in the specific features of the  $S_{10}$ -theory, which are important for the construction of the consistent dynamical scheme.

*Conclusions: the general scheme of the gauge gravity theory*

Let us summarize the essential points of the gauge approach to the gravity theory, distinguishing it from the theory of internal symmetries.

1. The spacetime  $M_4$  is not considered as a fixed background. The determination of its geometrical structure is a main issue of the gauge-theoretic scheme.

2. The gauge field (potential) in the gravity theory is the connection in the principal bundle  $P(M, G)$ .

3. The principal bundle of the gauge gravity theory is not abstract, but is realized as a certain structure on  $M_4$ . For the  $\mathcal{P}_{10}$ -theory, this is the bundle of affine frames  $A(M)$ .

4. The specific realization of bundles in the gauge gravity theory implies their soldering to the base. There exists a cross-section of the  $(G/H)$ -bundle and the soldering 1-form is defined by the translational part of the connection in  $P(M, G)$ .

5. The soldering establishes a homomorphism of the  $P(M, G)$  bundle structure onto the tangent bundle  $TM$ , thereby introducing the geometrical structure on the spacetime  $M_4$ , with the metric and connection defined in terms of the gauge gravity field.

6. The explicit construction of the geometrical spacetime structure is performed in terms of the connection in  $P(M, G)$  in the nonlinear realization of the gauge group. The Goldstone fields, arising in the reduction  $G \rightarrow H$  (the latter being closely related to the soldering), also contribute to the metric and the linear connection in  $TM$ .

It is worthwhile to note that the group of general coordinate transformations is not related directly to the construction of the gauge gravity theory. However, the gauge transformations of special form may induce the infinitesimal transformations of coordinates on  $M_4$ , if we introduce the latter with the help of the reduction map  $\varphi$  defined in (5.9). Indeed, using the cross-section  $\sigma : M \rightarrow A(M)$ , we can define the local coordinates  $x^\mu$  as  $x = \varphi \circ \sigma$ . Then

under the action of translations  $T = \begin{pmatrix} \mathbf{1} & -a \\ 0 & 1 \end{pmatrix}$ , the cross-section  $\sigma \rightarrow R_T^* \sigma$ ,

and we obtain  $x^\mu \rightarrow x^\mu + a^\mu(x)$  as the local general coordinate transformations.

Within the framework described above, the Riemann structure of GR cannot be obtained without additional restrictions of the natural geometrical structure arising on the spacetime. Such restrictions (for example, a priori putting the torsion equal zero) contradict the consistent gauge-theoretic scheme, and they do not follow from any (even non-gauge-theoretic) fundamental physical principles. In this relation, it is instructive to recall Einstein's words, that "...the question whether this continuum has a Euclidean, Riemannian, or any other structure is a question of physics proper which must be answered by experience, and not a question of a convention to be chosen on grounds of mere expediency." [39].

# 6

## Gauge gravity models with dynamical $(\Gamma$ - $S$ )-interaction

### 6.1. Principles for construction of dynamics

In the previous Chapter, we saw that the gauge approach based on the Poincaré group  $\mathcal{P}_{10} = T_4 \rtimes L_6$  in general yields a non-Einstein theory of gravity. One can choose the tetrad field  $h_\mu^a$  and the local Lorentz connection  $\Gamma^a_{b\mu}$  field on  $M_4$  as the main dynamical variables of the  $\mathcal{P}_{10}$ -theory. In accordance with the general gauge scheme, their source is the  $\mathcal{P}_{10}$ -current consisting of the canonical energy-momentum tensor  $t^\mu_a$  and the canonical spin tensor  $S^{\mu a}_b$ , which correspond to subgroups  $T_4$  and  $L_6$ , respectively. As compared to GR, the new feature of the  $\mathcal{P}_{10}$ -theory is the existence of an additional  $(\Gamma^a_{b\mu}$ - $S^{\mu b}_a)$ -interaction, along with Einstein's  $(h_\mu^a$ - $t^\mu_a)$ -interaction. Since the Poincaré group is not simple, these interactions are characterized by the two different coupling constants – one with the dimension  $[l_0^2] = \text{m}^2$  for translations and the dimensionless one  $\lambda$  for Lorentz rotations.

The prediction of a new type of the gravitational interaction (spin-connection) is the main physical consequence of the gauge gravity theory for the Poincaré group.

The realization of this interaction in the Einstein-Cartan model was reviewed in Chapters 3 and 4. However, one cannot consider the ECT as a satisfactory gauge gravity theory. The Lagrangian  $\mathcal{L}_{ECT} \sim R(\Gamma) \sim (\partial\Gamma + \Gamma\Gamma)hh$  is non-dynamical both for  $\Gamma$  and for  $h$ , in the sense that the field equations do not contain derivatives of the gauge field strengths, unlike the Maxwell or Yang-Mills theories. Due to the contact nature of the  $(\Gamma$ - $S)$ -interaction, one can eliminate the torsion from the theory, thereby arriving at the effective non-renormalizable

$S^2$ -self-interaction of matter. The degeneracy of ECT is also manifest in the absence of the second coupling constant for the  $(\Gamma-S)$ -interaction, as result of which the spin effects turn out to be negligibly small. Another sign of degeneracy is the mismatch of the gauge kinematics and the dynamics of the model: we have  $R \sim t$ ,  $Q \sim S$  in ECT field equations, linking the  $L_6$  field strength (the curvature) to the  $T_4$  current (the energy-momentum), and the  $T_4$  field strength (the torsion) to the  $L_6$  current (the spin).

The consistent nondegenerate dynamical scheme of the gauge gravity theory should be free of such deficiencies. However, its construction is complicated due to the essential non-uniqueness of theory's action, which is a consequence of the nonlinear realization of the gauge symmetry that allows for non-polynomial (in the nonlinear fields  $h_\mu^a$  and  $g_{\mu\nu}$ ) interactions. We can try to overcome this difficulty by supplementing the gauge theory with a number of conditions (natural from physical and geometrical viewpoints) that help to select the basic Lagrangian of the theory.

The appropriate general requirements can be formulated as follows:

1. Correspondence principle: in the absence of spin and torsion, the field equations should allow for a limiting transition to GR.
2. Both gravitational interactions,  $(h-t)$  and  $(\Gamma-S)$ , should be dynamical.
3. The dynamics of the theory should be consistent with the gauge kinematics in the sense that the source of the gauge field of a certain spacetime symmetry is the corresponding Noether current.

Let us apply these criteria to analyze the possible choices of the Lagrangians (more exactly, of the action functionals) of the gravitational theory considered earlier in the literature. To keep the correct dimension of the action,  $[S] = [\hbar]$ , we write down all the dimensional factors explicitly.

$$\text{a) } S_{GR} = \frac{1}{2\kappa c} \int d^4x \sqrt{g} R, \quad S_{ECT} = \frac{1}{2\kappa c} \int d^4x \sqrt{g} R(\Gamma).$$

These models with the Lagrangians linear in the curvature do not satisfy the requirements 2 and 3.

$$\text{b) } S_{YM} = -\frac{\hbar}{4} \int d^4x \sqrt{g} R^{ab}{}_{\mu\nu}(\Gamma) R_{ab}{}^{\mu\nu}(\Gamma).$$

The Yang-Mills Lagrangian obviously provides the fulfilment of the requirements 2 and 3. However, this model has two serious problems: the absence of GR limit and the conformal invariance of  $S_{YM}$  which imposes a strong constraint on matter (the vanishing of the trace of the energy-momentum tensor). Therefore, this model can be viewed only as a microscopic limit of a fundamental theory.

$$\text{c) } S_0 = S_{GR} + \beta S_{YM}, \quad \beta = \text{const.}$$

This choice does not completely satisfy the requirement 3, since this model arises in the gauge theory of the Lorentz group with the assumption that the metric is a priori defined on the base.

$$\text{d) } S_1 = S_{ECT} + \beta S_{YM}, \quad \beta = \text{const.}$$

Also in this model the requirement 3 is not fulfilled in the sense that the variation with respect to the tetrads yields non-dynamical equations.

$$\text{e) } S_2 = -\frac{\alpha}{2\kappa c} \int d^4x \sqrt{g} Q^a{}_{\mu\nu} Q_a{}^{\mu\nu} + \beta S_{YM}, \quad \alpha, \beta = \text{const.}$$

This model does not satisfy the principle of correspondence with GR. However, one can come up with a modification

$$S_3 = \frac{1}{2\kappa c} \int d^4x \sqrt{g} (-Q^a{}_{\mu\nu} Q_a{}^{\mu\nu} + 2Q_\mu Q^\mu) + \beta S_{YM},$$

that contains the Schwarzschild solution as a torsionless limiting case.

f) The model, satisfying all the requirements above, is given by the action

$$S_{\mathcal{P}} = \int d^4x \sqrt{g} \left\{ \frac{1}{2\kappa c} (R(\Gamma) - 2\alpha Q_a{}^{\mu\nu} Q^a{}_{\mu\nu}) - \frac{1}{4\lambda} R^{ab}{}_{\mu\nu}(\Gamma) R_{ab}{}^{\mu\nu}(\Gamma) \right\}. \quad (6.1)$$

Here  $\alpha$  is a dimensionless coupling constant, and the dimension of  $[\lambda] = [1/\hbar]$ .

Below we demonstrate, how this dynamical framework naturally arises in  $\mathcal{P}_{10}$ - and in  $S_{10}$ -theory, so that one can avoid a nontrivial analysis of all possible Lagrangians quadratic in the curvature and torsion with arbitrary coefficients.

## 6.2. Choice of the action in the gauge gravity theory

### *Difficulties of the Poincaré gauge theory*

Let  $\Omega$  be the 1-form of the generalized affine connection on  $M_4$  (we assume a cross-section  $\sigma: M \rightarrow O(M)$ ), interpreted as the linear  $\mathcal{P}_{10}$ -gauge gravitational field. The field strength is identified with the curvature 2-form

$$\mathcal{R} = d\Omega + \Omega \wedge \Omega = \begin{pmatrix} \bar{R} & \bar{\Theta} \\ 0 & 0 \end{pmatrix},$$

where  $\bar{R} = \widetilde{d\omega} + \widetilde{\omega} \wedge \widetilde{\omega}$ ,  $\bar{\Theta} = \widetilde{d\theta} + \widetilde{\omega} \wedge \widetilde{\theta}$ . However, the latter is not invariant with respect to the decomposition of the curvature into the translational and the Lorentz parts, and to improve this we proceed to the nonlinear realization. The nonlinear gauge field

$$A = \begin{pmatrix} \Gamma^a{}_b & \theta^a \\ 0 & 0 \end{pmatrix} \quad (6.2)$$

encompasses the local Lorentz connection  $\Gamma^a{}_b$  (the  $L_6$  gauge field) and the tetrad field  $h^a$  (the  $T_4$  nonlinear gauge field) is introduced by  $\theta^a = \frac{1}{l_0} h^a$ , where the factor  $l_0$  is included to get the correct dimension,  $[h^a] = 1$ . The curvature 2-form

$$\mathcal{R} = \begin{pmatrix} R^a{}_b & \frac{1}{l_0} \widehat{Q}^a \\ 0 & 0 \end{pmatrix}, \quad \begin{aligned} R^a{}_b &= d\Gamma^a{}_b + \Gamma^a{}_c \wedge \Gamma^c{}_b, \\ \widehat{Q}^a &= dh^a + \Gamma^a{}_b \wedge h^b \end{aligned} \quad (6.3)$$

consists of the Lorentz curvature 2-form  $R^a{}_b$  and the torsion 2-form  $\widehat{Q}^a$ , which are the  $L_6$  and the  $T_4$  field strengths of the  $\mathcal{P}_{10}$  gauge field, respectively. Note that  $\widehat{Q}^a = \frac{1}{2} \widehat{Q}^a{}_{\mu\nu} dx^\mu \wedge dx^\nu$  with  $\widehat{Q}^a{}_{\mu\nu} = -2Q^a{}_{\mu\nu}$ , cf. (1.3).

One can construct the dynamics of the  $\mathcal{P}_{10}$ -theory by two methods:

1. We can use the irreducible parts  $\Gamma$  and  $h$  of the nonlinear gauge field (6.2) to derive the covariant objects (6.3) and then to build the Lagrangian as an arbitrary polynomial of invariants formed from the curvature, torsion and tetrad,  $R, Q, h$ . This procedure is essentially non-unique.

2. Using the analogy with the Yang-Mills fields of the internal symmetries, one can try to construct the Lagrangian depending on  $A$  (and  $\mathcal{R}$ ) as a whole. In this case, the simplest choice is the Yang-Mills Lagrangian  $\frac{1}{2}\text{Tr}(\mathcal{R} \wedge *\mathcal{R})$ . The trace operation “Tr” refers to the  $\mathcal{P}_{10}$ -“internal” indices, and the Hodge operator (dualization,  $*$ ) is defined with the help of the metric  $g_{\mu\nu} = h_\mu^a h_\nu^b \eta_{ab}$ , resulting from the nonlinear realization of the  $\mathcal{P}_{10}$  group. Since the Poincaré group is semi-simple, its Cartan-Killing metric (encoded in the trace operation Tr) is degenerate, and as a result, the variational derivation of the field equation does not commute with the decomposition of  $A$  and  $\mathcal{R}$  into the  $L_6$ - and  $T_4$ -irreducible parts. Indeed, if we make such decomposition at first, then

$$\frac{1}{2}\text{Tr}(\mathcal{R} \wedge *\mathcal{R}) = \frac{1}{2}\text{Tr} \left\{ \begin{pmatrix} R & \frac{1}{l_0}\widehat{Q} \\ 0 & 0 \end{pmatrix} \wedge \begin{pmatrix} *R & \frac{1}{l_0}*\widehat{Q} \\ 0 & 0 \end{pmatrix} \right\} = \frac{1}{2}\text{Tr}(R \wedge *R),$$

hence the torsion completely drops out, and the remaining Lagrangian is described as the case b) in Sec. 6.1.. The resulting field equations are unsatisfactory for the macroscopic gravitation theory [100].

On the other hand, if the action  $\frac{1}{2\lambda} \int \text{Tr}(\mathcal{R} \wedge *\mathcal{R}) + S_{\text{mat}}$  is formally varied with respect to  $A$ , we find the field equation

$$\mathcal{D} * \mathcal{R} \equiv d * \mathcal{R} + A \wedge * \mathcal{R} - * \mathcal{R} \wedge A = \lambda * J. \quad (6.4)$$

Here the  $\mathcal{P}_{10}$ -current is obtained as a variational derivative  $*J := \delta S_{\text{mat}}/\delta A$  of the matter action with respect to  $A$ . The structure of the source reflects its relation to the Poincaré group,

$$J = \begin{pmatrix} S^a{}_b & l_0 t^a \\ 0 & 0 \end{pmatrix}, \quad (6.5)$$

where the 1-forms  $S^a{}_b$  and  $t^a$  are the  $L_6$ - and  $T_4$ -irreducible parts of  $J$ , respectively. With an account of (6.2) and (6.3), we recast (6.4) into the  $L_6$ - and  $T_4$ -components:

$$\mathcal{D} * R^a{}_b \equiv d * R^a{}_b + \Gamma^a{}_c \wedge * R^c{}_b - * R^a{}_c \wedge \Gamma^c{}_b = \lambda * S^a{}_b, \quad (6.6)$$

$$\mathcal{D} * \widehat{Q}^a - * R^a{}_b \wedge h^b = \lambda l_0^2 * t^a. \quad (6.7)$$

Let us investigate the vacuum solutions ( $S^a{}_b = 0$ ,  $t^a = 0$ ). We switch to the coordinate notation in components and use the decomposition (1.12) to write

the field equations (6.6) and (6.7) in terms of the metric and contortion tensors  $(g_{\mu\nu}, T^\lambda{}_{\mu\nu})$  instead of the original variables  $(h^a{}_\mu, \Gamma^a{}_{b\mu})$ :

$$R^\alpha{}_\beta{}^{\mu\nu}(\Gamma)_{;\nu} + T^\alpha{}_{\sigma\nu}R^\sigma{}_\beta{}^{\mu\nu}(\Gamma) - T^\sigma{}_{\beta\nu}R^\alpha{}_\sigma{}^{\mu\nu}(\Gamma) = 0, \quad (6.8)$$

$$T^{\alpha[\nu\beta]}{}_{;\nu} + T^\alpha{}_{\sigma\nu}T^{\sigma[\nu\beta]} + \frac{1}{2}R^{\alpha\beta}(\Gamma) = 0. \quad (6.9)$$

Recall that the semicolon  ${}_{;\nu}$  denotes the Riemannian covariant derivatives defined by the Christoffel symbols.

The field equations (6.6) and (6.7) is the system of differential second order equations for the tetrad and the local Lorentz connection. Therefore, both  $(h-t)$  and  $(\Gamma-S)$ -interactions are dynamical in this model. Decomposing the curvature tensor into the Riemannian (depending only on the metric) part and the post-Riemannian (torsion-dependent) part, we find the system (6.8) and (6.9) of differential equations of the second order for the metric and contortion  $(g_{\mu\nu}, T^\lambda{}_{\mu\nu})$ .

When the torsion equal to zero,  $T^\alpha{}_{\mu\nu} = 0$ , the vacuum system (6.8) and (6.9) reduces to the Einstein equation  $R_{\alpha\beta} = 0$ . But in general case, the torsion is propagating, and thus the  $(\Gamma-S)$ -interaction has a non-zero effective radius, in contrast to ECT.

The physical interpretation of the sources in the right-hand sides of the field equations (6.6) and (6.7) is a difficult issue. This primarily refers to (6.7), since the variational derivation of the equation (6.6), corresponding to  $L_6$ -subgroup, does not depend on the order of separation of translations in  $\mathcal{P}_{10}$ . Accordingly, the source  $S^a{}_{b\mu}$  can be consistently identified with the canonical spin density tensor of matter [99, 100]. However,  $t^\alpha{}_\mu$  in (6.5) cannot be identified with the energy-momentum tensor, since when deriving the system (6.4), the dependence of the metric  $g_{\mu\nu}$  (in the Hodge operator  $*$ ) on the translational field  $\theta$  was not taken into account (recall that the formal variation of the gravitational and matter actions was performed only with respect to the gauge field  $A$  as a whole).

Therefore, although the second method of construction of the  $\mathcal{P}_{10}$ -theory dynamics allows to highlight some important features typical for the gauge gravity theory (such as the correspondence with GR, the dynamical nature of all interactions, and the partial consistency of the dynamics with the kinematics), it does not lead to the physically satisfactory results. This is related to the fact that the Poincaré group  $\mathcal{P}_{10}$  is not non-semisimple.

Thus, we have to turn to the first method to construct the dynamics of the  $\mathcal{P}_{10}$ -theory. However, in this case, there are too many arbitrary coupling constants in the theory. For their concretization and for the fixation of the structure of the gravitational Lagrangian, we now analyse the construction of the gauge theory dynamics for de Sitter group  $S_{10}$  – the nearest semisimple extension of  $\mathcal{P}_{10}$ .

### *Dynamics of $S_{10}$ -theory of gravity*

In the discussion of kinematics of the gauge gravity theory in Sec. 5.4., we identified the gauge potential for de Sitter group theory with the nonlinear connection (5.32):

$$A = \begin{pmatrix} \Gamma^a_b & \frac{1}{l_0} h^a \\ -\frac{1}{l_0} h_b & 0 \end{pmatrix}.$$

The corresponding gauge field strength is identified with the curvature 2-form

$$\mathcal{R} = dA + A \wedge A = \begin{pmatrix} R^a_b - \frac{1}{l_0^2} \pi^a_b & \frac{1}{l_0} \widehat{Q}^a \\ -\frac{1}{l_0} \widehat{Q}_b & 0 \end{pmatrix}, \quad (6.10)$$

where  $\pi^a_b = h^a \wedge h_b = \frac{1}{2} \pi^a_{b\mu\nu} dx^\mu \wedge dx^\nu$ . Obviously,  $\pi^a_{b\mu\nu} = h^a_\mu h_{\nu b} - h^a_\nu h_{\mu b}$ .

Since the de Sitter group is semisimple, its Cartan-Killing form is non-degenerate, and one can choose the action of the theory in the standard Yang-Mills form

$$S_{dS} = \frac{1}{4\lambda} \int d^4x \sqrt{g} \text{Tr}(\mathcal{R} \wedge * \mathcal{R}).$$

With an account of (6.10), we have

$$S_{dS} = \int d^4x \sqrt{g} \left\{ \frac{1}{2\kappa c} [R(\Gamma) - 2\Lambda - 2Q^a_{\mu\nu} Q_a^{\mu\nu}] - \frac{1}{8\lambda} R^{ab}{}_{\mu\nu} R_{ab}{}^{\mu\nu} \right\}, \quad (6.11)$$

where we redefined the coupling constants as  $\kappa c = l_0^2 \lambda$ ,  $\Lambda = \frac{3}{l_0^2}$ .

The action (6.11) is invariant with respect to the nonlinear realization of the de Sitter group  $S_{10}$ . Its structure coincides, up to a  $\lambda$ -term, with that of  $S_{\mathcal{P}}$  (6.1) with  $\alpha=1$ . Therefore, the non-degeneracy of the Cartan-Killing form for  $S_{10}$  makes it possible to construct the gauge theory dynamics that satisfies the requirements 1-3 formulated in Sec. 6.1..

From the action (6.11), the equations of the gravitational field are derived with the help of variation with respect to the local Lorentz connection  $\Gamma^a_{b\mu}$  and the tetrad  $h^a_\mu$ . In vacuum (in the absence of matter) they read

$$-\frac{1}{2\lambda} \overline{\mathcal{D}}_\nu R_{ab}{}^{\mu\nu}(\Gamma) - \frac{1}{\kappa c} (Q^\mu{}_{ab} + h^a_\mu Q_b - h^a_\mu Q_a + 2Q_{[ab]}{}^\mu) = 0, \quad (6.12)$$

$$\frac{1}{\kappa c} (2\overline{\mathcal{D}}_\nu Q_a{}^{\mu\nu} - R^\mu{}_a(\Gamma) + \frac{1}{2} h^a_\mu R(\Gamma) - \Lambda h^a_\mu) + \frac{1}{c} \tau^\mu{}_a = 0, \quad (6.13)$$

where modified covariant derivative  $\overline{\mathcal{D}}_\mu$  is defined using  $\Gamma^a_{b\mu}$  for the Lorentz (tetrad) indices and using  $\{\lambda_{\mu\nu}\}$  for the world (coordinate) indices ( $\mu, \nu, \dots$ ).

The tensor  $\tau^\mu{}_a$  in (6.13) reads

$$\begin{aligned} \tau^\mu{}_a &= \frac{c}{2\lambda} \left( R^{bc}{}_{a\nu}(\Gamma) R_{bc}{}^{\mu\nu}(\Gamma) - \frac{1}{4} h^a_\mu R^{bc}{}_{\alpha\beta}(\Gamma) R_{bc}{}^{\alpha\beta}(\Gamma) \right) \\ &+ \frac{4}{\kappa} \left( Q_{ba\nu} Q^{b\mu\nu} - \frac{1}{4} h^a_\mu Q^b{}_{\alpha\beta} Q_b{}^{\alpha\beta} \right). \end{aligned} \quad (6.14)$$

This is the energy-momentum tensor of  $S_{10}$ -gauge field, that takes into account the influence on the spacetime geometry of the gravitational field itself. The field equation (6.13) differs from (6.7) by the term  $\tau_a^\mu$ , making the physical meaning of the nonlinear  $S_{10}$ -theory more clear. One can check that

$$\begin{aligned} \kappa(\overline{\mathcal{D}}_\mu \tau^\mu_a + 2Q^b_{a\mu} \tau^\mu_b) &= - \left( \frac{l_0^2}{2} \overline{\mathcal{D}}_\nu R_{cd}{}^{\mu\nu} + 2Q_{cd}{}^\nu \right) R^{cd}{}_{a\mu} \\ &\quad - 4Q^b_{a\mu} \overline{\mathcal{D}}_\nu Q_b{}^{\mu\nu} + R^b_{a\mu\nu} Q_b{}^{\mu\nu}. \end{aligned}$$

The interaction with matter is introduced in accordance with the minimal coupling principle (as a result of the gauge approach) by means of a substitution  $\partial_\mu \rightarrow h_a^\mu \mathcal{D}_\mu$ . This amounts to supplementing the action  $S_{dS}$  by the interaction term

$$\left. \begin{aligned} S_{int} &= - \int \text{Tr}(A \wedge *J) = \int d^4x \sqrt{g} \mathcal{L}_{int}, \\ \mathcal{L}_{int} &= - \Gamma^a{}_{b\mu} S^{\mu b}{}_a + \frac{1}{c} t^\mu_a h_a^\mu, \quad J = \begin{pmatrix} S^a{}_b & \frac{l_0}{2c} t^a \\ -\frac{l_0}{2c} t_b & 0 \end{pmatrix}. \end{aligned} \right\} \quad (6.15)$$

Variation of total action  $S_{dS} + S_{int}$  with respect to  $\Gamma^a_b$  and  $h^a$  gives

$$\overline{\mathcal{D}}_\nu R_{ab}{}^{\mu\nu}(\Gamma) + \frac{2}{l_0^2} (Q^\mu{}_{ab} + h_a^\mu Q_b - h_b^\mu Q_a + 2Q_{[ab]}{}^\mu) = 2\lambda S^\mu{}_{ab}, \quad (6.16)$$

$$- 2\overline{\mathcal{D}}_\nu Q_a{}^{\mu\nu} + R^\mu{}_a(\Gamma) - \frac{1}{2} h_a^\mu R(\Gamma) + \Lambda h_a^\mu = \kappa (t^\mu{}_a + \tau^\mu{}_a). \quad (6.17)$$

This is the final system of the field equations of the  $S_{10}$ -gauge gravity theory.

In terms of the metric and contortion, the equations (6.16), (6.17) read

$$\begin{aligned} \frac{l_0^2}{2} \{ R_{\alpha\beta}{}^{\mu\nu}(\Gamma)_{;\nu} + T^\lambda{}_{\alpha\nu} R_{\beta\lambda}{}^{\mu\nu}(\Gamma) + T^\lambda{}_{\beta\nu} R_{\lambda\alpha}{}^{\mu\nu}(\Gamma) \} \\ + 3g^{\mu\nu} T_{[\alpha\beta\nu]} + \delta_{[\alpha}^\mu T_{\beta]} = \kappa c S^\mu{}_{\alpha\beta}, \end{aligned} \quad (6.18)$$

$$\begin{aligned} R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + g_{\mu\nu} \Lambda - T_{(\mu;\nu)} + g_{\mu\nu} T^\lambda{}_{;\lambda} + T_\lambda T^\lambda{}_{(\mu\nu)} \\ - \frac{3}{2} T_{[\alpha\beta\mu]} T^{\alpha\beta}{}_\nu - \frac{3}{2} T_{[\alpha\beta\nu]} T^{\alpha\beta}{}_\mu + \frac{1}{2} g_{\mu\nu} (T_\lambda T^\lambda + 3T_{[\lambda\alpha\beta]} T^{\lambda\alpha\beta}) \\ - \frac{l_0^2}{2} \{ R^{\alpha\beta}{}_{\mu\lambda}(\Gamma) R_{\alpha\beta\nu}{}^\lambda(\Gamma) - \frac{1}{4} g_{\mu\nu} R^{\alpha\beta}{}_{\rho\sigma}(\Gamma) R_{\alpha\beta}{}^{\rho\sigma}(\Gamma) \} = \kappa t_{(\mu\nu)}, \end{aligned} \quad (6.19)$$

$$3T_{[\mu\nu\lambda]}{}^\lambda - T_{[\mu;\nu]} + T_\lambda T^\lambda{}_{[\mu\nu]} + 3T_{[\alpha\mu\nu]} T_{\beta}{}^{\mu\nu} - 3T_{[\beta\mu\nu]} T_{\alpha}{}^{\mu\nu} = \kappa t_{[\mu\nu]}. \quad (6.20)$$

Taking the covariant derivatives of the field equations (6.16) and (6.17), after some algebra we find

$$t_{[ab]} = c \overline{\mathcal{D}}_\mu S^\mu{}_{ab}, \quad (6.21)$$

$$\overline{\mathcal{D}}_\mu t^\mu{}_a + 2Q^b_{a\mu} t^\mu{}_b + c S^\mu{}_{cd} R^{cd}{}_{\mu a} = 0. \quad (6.22)$$

These are the conservation laws of the total angular momentum and the energy-momentum of matter, which coincide with (1.47) and (1.45) when we notice that  $\overline{\mathcal{D}}_\mu S^\mu{}_{ab} = (\nabla_\mu - 2Q_\mu) S^\mu{}_{ab}$  and  $\overline{\mathcal{D}}_\mu t^\mu{}_a = (\nabla_\mu - 2Q_\mu) t^\mu{}_a$ .

When the matter is spinless and the matter Lagrangian does not depend on the tetrad otherwise than in terms of the metric, (6.22) reduces to the ordinary Einstein's conservation law of the energy-momentum tensor.

### 6.3. Physical consequences of the gauge gravity theory

The study of the  $S_{10}$ -theory generalizing the gauge approach based on the Poincaré group, allows one to fix the structure of gravitational field action via (6.10) and (6.11).

Now we investigate the physical content of the gauge gravity theory with the dynamical  $(\Gamma$ - $S$ )-interaction described by the system (6.16) and (6.17). It is worthwhile to note that these field equations represent a particular case of the general gauge gravity model with the action  $S_g = \int d^4x \sqrt{g} \mathcal{L}_g$ , where the Lagrangian  $\mathcal{L}_g = \mathcal{L}_g(h^a, Q^a, R^a_b)$  is an arbitrary scalar function of the tetrad (the metric), and the gravitational field strengths – the curvature and the torsion. The general field equations read

$$\begin{aligned} \overline{\mathcal{D}}_\nu H_{ab}{}^{\mu\nu} - E_{ab}{}^\mu &= S^\mu{}_{ab}, \\ \overline{\mathcal{D}}_\nu H_a{}^{\mu\nu} - E_a{}^\mu &= \frac{1}{c} t^\mu{}_a. \end{aligned}$$

Here the Lorentz and the translational gauge momenta are defined by

$$H^a{}_b{}^{\mu\nu} := 2 \frac{\partial \mathcal{L}_g}{\partial R^b{}_{a\mu\nu}}, \quad H_a{}^{\mu\nu} := \frac{\partial \mathcal{L}_g}{\partial Q^a{}_{\mu\nu}},$$

and the gravitational energy-momentum and spin are introduced as

$$E_a{}^\mu := h^\mu_a \mathcal{L}_g - 2Q^b{}_{a\nu} H_b{}^{\mu\nu} - R^c{}_{b a \nu} H^b{}_{c}{}^{\mu\nu}, \quad E_{ab}{}^\mu := H_{[ab]}{}^\mu.$$

#### *Correspondence with GR*

Let us investigate the issue of the correspondence of the vacuum field equations (6.18)-(6.20) and the Einstein field equations under the assumption of vanishing torsion. For the sake of generality, we do not confine ourselves to the de Sitter gravity with the action (6.17), but study the most general Lagrangian without the parity-violating terms that leads to the second-order equations for the gauge gravitational fields  $(h^a_\mu, \Gamma^a_{b\mu})$ :

$$\begin{aligned} \mathcal{L}_g &= \frac{1}{2\kappa c} \{ a_0 R(\Gamma) - 2\Lambda + a_1 Q_{\alpha\mu\nu} Q^{\alpha\mu\nu} + a_2 Q_{\alpha\mu\nu} Q^{\mu\alpha\nu} + a_2 Q_\alpha Q^\alpha \} \\ &+ b_1 R_{\alpha\beta\mu\nu}(\Gamma) R^{\alpha\beta\mu\nu}(\Gamma) + b_2 R_{\alpha\beta\mu\nu}(\Gamma) R^{\mu\nu\alpha\beta}(\Gamma) + b_3 R_{\alpha\beta\mu\nu}(\Gamma) R^{\alpha\mu\beta\nu}(\Gamma) \\ &+ b_4 R_{\alpha\beta}(\Gamma) R^{\alpha\beta}(\Gamma) + b_5 R_{\alpha\beta}(\Gamma) R^{\beta\alpha}(\Gamma) + b_6 R^2(\Gamma). \end{aligned} \quad (6.23)$$

We have the 4 dimensionless coupling constants  $a_0, a_1, a_2, a_3$ , and the 6 coupling constants  $b_1, \dots, b_6$  with the dimension  $[\hbar]$ , in addition to the cosmological term  $\Lambda$ . When  $a_0 = 0$ , we have the purely quadratic model.

Not all terms in the Lagrangian (6.23) are independent, since the expression

$$\sqrt{g} \{ R_{\alpha\beta\mu\nu}(\Gamma) R^{\mu\nu\alpha\beta}(\Gamma) - 4R_{\alpha\beta}(\Gamma) R^{\beta\alpha}(\Gamma) + R^2(\Gamma) \}$$

is the complete divergence representing the Gauss-Bonnet identity. The integral of this scalar density describes the topological invariant. Therefore, any of the constants  $b_2$ ,  $b_5$ ,  $b_6$  may be eliminated. However, here this possibility is not used.

Let us analyse the field equations for the Lagrangian (6.23) for the case of the vanishing torsion,  $Q^\lambda{}_{\mu\nu} = 0$ . Then we find  $H_a{}^{\mu\nu} = 0$ , hence  $E_{ab}{}^\mu = 0$ , and in vacuum ( $S^\mu{}_{ab} = 0$  and  $t^\mu{}_a = 0$ ) the field equations reduce to

$$\begin{aligned} \bar{\mathcal{D}}_\nu H_{ab}{}^{\mu\nu} &= -h_a^\alpha h_b^\beta \left( \varphi_1 C_{\alpha\beta}{}^{\mu\nu}{}_{;\nu} + \varphi_2 \delta^\mu{}_{[\alpha} R_{;\beta]} \right) = 0, \\ -E^\mu{}_a &= \frac{1}{\kappa c} \left( a_0 R^\mu{}_a - \frac{a_0}{2} R h_a^\mu + \Lambda h_a^\mu \right) + \varphi_1 C_{a\nu}{}^{\mu b} \mathcal{R}^\nu{}_b + \varphi_2 R \mathcal{R}^\mu{}_a = 0. \end{aligned}$$

Here  $C_{\alpha\beta}{}^{\mu\nu}$  is the Weyl tensor introduced in the irreducible decomposition of the curvature tensor (1.14), and  $\mathcal{R}_{\mu\nu} = R_{\mu\nu} - \frac{1}{4}g_{\mu\nu}R$  is the traceless part of the Ricci tensor. It is worthwhile to mention that the Weyl tensor is double-self-dual, as a result, it satisfies the quadratic identity

$$C_{\alpha\sigma}{}^{\mu\nu} C^{\beta\sigma}{}_{\mu\nu} = \frac{1}{4} \delta_\alpha^\beta C_{\rho\sigma}{}^{\mu\nu} C^{\rho\sigma}{}_{\mu\nu},$$

which was taken into account in the derivation of the field equations.

The form of the dynamical equations essentially depends on the parameters  $\varphi_1$  and  $\varphi_2$  that are constructed from the coupling constants:

$$\begin{aligned} \varphi_1 &= 2(4\lambda_1 + \lambda_2), & \varphi_2 &= \frac{4}{3}(\lambda_1 + \lambda_2 + 3\lambda_3), \\ \lambda_1 &= b_1 + b_2 + \frac{1}{2}b_3, & \lambda_2 &= b_4 + b_5, & \lambda_3 &= b_6. \end{aligned}$$

The last line determines the structure of the effective Lagrangian obtained from (6.23) for the vanishing torsion,  $Q^\lambda{}_{\mu\nu} = 0$ :

$$\mathcal{L}_g = \frac{1}{2\kappa c} (a_0 R - 2\Lambda) + \lambda_1 R_{\alpha\beta\mu\nu} R^{\alpha\beta\mu\nu} + \lambda_2 R_{\alpha\beta} R^{\alpha\beta} + \lambda_3 R^2.$$

When  $\varphi_1 = 0$  and  $\varphi_2 = 0$ , the field equations reduce to Einstein's equation with the cosmological term. This happens when  $\lambda_1 = \lambda_3$  and  $\lambda_2 = -4\lambda_3$  which corresponds to the Gauss-Bonnet topological action.

The resulting system for all values of  $\varphi_1$  and  $\varphi_2$  yields  $a_0 R = 4\Lambda$  (to check this, contract the second equation). Consequently, the torsionless ( $Q^\lambda{}_{\mu\nu} = 0$ ) vacuum field equations for the whole class of models (6.23) read

$$a_0 R = 4\Lambda, \tag{6.24a}$$

$$\varphi_1 C_{\alpha\mu\beta\nu} \mathcal{R}^{\alpha\beta} + \left( \frac{a_0}{\kappa c} + \varphi_2 R \right) \mathcal{R}_{\mu\nu} = 0, \tag{6.24b}$$

$$\varphi_1 \mathcal{R}_{\mu[\alpha;\beta]} = 0. \tag{6.24c}$$

The last equation follows from the Bianchi identity (1.17).

It is obvious that the vacuum Einstein spaces (with  $\lambda$ -term)

$$a_0 R = 4\Lambda, \quad R_{\mu\nu} = \frac{1}{4} g_{\mu\nu} R, \quad (6.25)$$

are solutions of the system (6.24a)-(6.24c). The question is: are there other solutions or (6.25) represents the unique solution? This is a non-trivial problem [137, 187]. Let us analyse the case of a nonvanishing  $a_0 \neq 0$  first; we then can put  $a_0 = 1$  without the loss of generality. There are several situations depending on the values of  $\varphi_1$  and  $\varphi_2$ .

1. When  $\varphi_1 = 0$ , the system (6.24a)-(6.24c) reduces to

$$R = 4\Lambda, \quad \left( \frac{1}{\kappa c} + 4\varphi_2 \Lambda \right) \mathcal{R}_{\mu\nu} = 0. \quad (6.26)$$

Then we have one of the two possibilities.

1.1. If  $\frac{1}{\kappa c} + 4\varphi_2 \Lambda \neq 0$ , the system (6.26) coincides with Einstein's field equations (6.25).

1.2. In the special case  $\frac{1}{\kappa c} + 4\varphi_2 \Lambda = 0$ , equations (6.24b), (6.24c) are satisfied identically, and the solutions are arbitrary spaces subject to the only constraint  $R = 4\Lambda$ .

2. If  $\varphi_1 \neq 0$ , we introduce

$$\xi := \frac{\frac{1}{\kappa c} + 4\varphi_2 \Lambda}{\varphi_1},$$

and the system (6.24a)-(6.24c) is then recast into

$$R = 4\Lambda, \quad (6.27a)$$

$$C_{\alpha\mu\beta\nu} \mathcal{R}^{\alpha\beta} = -\xi \mathcal{R}_{\mu\nu}, \quad (6.27b)$$

$$\mathcal{R}_{\mu[\alpha;\beta]} = 0. \quad (6.27c)$$

One can prove [193] that the only solutions of the system (6.27a)-(6.27c) are Einstein spaces (6.25), unless  $\xi = 0$ , or  $\xi = \frac{2\Lambda}{3}$ , or  $\xi = -\frac{4\Lambda}{3}$ . In these exceptional cases, the solutions of the system (6.27a)-(6.27c) are known with  $\mathcal{R}_{\mu\nu} \neq 0$  which are not Einstein spaces [189].

For completeness, let us mention that similar conclusions can be derived for the purely quadratic model with  $a_0 = 0$  in which case  $\xi = \varphi_2 R / \varphi_1$ .

The  $S_{10}$ -theory with the action (6.11) belongs to the class of quadratic models (6.23). As one can immediately check,  $\xi = \frac{\Lambda}{3}$  in this de Sitter model, and hence it does not belong to the exceptional cases above. Accordingly, we conclude that the only torsionless solutions of the de Sitter theory (6.11) are Einstein spaces (6.25), thus demonstrating the correspondence of the  $S_{10}$ -theory with GR.

Let us estimate the coupling constants  $\lambda$  and  $l_0$  for the  $(\Gamma$ - $S)$ -interaction. Using the torsionless solutions (6.25) (Einstein spaces), one can try to identify  $\Lambda$  directly with the cosmological term of GR. According to the modern observations,  $\Lambda \approx 10^{-52} \text{ m}^{-2}$ , consequently we find  $l_0 \approx 10^{26} \text{ m}$ , which is of order of

a typical size of the de Sitter world. As a result, assuming that  $\kappa$  is the usual Einstein's gravitational constant, we obtain for the coupling constant of the ( $\Gamma$ - $S$ )-interaction an estimate  $1/\lambda \approx 10^{120}\hbar$ . Thus, we can conclude that the spin of matter under ordinary physical conditions practically does not affect the gravitational field, and with the high accuracy we can treat (6.16) as the vacuum field equation with the zero right-hand side. However, in this case, the last term in (6.11) quadratic in the curvature would dominate at macroscopic distances, which obviously contradicts the idea to identify  $\Lambda$  with the cosmological term. Consequently, such identification and the resulting estimate of the coupling constants are not consistent with observations.

One should expect the coupling constant  $\lambda$  to be much larger (with the upper limit for  $\lambda$  to be fixed by experiment), so that the curvature quadratic term dominates only at small distances, whereas at the macroscopic scales it is essentially smaller than the term  $(R + Q^2)$ . Then the resulting huge value of  $\Lambda$  does not allow to interpret it as a cosmological term. The correspondence with Einstein's equations was derived here in the absence of matter, which is a special physical situation. When  $t_{\mu\nu} \neq 0$ , the condition of the vanishing torsion turns out to be too strong even in the absence of spin  $S^\lambda_{\mu\nu} = 0$ , and it does not lead to reasonable results, in general [100, 129, 140, 141].

### *Instanton solutions*

The Yang-Mills structure of the action (6.10) of the de Sitter theory underlies the existence of the self-dual or the instanton solutions. To analyse this important feature, we turn to the Euclidean formulation of the theory. Technically this is done via the replacements  $SO(1,4) \rightarrow SO(5)$ ,  $SO(1,3) \rightarrow SO(4)$ , and  $\eta_{ab} \rightarrow \delta_{ab}$ ; one achieves this by the Wick rotation of the time coordinate into the purely imaginary domain  $t \rightarrow it_E$ . The self-duality conditions for the de Sitter curvature (6.10) read

$$\mathcal{R} = *\mathcal{R}, \quad (6.28)$$

and this is written explicitly as the system

$$\left(R^a{}_b - \frac{1}{l_0^2}\pi^a{}_b\right) = *\left(R^a{}_b - \frac{1}{l_0^2}\pi^a{}_b\right), \quad (6.29)$$

$$\widehat{Q}^a = *\widehat{Q}^a. \quad (6.30)$$

On the self-dual configurations (6.29), (6.30), the vacuum field equations (6.12) and (6.13) are satisfied due to generalized Ricci (1.15) and Bianchi (1.17) identities which can be conveniently recast into

$$2\overline{\mathcal{D}}_{[\lambda}Q^a{}_{\mu\nu]} = -R^a{}_{b[\mu\nu}h^b{}_{\lambda]}, \quad \overline{\mathcal{D}}_{[\lambda}R^a{}_{\mu\nu]} = 0.$$

The solutions of (6.29), (6.30) are called gravitational instantons. The Euclidean action

$$S = \frac{1}{8\lambda} \int \left\{ \left(R^{ab} - \frac{1}{l_0^2}\pi^{ab}\right) \wedge * \left(R_{ab} - \frac{1}{l_0^2}\pi_{ab}\right) + \frac{2}{l_0^2} \widehat{Q}^a \wedge *\widehat{Q}_a \right\}, \quad (6.31)$$

reaches a local extremum (minimum) on the instanton configurations, when it becomes proportional to the topological invariant

$$- \int \text{Tr}(\mathcal{R} \wedge \mathcal{R}) = \int \left\{ R^{ab} \wedge R_{ab} + \frac{2}{l_0^2} \left( \widehat{Q}^a \wedge * \widehat{Q}_a - R^{ab} \wedge \pi_{ab} \right) \right\}.$$

The first term (with an appropriate numeric coefficient) is equal to the Pontryagin index, whereas the second term describes the Nieh-Yan topological invariant.

In the theory under consideration, the Euclidean action is well defined (unlike the Euclidean action in GR and ECT), leading to the formally convergent path integral in the quantum theory.

The absolute minimum of (6.31) is realized on the Euclidean de Sitter space

$$\mathcal{R} = 0 \quad \left\{ Q^a{}_{\mu\nu} = 0, \quad R^{ab}{}_{\mu\nu} = \frac{2}{l_0^2} h_\mu^{[a} h_\nu^{b]} \right\}, \quad (6.32)$$

on which the action vanishes,  $S = 0$ .

It is worthwhile to note that using the ansatz  $R^\alpha{}_{\beta\mu\nu} = \frac{2}{l_0^2} \delta^\alpha_{[\mu} g_{\nu]\beta}$  when solving the complete system of the vacuum field equations (6.18)-(6.20) automatically yields the zero torsion. This result agrees with the original assumptions: in the  $S_{10}$ -theory, the simplest solution is the de Sitter space of the constant curvature, but not the Minkowski flat space.

One can straightforwardly describe the torsionless instantons. When  $Q = 0$ , the contraction of the equations (6.29) gives (due to the Ricci identity) the Einstein equations with the  $\lambda$ -term. Thus, in the theory under consideration, the Einstein spaces constitute the subset of the instanton solutions, and it is quite natural to interpret the solutions of Einstein's equations as gravitational instantons in this approach.

Let us consider in detail the important special case: the spherical  $SO(4)$ -symmetric instanton solutions of (6.29) and (6.30). The general spherically symmetric (in the four-dimensional  $SO(4)$  sense) ansatz for the local Lorentz connection and the tetrad reads as follows:

$$\Gamma^a{}_{b\mu} = A(x^a \delta_{b\mu} - x_b \delta_\mu^a) + B \varepsilon^a{}_{b\mu\nu} x^\nu, \quad (6.33)$$

$$h_\mu^a = f \delta_\mu^a + g x^a x_\mu, \quad (6.34)$$

where  $A = A(\rho)$ ,  $B = B(\rho)$ ,  $f = f(\rho)$ ,  $g = g(\rho)$ , are the scalar functions of  $\rho = x^a x_a = \delta_{ab} x^a x^b$ .

Substituting (6.33), (6.34) to (6.29), we obtain the system

$$2\rho A' f + 2A f \mp \left( 2B - 2AB\rho + \frac{f^2}{l_0^2} \right) (f + \rho g) = 0, \quad (6.35)$$

$$2(A \pm B)' f - 2(A \pm B)g + (A \pm B)^2 (f + \rho g) = 0. \quad (6.36)$$

Here the dash denotes derivatives with respect to  $\rho$ , and the upper signs correspond to the self-duality  $(R - l_0^{-2}\pi) = *(R - l_0^{-2}\pi)$ , whereas the lower signs

refer to the anti-self-duality  $(R - l_0^{-2}\pi) = - * (R - l_0^{-2}\pi)$ . In a similar way, for the torsion we derive from (6.30) one more equation

$$2f' - g + (A \pm 2B)(f + \rho g) = 0. \quad (6.37)$$

The spherical  $SO(4)$ -symmetric ansatz (6.33) and (6.34) excludes the traceless part of the torsion. However, the trace and pseudotrace are non-trivial:

$$Q_\mu = -\frac{3}{2f}x_\mu \{2f' - g + A(f + \rho g)\}, \quad \check{Q}_\mu = -\frac{3}{2f}x_\mu B(f + \rho g),$$

and the self-duality equation (6.37) establishes their relation (it is obvious that for such configuration the trace  $Q^\mu$  is proportional to the pseudotrace  $\check{Q}^\mu$ ). This implies the first result: any self-dual  $SO(4)$ -symmetric solution (6.33), (6.34) describes the Riemannian spacetime (there is no torsion), if either  $B = 0$  or  $f + g\rho = 0$ .

All Riemannian  $SO(4)$ -instantons may be described explicitly.

When  $f + g\rho = 0$ , we obtain a degenerate geometry. Indeed, then  $g = -f/\rho$  and thus the tetrad  $h_\mu^a = f(\delta_\mu^a - x^a x_\mu/\rho)$  is a projector with the vanishing determinant. The same applies to the corresponding Euclidean spacetime metric  $g_{\mu\nu} = f^2(\delta_{\mu\nu} - x_\mu x_\nu/\rho)$  which represents an unphysical geometry.

When  $B = 0$ , there exists an infinite family of the  $SO(4)$ -symmetric solutions of the system (6.35)-(6.37):

$$(A\rho - 1)^2 = 1 \mp \frac{f^2\rho}{l_0^2}, \quad g(1 - A\rho) = 2f' + Af.$$

We can express  $A$  and  $g$  in terms of an arbitrary function  $f$ , provided  $A\rho \neq 1$ . For the self-dual case, we have an exceptional solution  $A = 1/\rho$ ,  $f = l_0/\sqrt{\rho}$  with an arbitrary  $g$ . This is a singular solution, but the infinite family encompasses the regular configurations, as well. The famous de Sitter instanton is recovered from the family above by fixing  $g = 0$ :

$$A = \frac{2}{\rho \pm \rho_0}, \quad f = \frac{2l_0\sqrt{\rho_0}}{\rho \pm \rho_0}, \quad g = 0, \quad B = 0.$$

As usual, the upper (lower) sign refers to the self-duality (anti-self-duality). For the positive real integration constant  $\rho_0$ , the self-dual solution is everywhere regular, but the anti-self-dual solution diverges at the radius  $\rho = \rho_0$ .

The search for the non-Riemannian ( $Q^a{}_{\mu\nu} \neq 0$ )  $SO(4)$ -instantons represents a nontrivial task. There is no their complete description yet, and specific solutions are difficult to interpret. A preliminary analysis can be found in [194].

Possible existence of the instanton configurations is consistent with the idea of Hanson and Regge [145] who noticed the analogy of the gravity with  $Q \neq 0$  and the superconductivity theory. The vanishing torsion is then dynamically realized as a phase of the Meissner type effect, and the local regions with nontrivial torsion  $Q \neq 0$  are the analogs of Abrikosov's vortices.

*Torsion dynamics and universal spin-spin interaction*

Let us consider the physically most important particular type of matter – the fermion fields (describing quarks, leptons, hadrons) as a source of the gauge gravitational field. Then the tensor of spin is totally antisymmetric and has the following form:

$$S_{\lambda\mu\nu} = \varepsilon_{\lambda\mu\nu\kappa} \check{S}^\kappa, \quad \check{S}_\mu = \frac{i\hbar}{4} \bar{\Psi} \gamma_\mu \gamma_5 \Psi.$$

Substitution of the ansatz  $R^a_b = \frac{1}{l_0^2} \pi^a_b$  into the field equations of the de Sitter model (6.18)-(6.20) yields the Riemannian de Sitter space in the absence of matter, but for the nontrivial fermion sources this ansatz gives rise to the algebraic relation between the torsion and the spin  $\check{T}_\mu = \frac{\kappa c}{3} \check{S}_\mu$  via (6.18), with the “translational” equation (6.19) reducing to the Einstein equation with the  $\lambda$ -term and the symmetrical part of the canonical energy-momentum tensor on the right-hand side.

Let us qualitatively investigate the dynamics of the torsion field in the de Sitter gauge gravity theory. For this purpose, we consider a simplified model: we assume the metric to be flat  $g_{\mu\nu} = \eta_{\mu\nu}$ ,  $h_\mu^\alpha = \delta_\mu^\alpha$ , and the torsion is represented only by the pseudotraces vector  $\check{T}_\mu$ , which is natural if the sources are fermions. Then the action describing the torsion dynamics takes the following form:

$$S_{\check{T}} = \int d^4x \sqrt{g} \left\{ \frac{1}{\lambda} \left[ \partial_\mu \check{T}_\nu \partial^\mu \check{T}^\nu + \frac{1}{2} (\partial_\mu \check{T}^\mu)^2 + \mu^2 \check{T}_\mu \check{T}^\mu - \frac{3}{2} (\check{T}_\mu \check{T}^\mu)^2 \right] + \check{S}_\mu \check{T}^\mu \right\}, \quad (6.38)$$

where  $\mu^2 = 9\lambda/\kappa c = 9/l_0^2$ .

We can use the method of Gupta [195] to calculate the static non-relativistic interaction potential between the two fermions with spins  $\sigma_1$  and  $\sigma_2$  exchanging by the  $\check{T}$ -quanta. In the linear approximation we obtain

$$V_{\check{T}}(\mathbf{r}) = -\frac{\lambda c \hbar^2}{8\pi} \left\{ (\sigma_1 \cdot \sigma_2) \frac{e^{-\mu r}}{r} - \frac{1}{\mu^2} (\sigma_1 \cdot \nabla) (\sigma_2 \cdot \nabla) \left( \frac{e^{-\mu r}}{r} - \frac{e^{-\sqrt{\frac{2}{3}} \mu r}}{r} \right) \right\}.$$

In the limit of the zero transferred momentum in  $\check{T}$ -propagators ( $\mu \neq 0$ ), one finds the contact potential like in the ECT. Therefore, the presence of the Yang-Mills (quadratic in the curvature) term in (6.11) modifies ECT in the same way as the theory of intermediate bosons modifies the Fermi theory of weak interactions.

Qualitatively the same result for the interaction potential arises also for the most general quadratic gauge gravity model (6.23), where one needs to replace the parameters  $\lambda$ ,  $\mu^2$  and  $2/3$  with the appropriate algebraic expressions constructed from the coupling constants  $a_1, \dots, a_3$  and  $b_1, \dots, b_6$ , see the details in [196, 197, 198].

The spin-spin potential is also found [146] in GR for the fermions interacting via the graviton exchange, where  $V_G \sim \kappa c^2 \hbar^2 (\sigma_1 \cdot \sigma_2) / r^3$  (analog of the Breit potential). In contrast to  $V_G$ , the dynamical gauge ( $\Gamma$ - $S$ )-interaction predicts the  $1/r$  spin-spin interaction (similar to Coulomb’s and Newton’s potentials).

## 6.4. Model description of microscopic gravitational interactions

We now turn to the study of the microscopic limit of the gauge gravity theory. Addressing this problem one can relax the requirements formulated in Sec. 6.1. for the construction of the dynamics, in particular it makes no sense to talk about the correspondence with GR at small distances. Moreover, in accordance with the gauge principles in the framework of the geometrodynamical approach (following Einstein's idea that the spacetime geometry is not fixed ad hoc, but is determined by the interaction and the motion of matter), one can assume that at small scales we may need to consider new microscopic physical properties of matter, which did not manifest themselves at the macroscopic level. Understanding the geometrization program in a general sense, the new physics should lead to the new geometry. Accordingly, the minimal model based on the Poincaré group (or its extension to the de Sitter group) may turn out to be too narrow in the area of the high energies (very small distances).

Most probably, the  $\mathcal{P}_{10}$ -theory provides a good description for the gravitational interactions of leptons, since their spacetime symmetries are exhausted by the Poincaré group. However, there are additional symmetries in the physics of strongly interacting particles [147] besides the  $\mathcal{P}_{10}$ . In particular, the systematics of hadrons within the Regge approach [148] suggests the description of all particles lying on a Regge trajectory as excitations of one physical object. Such states are classified by the unitary representations of  $SL(3, R)$  group, or in relativistic theory by  $SL(4, R)$ . In addition, in the high energy strong interactions one observes another (asymptotic) regularity – the Bjorken scaling [149] as a manifestation of the dilational symmetry [140]. Together, the dilations and the  $SL(4, R)$  constitute the general linear group and, adding the translations, we end up with the general affine group  $GA(4, R) = GL(4, R) \rtimes T_4$ .

Therefore, one can suggest that the gravitational interactions of the hadrons should be described not by the  $\mathcal{P}_{10}$ -theory, but rather by a more general theory based on  $GA(4, R)$ . The corresponding gauge theory is called a metric-affine gravity; for its comprehensive overview see [199].

In Secs. 5.3. and 5.4., we constructed the kinematics of the gauge theory for the general affine group. The gauge potential is identified with the generalized affine connection in the principal bundle of affine frames  $A(M)$ , which introduces the most general geometrical structure on the spacetime manifold  $M_4$ : the independent linear connection  $\Gamma^\alpha_{\beta\mu}$  and the metric  $g_{\mu\nu}$  (the tetrad fields  $h^a_\mu$ ). The sources of such a microscopic gravitational field are the corresponding Noether  $GA(4, R)$ -currents of matter: the energy-momentum and the “hypermomentum” [147, 199]

$$J^\mu_{ab} = \frac{\partial \mathcal{L}_m}{\partial \partial_\mu \Phi} I_{ab} \Phi. \quad (6.39)$$

Here  $\mathcal{L}_m$  is the Lagrangian of the matter fields  $\Phi$ , and  $I_{ab}$  are the generators of  $GL(4, R)$ . The irreducible parts of the hypermomentum (6.39) are the spin  $S^{\mu ab} = J^{\mu [ab]}$ , the dilation current  $J^\mu = J^{\mu ab}\eta^{ab}$ , and the ‘‘proper hypermomentum’’  $\bar{J}^{\mu ab} = J^{\mu (ab)} - \frac{1}{4}J^\mu\eta_{ab}$ , corresponding to shear deformations of the frame.

### *Microscopic gravitation as a Yang-Mills field*

Now we consider the microscopic gravitation model  $(\Gamma^a_{b\mu}, h^a_\mu)$  assuming that the linear connection  $\Gamma^a_{b\mu}$  is the most general one. The dynamics of the theory at small distances can be formally obtained by discarding the ‘‘translational’’ low-energy terms in the basic Lagrangian (6.11) or taking a formal limit  $l_0 \rightarrow \infty$ . Thus, we will describe the model microscopic theory of the gravitational interactions by the Lagrangian of the Yang-Mills type for the  $GL(4, R)$ -gauge field

$$\mathcal{L}_{YM} = \frac{1}{8\lambda} R^a_{b\mu\nu}(\Gamma) R^b_a{}^{\mu\nu}(\Gamma). \quad (6.40)$$

The invariance group of the theory (6.40) is very wide. It includes the conformal Weyl group  $g_{\mu\nu} \rightarrow g'_{\mu\nu} = e^{2\sigma(x)}g_{\mu\nu}$  (see Sec. 4.2.); the local linear  $GL(4, R)$ -group  $h^a_\mu \rightarrow h'^a_\mu = L^a_b h^b_\mu$ ,  $\Gamma^a_{b\mu} \rightarrow \Gamma'^a_{b'\mu} = L^a'_a \Gamma^a_{b\mu} L^{-1b}_{b'} + L^a'_c \partial_\mu L^{-1c}_{b'}$  with  $L \in GL(4, R)$  and the group of general coordinate transformations: for example,  $h^a_\mu \rightarrow h^a_{\mu'} = \frac{\partial x^\mu}{\partial x^{\mu'}} h^a_\mu$ .

The field equations are derived by the variation of (6.40) with respect to the independent  $\Gamma^a_{b\mu}$  and  $g_{\mu\nu}$ . In vacuum, they read

$$\begin{aligned} \bar{\mathcal{D}}_\mu R^a_b{}^{\mu\nu}(\Gamma) &\equiv \frac{1}{\sqrt{g}} \partial_\mu \{ \sqrt{g} R^a_{b\alpha\beta}(\Gamma) g^{\alpha\mu} g^{\beta\nu} \} \\ &+ g^{\alpha\mu} g^{\beta\nu} \{ \Gamma^a_{c\mu} R^c_{b\alpha\beta}(\Gamma) - \Gamma^c_{b\mu} R^a_{c\alpha\beta}(\Gamma) \} = 0, \end{aligned} \quad (6.41)$$

$$\tau_{\mu\nu} = -\frac{c}{2\lambda} \left\{ R^a_{b\mu\lambda}(\Gamma) R^b_{a\nu}{}^\lambda(\Gamma) - \frac{1}{4} g_{\mu\nu} R^a_{b\alpha\beta}(\Gamma) R^b_a{}^{\alpha\beta}(\Gamma) \right\} = 0. \quad (6.42)$$

Compare the last equation with (6.14).

Technically, the equation (6.42) means that the energy-momentum tensor of the Yang-Mills field vanishes, which therefore is often called a ghost field.

It is convenient to introduce the conformally invariant tensor density  $\pi^{\mu\nu\alpha\beta} = \sqrt{g} (g^{\mu\alpha} g^{\nu\beta} - g^{\nu\alpha} g^{\mu\beta})$ . The metric enters the theory only in terms of  $\pi^{\mu\nu\alpha\beta}$ , in particular, the action reads  $S_{YM} = \frac{1}{16\lambda} \int d^4x \pi^{\mu\nu\alpha\beta} R^a_{b\mu\nu}(\Gamma) R^b_{a\alpha\beta}(\Gamma)$  and the field equation (6.41) can be rewritten as

$$\partial_\mu \{ R^a_{b\alpha\beta}(\Gamma) \pi^{\mu\nu\alpha\beta} \} + \{ \Gamma^a_{c\mu} R^c_{b\alpha\beta}(\Gamma) - \Gamma^c_{b\mu} R^a_{c\alpha\beta}(\Gamma) \} \pi^{\mu\nu\alpha\beta} = 0. \quad (6.43)$$

The importance of the Yang-Mills fields motivates the interest in the exact solutions of the classical gauge field equations. It is known [150, 151] that any spherically symmetric solution of the Yang-Mills equations for the internal groups [ $SU(2)$  etc.] in the Minkowski space has the ‘‘Coulomb’’ type behaviour.

Let us study the spherically symmetric solutions of (6.43) in the Riemannian space-time with a given background metric  $g_{\mu\nu}(x^\lambda)$ . In the local spherical coordinates  $(x^0 = t, x^1 = r, x^2 = \theta, x^3 = \varphi)$  for an arbitrary spherically symmetric metric  $ds^2 = -\xi(r, t)dt^2 + \eta(r, t)dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\varphi^2$ , the most important component of the density  $\pi^{\mu\nu\alpha\beta}$  reads as follows:

$$\pi^{1010}(t, r, \theta, \varphi) = f(r, t)\chi(\theta, \varphi), \quad (6.44)$$

where  $\xi, \eta, f, \chi$  are the functions of the specified arguments. Under the condition of the spherical symmetry, we assume that  $\Gamma^a{}_{b0}(r, t)$  depends on the time and the radial coordinates, whereas  $\Gamma^a{}_{b1} = \Gamma^a{}_{b2} = \Gamma^a{}_{b3} = 0$ . Then one can check that the solution of (6.43) reads

$$\Gamma^a{}_{b0}(r, t) = A^a{}_b \int_{r_0}^r \frac{dr'}{f(r', t)}, \quad R^a{}_{b10} = \Gamma^a{}_{b0} = \frac{\partial \Gamma^a{}_{b0}}{\partial r}, \quad (6.45)$$

where the constant matrix  $A \in GL(4, R)$ . In [152], it is proved that this is a general (up to a gauge transformation) spherically symmetric solution of the equations (6.43).

In addition, when the matrix  $A$  is such that  $A^a{}_b A^b{}_a = 0$ , eq. (6.45) describes the exact solutions of the complete system of the microscopic gravitation (6.41) and (6.42), since then  $\tau_{\mu\nu} = 0$  identically. Such matrices do exist in  $GL(4, R)$ .

To illustrate (6.45), we consider several special background metrics.

1. Let  $f(r, t) = -r^2$ . This is true for the Schwarzschild, the Reissner-Nordström, the de Sitter, the Kottler metrics (the spherical vacuum solution of Einstein's equations with the cosmological term), the conformally flat metrics (in particular, for the Friedman metric) and for some others. In this case,  $R^a{}_{b10} = -A^a{}_b/r^2$ , and  $\Gamma^a{}_{b0} = A^a{}_b(1/r - 1/r_0)$ . The solution thus has the "Coulomb" form, generalizing the result of [150] to the aforementioned Riemannian spaces.

2. Let us consider the cosmological metrics

$$ds^2 = -dt^2 + a^2(t) \left\{ \frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \right\}, \quad k = \text{const.}$$

Then  $f(r, t) = -a(t)r^2\sqrt{1 - kr^2}$  and

$$\Gamma^a{}_{b0}(r, t) = \frac{A^a{}_b}{a(t)} \left( \frac{\sqrt{1 - kr^2}}{r} - \text{const} \right), \quad R^a{}_{b10} = -\frac{A^a{}_b}{a(t)r^2\sqrt{1 - kr^2}}.$$

For particular cases, we have:

a)  $k = 1/r_0^2$  is a closed universe, with  $a(t) = 1$  this is the static Einstein's world. The gravitational field strength  $R^a{}_{b10}(r, t)$  has singularities at  $r = 0$  and at  $r = r_0$ , at the boundary or the world. The region  $r > r_0$  is unphysical. Thus, in the Einstein world, the spherically symmetric solutions of the Yang-Mills equations have a non-Coulomb short-range behaviour.

b)  $k = -1/r_0^2$  is an open universe,  $a(t) = 1$ . The solution is asymptotically Coulomb one.

c)  $k = 0$ ,  $a(t) = e^{Ht}$ , ( $H = \text{const}$ ) is the inflationary cosmological de Sitter metric in the Lemaitre-Robertson form [120]. In this case

$$R^a{}_{b10}(r, t) = -\frac{A^a{}_b}{r^2} e^{-Ht}, \quad \Gamma^a{}_{b0}(r, t) = A^a{}_b(1/r - 1/r_0) e^{-Ht}.$$

A local observer interprets this solution as a Coulomb type configuration with a decreasing gauge charge (compare this with Dirac's hypothesis [68] about the change of constants in time).

Introducing the interaction of the microscopic gravity with matter, the field equations are modified by the relevant sources: the hypermomentum (6.39) is added in the right-hand side of (6.41) and the energy-momentum tensor  $t_{\mu\nu}$  in (6.42). The conformal symmetry of the model (6.40) give rise to the restriction  $t^\mu{}_\mu = 0$  that allows only the massless matter fields as the sources of the  $(\Gamma, g)$ -gravity. Examples of such fields are the massless spinor fields and the Yang-Mills fields in the theories of "grand unification" (before a spontaneous symmetry breaking), which underlines the essentially microscopic nature of model under consideration.

Below, we investigate the question how this microscopic picture is related to the macroscopic theory of gravity.

### *Spontaneous breaking of conformal symmetry and macroscopic limit*

The structure of the microscopic gravity theory does not formally allow for the interaction with the macroscopic, massive matter. Therefore, in order to preserve the conformal invariance of a massive field action, we introduce an auxiliary scalar field  $\varphi(x^\lambda)$  which transforms under the action of the Weyl group as

$$\varphi \longrightarrow \varphi' = e^{-\sigma(x)} \varphi(x).$$

If we now define the masses of a vector  $A_\mu$  and a spinor  $\Psi$  fields as  $m = \varphi \overset{\circ}{m}$ , with  $\overset{\circ}{m} = \text{const}$ , one can add the corresponding mass term to the complete matter Lagrangian.

It is important to determine the Lagrangian of the dynamical field  $\varphi$  to set the transition to the macroscopic theory. There are two possibilities, which are based on the different interpretation of the Weyl symmetry that we analyzed in Sec. 4.2.. Let us consider both cases.

#### *a) Canonical Weyl approach and correspondence with GR*

The usual interpretation of Weyl's transformations  $g_{\mu\nu} \rightarrow e^{2\sigma} g_{\mu\nu}$  is based on the *expansion of lengths*, and in particular, on the scaling of the interval  $ds \rightarrow ds' = e^\sigma ds$ . If we understand the conformal symmetry of the microscopic gravity theory in this sense, the most general Weyl-invariant action for the

scalar field  $\varphi$  has the form:

$$S_\varphi = \int d^4x \sqrt{g} \hbar \left\{ \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi + \frac{R}{12} \varphi^2 - \Lambda_0 \varphi^4 \right\}, \quad \Lambda_0 = \text{const.}$$

We assume that the scalar field has the canonical dimension of an inverse length, whereas  $\Lambda_0$  is dimensionless. Fixing the Weyl freedom by the gauge condition  $\varphi = \varphi_0 = \text{const}$  [73, 149] (which corresponds to spontaneous breaking of conformal symmetry) one finds the complete action of the theory

$$S = \int d^4x \sqrt{g} \left\{ \mathcal{L}_{YM}(\Gamma) + \frac{1}{c} \mathcal{L}_m(\Psi, A) + \frac{1}{2\kappa c} (R - 2\Lambda) - \frac{1}{2} \varphi_0^2 \overset{\circ}{m}_A^2 A_\mu A^\mu + \varphi_0 \overset{\circ}{m}_\Psi \bar{\Psi} \Psi \right\}. \quad (6.46)$$

Here  $\mathcal{L}_m(\Psi, A)$  is the free Lagrangian of the massless vector  $A_\mu$  and spinor  $\Psi$  fields, and  $\kappa c = 6/(\hbar \varphi_0^2)$ ,  $\Lambda = 6\Lambda_0 \varphi_0^2$ .

On the macroscopic scales  $\gg 1/\varphi_0$ , we have  $\frac{1}{2\kappa c} R \gg \mathcal{L}_{YM}(\Gamma)$ , i.e., the Hilbert-Einstein term is dominating. Consequently, the macroscopic limit in this case is described by GR with the cosmological term.

This procedure, that introduces the masses and the Lagrangian  $R$ , is the classical analog of the quantum mechanism of the dynamical breaking of the conformal symmetry, and the field  $\varphi$  plays the role of the Goldstone boson [153].

Assuming that the coupling self-interaction constant of the scalar field is small ( $\Lambda_0 \rightarrow 0$ ), one can ignore the  $\Lambda$ -term in (6.46), and hence the gravitational Lagrangian has the form  $\mathcal{L}_0 = \frac{1}{2\kappa c} R + \frac{1}{8\lambda} R^a{}_{b\mu\nu}(\Gamma) R^b{}_{a}{}^{\mu\nu}(\Gamma)$ , described as the case (c) in Sec. 6.1.. The corresponding field equations read:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa (t_{\mu\nu} + \tau_{\mu\nu}), \quad (6.47)$$

$$\overline{D}_\nu R^a{}_{b}{}^{\mu\nu}(\Gamma) = 2\lambda J^{\mu a}{}_b. \quad (6.48)$$

The exact spherically symmetric solution of the vacuum equations (6.47), (6.48) (for the similar solutions for an arbitrary gauge group see in [99, 100, 152]) is straightforwardly derived:

$$ds^2 = -e^\nu dt^2 + e^{-\nu} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2, \\ e^\nu = 1 - \frac{2Gm}{c^2 r} + \frac{\kappa c q^2}{4\lambda r^2}$$

for the metric, and for the gauge field  $\Gamma^a{}_{b\mu}$ :

$$R^a{}_{b01} = \frac{u^a{}_b}{r^2}, \quad R^a{}_{b23} = v^a{}_b \sin \theta,$$

where  $q^2 = -u^a{}_b u^b{}_a - v^a{}_b v^b{}_a$  and the dimensionless constant matrices  $u^a{}_b, v^a{}_b \in GL(4, R)$ . This solution is of the Reissner-Nordström type. The essential difference is that the value of the “effective charge”  $q^2$  may be greater, equal or

smaller than zero, in view of the non-compactness of the group  $GL(4, R)$ . Formally,  $q^2$  is composed of the three irreducible parts, since  $u$  and  $v$  are connected with the spin, the dilation current and the proper hypermomentum of the central body:  $u_{ab} = \frac{1}{4}\eta_{ab}u + u_{[ab]} + \bar{u}_{ab}$ ,  $\bar{u}^a{}_a = 0$ . The case  $q^2 = 0$  belongs to the class of the abovementioned ghost solutions of the Yang-Mills equations.

The non-compactness of the symmetry group may have some specific consequences in the gauge gravity theory. For example, the indefiniteness of the sign of  $q^2$  may prevent the formation of a horizon, as well as singularities may be avoided in cosmology. This is related to the fact that the sign of the energy  $\tau_{00}$  of the field  $\Gamma$  is indefinite, which leads to the violation of the energy conditions of the Penrose-Hawking theorems. Indeed, denoting  $E^a{}_i = R^{ab}{}_{0i}$  and  $B^{ab}{}_{ij} = R^{ab}{}_{ij}$  (Latin indices from the middle of the alphabet  $i, j, k, \dots = 1, 2, 3$  label 3-space components), we can decompose the curvature into the four pieces

$$\overset{e}{\mathcal{E}}{}^k{}_i = E^{0k}{}_i, \quad \overset{e}{\mathcal{B}}{}^{kj}{}_i = E^{kj}{}_i, \quad \overset{m}{\mathcal{E}}{}^k{}_{ij} = B^{0k}{}_{ij}, \quad \overset{m}{\mathcal{B}}{}^{kl}{}_{ij} = B^{kl}{}_{ij}.$$

Then using the self-evident symbolic notation we find

$$\tau_{00} = \frac{c}{2\lambda} \left( \overset{e}{\mathcal{B}}{}^2 + \overset{m}{\mathcal{B}}{}^2 - \overset{e}{\mathcal{E}}{}^2 - \overset{m}{\mathcal{E}}{}^2 \right),$$

that clearly demonstrates that the energy density is not positive definite.

This can be observed as an ‘‘antigravity’’ effect, or a gravitational repulsion, that would be manifest as an effective reduction of mass. One can qualitatively illustrate this as follows. Suppose the matter source in (6.47), (6.48) is a massive cylinder made of a ferromagnetic material that can be magnetized along its axis  $z$ . The dilation current and the proper hypermomentum are both equal zero and the model thus reduces to an effective  $T_4 \times L_6$ -theory. Under the assumption of an approximately flat metric  $g_{\mu\nu} \approx \eta_{\mu\nu}$ , the polarized state of a homogeneous sample can be modeled by the spinor field  $\Psi = \sqrt{n} e^{imt} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$ , so that the spin pseudovector is  $\check{S}^\mu = \{0, 0, 0, -\frac{\hbar}{2}n\}$ , where  $n$  is the particle density of the polarized spins. In the linear approximation, neglecting the boundary effects, we then find the internal solution of the equation (6.48) for the components of the curvature:  $R_{ab}{}^{0i} = \frac{2}{3}\lambda x^i S^0{}_{ab}$ ,  $R_{ab}{}^{ik} = \lambda(x^i S^k{}_{ab} - x^k S^i{}_{ab})$ . This yields  $\tau_{00} = -\lambda c \hbar^2 n^2 (5r^2 + 9z^2)/36$ , and we obtain the correction to the energy caused by the  $\Gamma$ -field by evaluating the integral over the cylinder source:

$$\Delta E = \int d^3x \tau_{00} = -\frac{\lambda c \hbar^2 n^2}{36} V \left( \frac{5\rho_0^2}{2} + \frac{14\ell_0^2}{3} \right) < 0,$$

where sample’s geometry is specified by the radius  $\rho_0$ , the length  $\ell_0$ , and the volume  $V = \pi\rho_0^2\ell_0$ . To make a numeric estimate, let us take  $\rho_0 \sim 10^{-2}$  m,  $\ell_0 \sim 10^{-1}$  m, and  $n = 10^{29} \text{ m}^{-3}$ . Then we conclude, that upon an instantaneous magnetization of the sample, its mass should effectively reduce by  $\Delta M = \Delta E/c^2 \approx -\lambda \hbar \times 10^{-8}$  kg, or the relative reduction should be

$\Delta M/M \approx -\lambda\hbar \times 10^{-7}$ . This is a tiny effect for the value  $\lambda\hbar = 10^{-120}$  of the coupling constant estimate obtained in Sec. 6.3.. The current qualitative analysis shows that the gauge gravity coupling constant can be significantly larger, and even  $\lambda\hbar \sim 1$  is a viable estimate. The actual value of  $\lambda$  should be fixed by future experiments.

b) *Internal conformal symmetry and model GA(4, R)-theory*

In another approach, which was developed earlier in Sec. 4.2., the Weyl transformations are interpreted as the tetrad scaling  $h_\mu^a \rightarrow e^\sigma h_\mu^a$ . In this case, the most general conformally invariant action for the auxilliary scalar field  $\varphi$  has the form  $\tilde{S}_\varphi = \int d^4x \sqrt{g} \hbar \{R(\Gamma)\varphi^2 - \Lambda_0\varphi^2\}$  (the kinetic term for  $\varphi$  is not added since it violates the conformal invariance). Then, fixing the gauge  $\varphi = \varphi_0 = \text{const}$  as a result of the spontaneous symmetry breaking, one introduces masses for the matter fields and finds the effective gravitational Lagrangian, which now has the following form (it was described as the case (d) in Sec. 6.1.):

$$\mathcal{L}_1 = \frac{1}{2\kappa c} R(\Gamma) + \frac{1}{8\lambda} R^a{}_{b\mu\nu}(\Gamma) R^b{}_{a}{}^{\mu\nu}(\Gamma). \quad (6.49)$$

Thereby, we obtain the model gauge theory for the general affine group investigated in [18]. The peculiar feature of this  $(g_{\mu\nu}, \Gamma^a{}_{b\mu})$ -theory is the presence of an additional Abelian long-distance field in it – the Weyl vector  $K_\mu$ . Using the irreducible decomposition of  $GL(4, R)$ -connection  $\Gamma^a{}_b = \Gamma^a{}_{b\mu} dx^\mu$  1-form

$$\Gamma^{ab} = \hat{\Gamma}^{ab} - \frac{1}{2} K^{ab}$$

into the local Lorentz connection  $\hat{\Gamma}^{ab} = \Gamma^{[ab]}$  and the nonmetricity<sup>1</sup> 1-form  $K_{ab} := D\eta_{ab} = -2\Gamma^{(ab)}$ , we find the relevant decomposition of  $GL(4, R)$ -curvature  $R^a{}_b = \frac{1}{2} R^a{}_{b\mu\nu}(\Gamma) dx^\mu \wedge dx^\nu$  2-form

$$R^a{}_b = \hat{R}^a{}_b + \frac{1}{4} \bar{K}^a{}_c \wedge \bar{K}^c{}_b - \frac{1}{2} \hat{D}\bar{K}^a{}_b + \frac{1}{4} \delta_b^a \Omega \quad (6.50)$$

into the local Lorentz curvature 2-form  $\hat{R}^a{}_b = d\hat{\Gamma}^a{}_b + \hat{\Gamma}^a{}_c \wedge \hat{\Gamma}^c{}_b$  and the homothetic curvature<sup>2</sup> 2-form  $\Omega = -\frac{1}{2} dK = -\frac{1}{2} \partial_{[\mu} K_{\nu]} dx^\mu \wedge dx^\nu$ . Here we denoted  $K = K^a{}_a = K_\mu dx^\mu$ , and the traceless 1-form  $\bar{K}^a{}_b = K^a{}_b - \frac{1}{4} \delta_b^a K$ . It is easy to show that the vector field  $K_\mu$  decouples from the other fields and interacts with matter and with the metric similarly to the electromagnetic field. Furthermore, the homothetic curvature plays the role of the “Maxwell tensor”. Indeed, substituting (6.50) into (6.49) we recast the Yang-Mills part of the Lagrangian

<sup>1</sup>Note that  $K_{ab} = K_{\mu ab} dx^\mu$  where  $K_{\mu ab} = h_\alpha^a h_b^\beta \nabla_\mu g_{\alpha\beta}$ , cf. the definition of the tensor of nonmetricity (1.7). Furthermore,  $2\Gamma^{(ab)} = D\eta^{ab} = -\eta^{ac} \eta^{bd} D\eta_{cd}$ .

<sup>2</sup> $\Omega = \frac{1}{2} \Omega_{\mu\nu} dx^\mu \wedge dx^\nu$ , see (1.2) and (1.10) for the definition of the homothetic curvature tensor.

into

$$R^a{}_b \wedge *R^b{}_a = \frac{1}{4}\Omega \wedge *\Omega + \frac{1}{4}\widehat{D}\overline{K}^a{}_b \wedge *\widehat{D}\overline{K}^b{}_a \\ + \left( \widehat{R}^a{}_b + \frac{1}{4}\overline{K}^a{}_c \wedge \overline{K}^c{}_b \right) \wedge * \left( \widehat{R}^b{}_a + \frac{1}{4}\overline{K}^b{}_d \wedge \overline{K}^d{}_a \right),$$

while the curvature scalar  $R(\Gamma)$ , depends only on the first term from the right-hand side of (6.50), not containing  $K_\mu$ . The matter source of the quasi-Maxwell Weyl field is identified with the dilation current  $J^\mu$  which is the trace part of the hypermomentum (6.39). The latter is conserved due to the invariance of the  $\mathcal{L}_1$  Lagrangian (6.49) with respect to Einstein's  $\lambda$ -transformations [70].

The remaining irreducible non-Riemannian parts of  $GL(4, R)$ -connection  $\Gamma$  have the short-range behaviour due to the effective mass arising for them from the Hilbert-Einstein term in (6.49). One can show that in the weak field approximation for the Lagrangian (6.49), the translational field  $h_\mu^a$  has the standard Einstein dynamics, thus providing the correspondence with GR.

# 7

## Quantization of gravity

### 7.1. Methods of quantum gravity

Construction of the quantum theory of a physical system starts with the conversion of its degrees of freedom into operators on which the canonical commutation relations are then imposed. Obviously this should be preceded by the construction of the canonical formalism which was analyzed at the classical level in the previous chapters. The results of this analysis, caused by the special features of the canonical formalism in the curved spacetime and the degeneracy of the gravitational action, reveal a number of new features of the quantum theory.

A basic feature of gravity theory is that it is degenerate in virtue of its gauge invariance under local diffeomorphisms, so that we need a method of quantization of degenerate systems. As was shown in Chapter 2, the degenerate theory described by the canonical action (2.27) with the first-class constraints (2.28) after choosing the canonical gauges (2.32) reduces to the effective non-degenerate system with the action (2.40) in terms of independent physical degrees of freedom  $\Phi^* = (q^*, p^*)$  satisfying (2.39). The quantization of this theory means that  $q^*$  and  $p^*$  become operators in the Hilbert space of states and are subject to canonical commutation relations

$$\left[ \hat{\Phi}^*, \hat{\Phi}^{*'} \right] = i\hbar \{ \Phi^*, \Phi^{*'} \}^*, \quad (7.1)$$

where  $\{ , \}^*$  are the Poisson brackets calculated for the set of the physical phase variables  $\Phi^*$ . In what follows we consider the Planck constant  $\hbar$  to be one in universal units, but later will reinstate it explicitly when considering the semiclassical expansion in powers of  $\hbar$ .

The variables of the original total phase space  $\Phi = (q, p)$  also become operators as the operator functions of  $\hat{\Phi}^*$  (2.38). However, their commutation relations differ essentially from the “naive” canonical commutators defined relative to the Poisson brackets  $\{ , \}$  in  $(q, p)$ -space

$$\left[ \hat{\Phi}, \hat{\Phi}' \right] = i \{ \Phi(\Phi^*), \Phi'(\Phi^*) \}^* \neq i \{ \Phi, \Phi' \} \quad (7.2)$$

and essentially depend on a choice of a gauge in the complete set of constraints  $\psi^a = (\chi^\mu, H_\mu)$ .

Thus, within the framework of a certain set of gauges, the quantum system is described by the operators  $\hat{\Phi}^*$ , and its state is determined in the Schrödinger picture by the vector of the space of states  $|\Psi_{\text{phys}}(t)\rangle$ , evolving according to the Schrödinger equation with the physical Hamiltonian

$$i \frac{\partial |\Psi_{\text{phys}}(t)\rangle}{\partial t} = \hat{H}_{\text{phys}}(\hat{\Phi}^*) |\Psi_{\text{phys}}(t)\rangle. \quad (7.3)$$

When applied to the gravity, this approach is a direct realization of the quantization program in the ADM procedure. Solution of the constraints in any given gauge defines the chronologically ordered family of space-like hypersurfaces and the corresponding family of quantum states representing the quantum evolution.

However, this approach has serious disadvantages. Solution of the constraints is in general a complicated problem, the resulting physical Hamiltonian turns out to be non-local, and the whole method is not manifestly covariant. Moreover, a priori there is no guarantee that such quantization performed with different gauge conditions will give unitarily equivalent results. The first step to the understanding how this equivalence can be attained was done in [202] within the path integral formalism.

The time evolution operator for the Schrödinger equation (7.3) has a form of the path integral over the physical phase space [154, 155]:

$$U_{\text{phys}}(g_1, t_1 | g_0, t_0) = \int D[g^A, p_A] \exp i \int_{t_0}^{t_1} dt \left( p_A \dot{g}^A - H_{\text{phys}}(g^A, p_A) \right), \quad (7.4)$$

$$D[g^A, p_A] = \prod_{t,A} dg^A(t) \prod_{t',B} dp_B(t'), \quad (7.5)$$

where the integration is done with respect to the Liouville measure<sup>1</sup> and with the boundary conditions for the phase space coordinate  $g^A(t)$  at the initial  $t_0$  and final  $t_1$  moments of time

$$g^A(t_0) = g_0^A, \quad g^A(t_1) = g_1^A. \quad (7.6)$$

---

<sup>1</sup>Here and in what follows we will disregard trivial purely numerical factors like  $\prod_t (1/2\pi)$  in the path integral measure, in the Fourier representation of relevant functional delta functions, etc.

One should use the relation between the phase space integration measure on the physical subspace of  $g^A$  and  $p_A$  and the delta-function type measure on the original phase space of  $q^i$  and  $p_i$

$$\prod_A dq^A dp_A = \prod_i dq^i dp_i \delta(\chi(q, p)) \delta(H(q, p)) J(q, p). \quad (7.7)$$

Here the delta functions of gauge conditions and constraints have a support on the subspace where they are vanishing,

$$\delta(\chi(q, p)) \equiv \prod_\mu \delta(\chi^\mu(q, p)), \quad (7.8)$$

$$\delta(H(q, p)) \equiv \prod_\mu \delta(H_\mu(q, p)) = \int \left( \prod_\mu dN^\mu \right) e^{iN^\mu H_\mu(q, p)}, \quad (7.9)$$

and the factor  $J(q, p)$  is the determinant of the functional matrix of the canonical Faddeev-Popov operator  $J_\nu^\mu(q, p)$  given by the Poisson brackets of the set of gauge conditions and constraints (cf. equation (2.33) or equation (3.7) for the case of a special choice of canonical coordinates  $\Phi = (X^\mu, \Pi_\mu; g^A, p_A)$ )

$$J(q, p) = \det J_\nu^\mu(q, p), \quad J_\nu^\mu(q, p) = \{\chi^\mu, H_\nu\}. \quad (7.10)$$

Therefore, making according to (7.7) the change of integration variables in (7.3) from  $(g^A, p_A)$  to  $(q, p)$  and using the relation (2.39) we get

$$U_{\text{phys}} \sim \int D[q, p] DN \delta[\chi(q, p)] J[q, p] \exp i \int_{t_0}^{t_1} dt (p_i \dot{q}^i - H_0 - N^\mu H_\mu), \quad (7.11)$$

where the linear combination of first class constraints in the exponentiated canonical action was achieved due to the integral representation of their delta function (7.9). Here the *functional* delta function  $\delta[\chi(q, p)]$  and the functional determinant  $J[q, p]$  obviously represent products over time moments of their local analogues,

$$\delta[\chi(q, p)] = \prod_t \delta(\chi(q(t), p(t))); \quad J[q, p] = \prod_t J(q(t), p(t)), \quad (7.12)$$

while the path integral measure  $D[q, p]$  is of course similar to that of the physical phase space (7.5),

$$D[q, p] = \prod_{t,i} dg^i(t) \prod_{t',k} dp_k(t'), \quad DN = \prod_{t',\mu} dN^\mu(t'). \quad (7.13)$$

The canonical path integral representation of the unitary evolution operator (7.11) allows one to perform the gauge dependence analysis, because the variation of the gauge in this expression can be achieved by a canonical transformation of integration variables [202], and thus its gauge independence can

formally be proven. This gives a first hint on the unitary equivalence of quantization schemes in different sets of gauge conditions. Moreover, the contribution of the functional determinant can be represented as a functional integral over the anticommuting (Grassman) variables – the Faddeev-Popov ghost fields  $C^\nu(t)$  and  $\bar{C}_\mu(t)$  [203],

$$J[q, p] = \int D\bar{C}DC \exp \left\{ i \int dt \bar{C}_\mu(t) J_\nu^\mu(q(t), p(t)) C^\nu(t) \right\}, \quad (7.14)$$

so that the total action becomes local in spacetime, and the path integral becomes manageable by a standard technique of renormalization of ultraviolet divergences. Integration over the canonical momenta converts it to the Lagrangian form which within the generalized set of special (relativistic) gauges becomes manifestly covariant and admits covariant renormalization.

All these advantages of the path integral representation remain, however, incomplete because the change of integration variables  $(g^A, p_A) \rightarrow (q^i, p_i)$  is not consistent at the end points of the time interval  $[t_0, t_1]$ . This happens in virtue of the boundary conditions (7.6) and unequal footing of the coordinates and momenta at these points – while the coordinates are kept fixed, the momenta are being integrated over. For these reasons strict equality in (7.11) was replaced by the similarity sign. In scattering theory applications this difficulty is easily circumvented due to the simplicity of the linear regime for asymptotic states at  $t_0 \rightarrow -\infty$  and  $t_1 \rightarrow +\infty$  ( $S$ -matrix theory), but for finite times and nontrivial quantum states this issue represents a real problem. And this problem becomes critically important in the cosmological context because the initial state of the very early quantum Universe cannot be formulated in terms of asymptotic states of scattering theory.

This problem, in particular, leads to the question of how the boundary conditions on integration histories in the right hand side of (7.11) should be associated with the arguments of the kernel  $U_{\text{phys}}$  on the left and side, how the composition law for the path integrals over  $(q, p)$  in the time intervals  $[t_0, t_1]$  and  $[t_1, t_2]$  should look like, or what will be the analogue of the unitary propagation law for quantum states in the physical (ADM) sector:

$$\Psi_{\text{phys}}(g_1, t_1) = \int dg_0 U_{\text{phys}}(g_1, t_1 | g_0, t_0) \Psi_{\text{phys}}(g_0, t_0), \quad (7.15)$$

not to mention how the physical wave functions  $\Psi_{\text{phys}}(g, t)$  should be associated with the quantum states  $\Psi(q, t)$  in the coordinate representation of the operators  $(\hat{q}, \hat{p})$ . The answers to this set of questions will be partially given in the rest of this chapter, but before that we will briefly consider the ADM quantization method in terms of the physical sector of phase space variables  $\Phi^*$ .

## 7.2. Arnowitt-Deser-Misner quantization method

In the ADM formalism (see Chapter 3), the physical degrees of freedom of arbitrary gravitating system are the pairs of the phase variables  $\Phi^A = (g^A, p_A)$  (3.4), which in accordance with the previous section become operators satisfying the canonical commutation relations (7.1). The state of the system in the set of the gauges (3.11) is the quantum state on one of the space-like hypersurfaces defined in the four-dimensional manifold with the help of the same gauge conditions. This state is described by the vector  $|\Psi_{\text{phys}}(t)\rangle$  which satisfies the Schrödinger equation

$$i \frac{\partial |\Psi_{\text{phys}}(t)\rangle}{\partial t} = \hat{H}_{\text{phys}}(\hat{g}^A, \hat{p}_A) |\Psi_{\text{phys}}(t)\rangle, \quad (7.16)$$

where the Hamiltonian is given by the formula (3.14).

With the solution of this equation, one can find the quantum averages of all phase variables  $\hat{\Phi} = (\hat{g}_{ab}, \hat{p}^{ab}; \hat{\varphi}, \hat{p})$  of the system, as well as of the lapse and shift functions. Indeed, according to (3.5), (3.12) and (3.15), we have

$$\begin{aligned} \langle \hat{\Phi}(t) \rangle &= \langle \Psi_{\text{phys}}(t) | \Phi[f^\mu, -P_\mu[f; \hat{g}^A, \hat{p}_A]; \hat{g}^A, \hat{p}_A] | \Psi_{\text{phys}}(t) \rangle, \quad (7.17) \\ \langle \hat{N}^\mu(\mathbf{x}) \rangle(t) &= \langle \Psi_{\text{phys}}(t) | \int d^3y \hat{J}^{-1(\mu)}_{(\nu)}(\mathbf{x}, \mathbf{y}) | \Psi_{\text{phys}}(t) \rangle \dot{f}^\nu(t, \mathbf{y}), \end{aligned}$$

where in the last equation the operator nature of  $\hat{J}^{-1(\mu)}_{(\nu)}(\mathbf{x}, \mathbf{y})$  implies that it is taken as a function of operators of physical variables and  $X^\mu = f^\mu(t, \mathbf{x})$  and  $\Pi_\mu = -P_\mu[f^\nu; \hat{g}^A, \hat{p}_A]$ . The proper time of the observer, whose world line is described by the equation  $\mathbf{x} = \text{const}$ , becomes the physical observable. Its average is given by the expression

$$\langle \tau \rangle = \int dt \langle N \sqrt{1 - N_a N^a / N^2} \rangle. \quad (7.18)$$

Similarly, the spatial intervals become the quantum observables. In accordance with the operator nature of the lapse and shift functions in (7.17), the position of a space-like hypersurface itself is an observable. This fact reflects special features of the quantization of the gravitational field, whose universal interaction determines the properties of space and time, and thereby affects the observer who cannot be shielded from this influence that has a quantum nature.

### *Example of ADM quantization: Friedmann universe with a scalar field*

The dynamics of the closed homogeneous universe filled with the scalar matter will be briefly considered here as a demonstration of the ADM quantization formalism. The subject of quantization will not only be the matter field but

also the gravitational configuration of the Friedman model [164] described by the metric (3.39).

The physical setup we will consider has basically a toy-model nature designed more for the demonstration of the limitations of the ADM approach, rather than for revealing some interesting new quantum physics undoubtedly inherent to the early quantum cosmology. Mainly we will consider the situation when the dynamics of the model is only slightly different from the evolution of the classical Universe. The qualitative behaviour of closed Friedmann models in the Einstein theory is the picture of the world that expands up to some maximal cosmological radius from the initial singularity, and then contracts back to the singular state. The expansion process is characterized by the suppression of the originally present inhomogeneous gravitational modes. Conversely, when contraction to the singularity, the Universe becomes essentially inhomogeneous and anisotropic. If we restrict ourselves to the dynamical stage ranging from the maximal radius to the domains which are sufficiently far from the singularity, one can take the homogeneous and isotropic model as an approximation of the closed Universe.

We can assume that the homogeneous gravitational field at the maximal expansion is generated by the large number of randomly moving particles that fill uniformly the entire volume of the Universe being in a thermodynamic equilibrium. To describe particles, it is necessary to introduce the inhomogeneous matter field, whose separate oscillatory modes correspond to the particles of different energies. Thus, the consistency condition of the model reduces not to the requirement of the homogeneity of the matter field, as it is done in the majority of the toy models [178] which claim to give a self-consistent dynamics of the homogeneous matter and gravitational fields, but rather to the requirement of constancy in the three-dimensional space of such characteristics as the energy density, the number of particles, etc.

The action of the massless scalar field  $\varphi$  reads

$$S_m = -\frac{1}{2} \int d^4x g^{1/2} g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi. \quad (7.19)$$

In view of the form of the metric (3.39), we derive the Lagrange function in terms of the dimensionless scalar field  $\varphi = l_0 \varphi$  and the dimensionless cosmological radius  $a$  (3.41)

$$L_m = \frac{\varepsilon_0}{2} \int d^3x \sqrt{\gamma} \left\{ \frac{\chi^2}{N} l_0^2 a^3 + a \varphi \Delta \varphi \right\}, \quad (7.20)$$

where the covariant Laplacian  $\Delta$  is defined by the constant three-dimensional metric  $\gamma_{ik}$ . Decomposing the field  $\varphi(\mathbf{x}, t)$  with respect to the orthonormal system of eigenfunctions  $Z_n(\mathbf{x})$  similarly to (4.56),  $\varphi(\mathbf{x}, t) = \sum_n \varphi'_n(t) Z_n(\mathbf{x})$ , we obtain the Lagrange function (7.20) in terms of discrete set of degrees of freedom of the scalar field

$$L_m = \frac{\varepsilon_0}{2} \sum_n \left\{ \frac{\dot{\varphi}'_n{}^2}{N} l_0^2 a^3 - N a \omega_n^2 \varphi'_n{}^2 \right\}. \quad (7.21)$$

The superhamiltonian which corresponds (7.21), reads

$$\int d^3x \mathcal{H}_\varphi = \frac{\varepsilon_0}{2a} \sum_n \left\{ \frac{\mathcal{P}_n^2}{a^2} + \omega_n^2 a^2 \varphi_n'^2 \right\}, \quad (7.22)$$

where  $\mathcal{P}_n$  are momenta conjugated to  $\varphi_n'$ .

We take the Hamiltonian constraint (3.42), or (2.125) (rather, its integral analogue) in the form

$$H_\perp = \frac{\varepsilon_0}{2} \left\{ -\frac{1}{2}(p^2 + a^2) + \frac{1}{2} \sum_n \left[ \frac{\mathcal{P}_n^2}{a^2} + \omega_n^2 a^2 \varphi_n'^2 \right] \right\} \quad (7.23)$$

and perform the canonical transformation of the general ADM procedure (3.4) in two stages. At first, we make the transformation

$$\begin{aligned} (p, a; \mathcal{P}_n, \varphi_n') &\longrightarrow (p', a'; p_n, \varphi_n), \\ p &= p' + \frac{V}{a'}, \quad a = a', \quad V \equiv \sum_n p_n \varphi_n, \\ \mathcal{P}_n &= a' p_n, \quad \varphi_n' = \varphi_n / a', \end{aligned} \quad (7.24)$$

and then similarly to (3.48)-(3.49), we substitute  $(a', p')$  by the new phase variables  $T$  and  $\Pi$ :

$$a' = \sqrt{-2\Pi} \cos T, \quad p' = \sqrt{-2\Pi} \sin T. \quad (7.25)$$

As a result, the constraint (7.23) takes the following form:

$$H_\perp = \frac{\varepsilon_0}{\sqrt{-2\Pi} \cos T} \left\{ \Pi + H + \frac{V^2}{4\Pi \cos^2 T} \right\}, \quad (7.26)$$

where

$$H = \frac{1}{2} \sum_n (p_n^2 + \omega_n^2 \varphi_n^2 - 2 \varphi_n p_n \tan T). \quad (7.27)$$

In accordance with the general ADM procedure, let us impose the gauge condition

$$T = t \quad (7.28)$$

and solve the constraint equation (7.26)  $H_\perp = 0$  with respect to  $\Pi$  approximately in the region of  $H \gg 1$ . Then with an accuracy  $O(H^{-1})$ , we obtain the action

$$S = \int dt \left\{ \sum_n p_n \dot{\varphi}_n - H(t) + \frac{V^2}{4H(t) \cos^2 t} + \dots \right\} \quad (7.29)$$

in terms of the physical degrees of freedom  $(\varphi_n, p_n)$ .

From now on, we will consider such a range of the parameter  $t$  and such states for which  $H(t) \cos^2 t \gg 1$ , therefore, we can neglect the last term in (7.29).

Consequently, the physical Hamiltonian of the problem may be approximated by

$$H(t) = \frac{1}{2} \sum_n (p_n^2 + \omega_n^2 \varphi_n^2 - 2 p_n \varphi_n \tan t). \quad (7.30)$$

According to (3.54), (7.26), and (7.25) with an accuracy of  $O(H^{-1})$ , the expressions for the lapse function and scale factor read

$$N = l_0(\sqrt{2H(t)} + O(H^{-1}(t))) \cos t, \quad a = (\sqrt{2H(t)} + O(H^{-1}(t))) \cos t, \quad (7.31)$$

since from the solution of the constraint equation we have  $-H = H(t) + O(H^{-1})$ . We see that in this approximation the gauge (7.28) is equivalent to the choice of the coordinate condition (3.46),  $N = R$ . The condition  $H(t) \cos^2 t \gg 1$ , imposed on the range of applicability of our approximation, is equivalent to the restriction

$$R \gg l_0, \quad (7.32)$$

i.e., to the values of the cosmological radius which are essentially greater than the Planck length.

We will quantize the system described by the Hamiltonian (7.30) in the Heisenberg representation, and as a first step we write the Heisenberg equation for  $\varphi_n$  that follows from (7.30):

$$\ddot{\varphi}_n + \Omega_n^2 \varphi_n = 0, \quad (7.33)$$

where  $\Omega_n^2 = \omega_n^2 + 1$ . As in Sec. 4.4., in order to introduce the particle interpretation of the system, we decompose the Heisenberg operator  $\varphi_n(t)$  with respect to the two complex-conjugated solutions of this equation  $u_n$  and  $u_n^*$ :

$$\varphi_n(t) = a_n u_n(t) + a_n^* u_n^*(t). \quad (7.34)$$

The basis functions  $u_n$  and  $u_n^*$  are chosen so that they satisfy the normalization conditions

$$u_n \dot{u}_n^* - \dot{u}_n u_n^* = -i,$$

therefore, for the creation and annihilation operators  $a_n^*$  and  $a_n$  the commutator reads  $[a_n, a_m^*] = \delta_{nm}$ .

Further analysis of the particle interpretation of the system can be done with the help of the standard procedure of Hamiltonian's diagonalization [61] described in Sec. 4.4.. Diagonalization of the Hamiltonian at the moment of the maximal expansion of the Universe  $t = 0$  takes place in terms of basis functions of the equation (7.33)

$$u_n(t) = \frac{1}{\sqrt{2\Omega_n}} \left\{ \sqrt{\frac{\Omega_n}{\omega_n}} \cos \Omega_n t - i \sqrt{\frac{\omega_n}{\Omega_n}} \sin \Omega_n t \right\}, \quad (7.35)$$

which are nontrivial because the frequency  $\omega_n$  in the Hamiltonian and the frequency in the Heisenberg equation of motion  $\Omega_n$  do not coincide (for  $\Omega_n = \omega_n$

these basis functions would reduce to the standard positive frequency ones  $\exp(-i\omega_n t)/\sqrt{2\omega_n}$ . For  $n = 0$  with  $\omega_0 = 0$  the basis function is not well defined, but this mode does not have an oscillator nature and should be quantized in the coordinate (or momentum) representation which we will not consider in much detail<sup>2</sup>. The modes with  $n > 0$  have a particle interpretation in terms of the creation and annihilation operators associated with the modes which would diagonalize the Hamiltonian at later moments of time and which do not coincide with (7.35).

In our toy model setup the initial state of the system is prescribed at the moment  $t = 0$  of the maximal expansion of the Universe when the non-stationarity of the metric and of spacetime curvature is minimal. The proper cosmological time that the Universe spends in the region of the maximal expansion  $\tau = \int dt N = \int dt R(t)$  is large, therefore, despite the fact that we neglected the interaction of particles from the very beginning, this interaction is sufficient for the collision relaxation to bring the system of particles in the Universe into the state of the thermal equilibrium. Therefore, it is reasonable to define the initial state as the equilibrium or thermal density matrix  $\hat{\rho}$  at some temperature.

The average value of any physical observable  $\hat{\mathcal{O}}$  may be defined by formula

$$\langle \hat{\mathcal{O}}(t) \rangle = \text{tr}(\hat{\rho} \mathcal{O}(t)) = \langle \overline{\mathcal{O}(t)} \rangle, \quad (7.36)$$

where  $\langle \mathcal{O}(t) \rangle$  should be understood as the quantum averaging of the Heisenberg operator  $\mathcal{O}(t)$  over a pure multiparticle state from the space of occupation numbers, and the bar means the statistical averaging with respect to  $\hat{\rho}$ . The details of such calculations for a thermal density matrix can be found in [71, 168].

Here we want to stress that in the approximation of large  $H(t)$  the calculation of the quantum average of the scale factor gives

$$\langle \hat{a}(t) \rangle = \langle \overline{\sqrt{2H(t)} + O(H^{-1}(t))} \rangle \cos t = \sqrt{2\mathcal{E}(t)} \cos t \{1 + O(\mathcal{E}^{-1}(t))\}, \quad (7.37)$$

where  $\mathcal{E}(t)$  is the evolving in time quantum average of the Hamiltonian

$$\mathcal{E}(t) = \langle \overline{\hat{H}(t)} \rangle. \quad (7.38)$$

This quantum average is ultraviolet divergent, and because of the essential lack of covariance and nonlinearity of the physical observable  $\hat{a}$  it is hopeless to have a renormalization of the arising divergences by local covariant counterterms in the original action of the theory. Therefore, the obtained result makes sense only in the situation when the contribution of large occupation numbers in  $\mathcal{E}(t)$  is a priori dominating the vacuum polarization contribution subject to UV renormalization. This setup is not applicable in the physically interesting

---

<sup>2</sup>One can check that the dynamics of the quasi-Gaussian wave packet in the coordinate representation of  $\varphi_0$  with the growth of time will be qualitatively described by a quantum spreading of this packet.

case of the early quantum cosmology when the vacuum polarization effects and nontrivial scaling behavior play the dominant role. ADM quantization in view of its nonlocality, nonlinearity and lack of manifest covariance (which can hardly be restored even at the implicit level) is incapable of solving this important class of quantum problems. For this reason, we go over to another quantization approach to gravity theory.

### 7.3. Dirac-Wheeler-DeWitt quantum geometrodynamics

The Einstein-Hamilton-Jacobi theory provides an intermediate link between the classical theory of gravity and Dirac-Wheeler-DeWitt quantization, historically called a quantum geometrodynamics, and we begin the review of this approach with its analysis. For simplicity of presentation, we restrict ourselves with the case of the pure gravitational field and consider only the closed cosmology systems with  $H_0 = 0$ .

#### *Einstein-Hamilton-Jacobi theory as a semiclassical approximation to quantum geometrodynamics*

The Hamilton-Jacobi equations (2.44), (2.45) for spatially closed cosmological systems with  $q^i = g_{ab}(\mathbf{x})$  and  $\partial/\partial q^i = \delta/\delta g_{ab}(\mathbf{x})$ , in accordance with (2.94) and (2.95), take the following form:

$$G_{abcd} \frac{\delta S}{\delta g_{ab}} \frac{\delta S}{\delta g_{cd}} - g^{1/2} {}^3R = 0, \quad (7.39)$$

$$-2g_{ac} \left[ \frac{\delta S}{\delta g_{cb}} \right]_{|b} = 0, \quad (7.40)$$

$$\frac{\partial S}{\partial t} = 0. \quad (7.41)$$

The last equation means that the Hamilton-Jacobi function  $S$  of the gravitational field does not depend on time  $t$ . This is a manifestation of the “frozen” formalism for closed cosmological systems, as described in Chapter 3. Time independence of  $S$  means that formally the time evolution can be incorporated in the system only with the aid of the gauge conditions (2.48) that by construction explicitly depend on time.

Thus,  $S$  is a functional of the three-dimensional metric and of the  $6 - 4 = 2$  functions  $\alpha^A(\mathbf{x})$  which represent the independent integration constants in the complete integral of the system of equations (7.39), (7.40).

The equations (7.40) have a simple geometrical meaning since they reflect the invariance of the functional  $S[g_{ab}(\mathbf{x}), \alpha^A(\mathbf{x})]$  with respect to the coordinate

transformations of the metric

$$\Delta^f g_{ab}(\mathbf{x}) = 2f_{(a|b)}(\mathbf{x}), \quad (7.42)$$

$$\Delta^f S = \int d^3x \frac{\delta S}{\delta g_{ab}(\mathbf{x})} \Delta^f g_{ab}(\mathbf{x}) = -2 \int d^3x \left[ \frac{\delta S}{\delta g_{ab}(\mathbf{x})} \right]_{|b} f_a(\mathbf{x}) \equiv 0. \quad (7.43)$$

Therefore,  $S$  is actually the functional not of the three-dimensional metric, but of three-dimensional geometry that encompasses the equivalence class of the three-dimensional metrics connected by the coordinate transformations.

The equation (7.39) is called the Einstein-Hamilton-Jacobi equation (EHJ) and it represents the basic equation of the classical geometrodynamics. Actually, this equation contains the entire dynamics of the classical theory of gravity. In accordance with the results of Sec. 2.1., the evolution of a gravitating system is restored as a solution of systems (2.47), (2.48), where the gauges are chosen to be explicitly time-dependent.

The Einstein-Hamilton-Jacobi theory is a semiclassical approximation to the quantum theory. However, the gravitation is a degenerate system with constraints and this approximation has the special gauge-theoretic features which are follows.

In the semiclassical approximation, the amplitude of the state of the quantum system is described in the coordinate representation by the superposition of the functions of the form  $\exp \{iS(q, \alpha)\}$  with different values of parameters  $\alpha$ , and with the corresponding amplitudes  $P(\alpha)$ . This superposition forms the wave-package

$$\Psi = \int d\alpha P(\alpha) e^{iS(q, \alpha)}. \quad (7.44)$$

The center of gravity of such package or the point in the coordinate space describing the classical system position is determined as a stationary point of the integral (7.44) with respect to the variables  $\alpha$ ,

$$\frac{\partial S}{\partial \alpha^A} = \beta_A, \quad (7.45)$$

where the parameters  $\beta_A = -\frac{1}{i} \partial \ln P(\alpha) / \partial \alpha^A$  depend on the form of the amplitude  $A(\alpha)$ . In non-degenerate systems, the equations (7.45) contain all the necessary information to find the classical evolution. In systems with constraints, the equations (7.45) or (2.47) are not sufficient, since they define the surface in the space of the phase coordinates that consists of a set of the classical trajectories describing the same physical state and connected by the transformations of the “group” of invariance of the action (2.30). To select a single physical representative, it is necessary to impose the gauge (2.32), i.e., to introduce another surface in the space of the variables  $q$ , described by the equation (2.48), that have a single intersection point with the initial one.

The space of three-dimensional geometries, where the action  $S$  is determined as a solution of EHJ equation (7.40), is usually called a superspace<sup>3</sup> [167]. However, since we can operate in a constructive way only in terms of the three-dimensional metrics, we use the term “superspace” to call the manifold of the three-dimensional metrics  $g_{ab}(\mathbf{x})$ , where the action  $S$  satisfying equations (7.40)-(7.42) is determined.

Now let us proceed to the construction of the exact quantum theory, whose quasi-classical approximation would give the EHJ theory.

### *Dirac quantization method and the Wheeler-DeWitt equation*

An alternative to the ADM quantization approach is the Dirac quantization method for theories subject to constraints [40]. The operators of the original phase space  $\hat{\Phi} \equiv (\hat{q}, \hat{p})$ , in contrast to the previous ADM method, are treated as independent and satisfying the usual canonical commutation relations

$$[\hat{\Phi}, \hat{\Phi}'] = i\{\Phi, \Phi'\}. \quad (7.46)$$

These operators are defined on a maximally wide space of vectors. However, the vectors of the physically admissible states may not be arbitrary, but only those that satisfy the quantum constraints as the equations

$$\hat{H}_\mu(\hat{q}, \hat{p}) |\Psi\rangle = 0 \quad (7.47)$$

and the Schrödinger equation

$$\frac{\partial |\Psi\rangle}{\partial t} = \hat{H}_0(\hat{q}, \hat{p}) |\Psi\rangle. \quad (7.48)$$

The latter reduces to the independence  $|\Psi\rangle$  on  $t$  for the special case of a closed world with  $H_0 \equiv 0$ .

The consistency of the system of equations (7.47), (7.48) is guaranteed by the involution relations (2.29), provided that one choose such order of operators in the constraints  $\hat{H}_\mu$  and in the Hamiltonian  $\hat{H}_0$ , that the Poisson brackets in (2.29) are mapped into the quantum commutators constructed on the basis of (7.46):

$$[\hat{H}_\mu, \hat{H}_\nu] = \hat{U}_{\mu\nu}^\alpha \hat{H}_\alpha, \quad [\hat{H}_0, \hat{H}_\mu] = \hat{V}_\mu^\nu \hat{H}_\nu, \quad (7.49)$$

with the *operators* of structure functions  $\hat{U}_{\mu\nu}^\alpha$  and  $\hat{V}_\mu^\nu$  standing to the left of the operators of constraints.

In the coordinate representation of the commutation relations (7.46), the momenta  $\hat{p} = \partial/i\partial q$ , the coordinates  $\hat{q}$  are  $c$ -numbers, and  $|\Psi\rangle = \Psi(q) = \Psi[g_{ab}(\mathbf{x})]$ , so that the equation (7.47) takes the form

$$H_\mu\left(q, \frac{\partial}{i\partial q}\right) \Psi(q) = 0, \quad (7.50)$$

---

<sup>3</sup>It should not be confused with the superspace in the modern supersymmetric theories. These are different notions.

with some particular choice of the operator realization  $H_\mu(q, \partial/i\partial q)$  of  $\hat{H}_\mu$ , which includes a particular operator ordering of  $q$  and  $\hat{p}$  and possible quantum correction terms  $O(\hbar)$ .<sup>4</sup>

Thus, in quantum geometrodynamics with  $\partial/i\partial q \equiv \delta/i\delta g_{ab}(\mathbf{x})$  at least naively we have

$$\left\{ -G_{abcd} \frac{\delta}{\delta g_{ab}} \frac{\delta}{\delta g_{cd}} - g^{1/2} {}^3R \right\} \Psi[g_{ab}] = 0, \quad (7.51)$$

$$-2g_{ac} \left[ \frac{\delta \Psi[g_{ab}]}{\delta g_{cb}} \right]_{|b} = 0. \quad (7.52)$$

The first of these equations is a widely acclaimed Wheeler-DeWitt equation [53] which, of course, represents an infinite set of equations – one per each spatial point. Its operator realization is thus far purely symbolic and will be discussed later. On the contrary, the operator ordering in (7.52) has a sufficiently strong geometrical ground because the equation (7.52), similarly to (7.43), incorporates the local coordinate invariance of  $\Psi[g_{ab}(\mathbf{x})]$ , which rather firmly fixes the operator realization of quantum momentum constraints.

The Wheeler-DeWitt equation is a basic equation of quantum geometrodynamics. Obviously, the quasi-classical approximation leads to its solution of the form

$$\Psi = e^{iS},$$

where  $S$  satisfies the equations (7.40) and (7.42).

Thus, comparing this approach with the ADM quantization in reduced phase space, we find the two different situations. In quantum geometrodynamics – the Dirac quantization scheme – the operators  $(\hat{q}, \hat{p})$  are independent but the physical states are subject to the constraints (7.47). In the ADM (or reduced phase space) quantization the operators  $(q, p)$  identically satisfy the constraints but the states are arbitrary. And no unitary equivalence exists between these two approaches, because the unitary equivalent representations have the same commutation relations, whereas the commutation relations (7.46) are fundamentally different from (7.2).

Another difficulty is that the Dirac quantization scheme is intrinsically incomplete, because a priori it lacks the algorithm for the physical inner product of quantum states that would give quantum expectation values and probabilities. The usual  $L^2$  inner product

$$\langle \Psi' | \Psi \rangle = \int dq \Psi'^*(q) \Psi(q) \quad (7.53)$$

does not make sense, because the physical states  $\Psi(q)$ , which are annihilated by constraints, have a distributional nature  $\sim \delta(\hat{H}_\mu)$  and make this integral

---

<sup>4</sup>The inclusion of these terms will require to reinstate explicitly the Planck constant  $\hbar$  which will be done later.

divergent,  $[\delta(\hat{H})]^2 \sim \infty$ . This inner product should have a nontrivial integration measure restricting the range of integration over  $q$  to a certain subspace corresponding to a given time slice of the whole spacetime (as long as physical time is hidden among the variables  $q$ ). The attempt to interpret as the physical inner product the flux of the conserved current of the Wheeler-DeWitt equation (7.51) [53, 165, 182] – the analogue of the conserved current of the second-order Klein-Gordon equation,

$$j_{ab}(\mathbf{x}) = G_{abcd} \left\{ \Psi^* \frac{\delta\Psi}{i\delta g_{cd}(\mathbf{x})} - \frac{\delta\Psi^*}{i\delta g_{cd}(\mathbf{x})} \Psi \right\}, \quad (7.54)$$

requires justification from first principles of quantization. Moreover, it demands specification of the surface in the space of 3-metrics  $q = g_{ab}(\mathbf{x})$  through which the flux of this current is running, this specification establishing the above mentioned association with the time slice of spacetime.

Consistency between the ADM quantization and quantum geometrodynamics as the Dirac quantization of gravity theory was attained in the series of works [204, 205, 206, 207, 208, 209, 210] where the questions of the above type were clarified and checked at least in the first nontrivial order of semiclassical expansion – one-loop approximation. It turned out that direct unitary equivalence between these quantization methods is impossible and can only be achieved by embedding the Dirac scheme into a wider quantization framework. This framework is the Batalin-Fradkin-Vilkovisky (BFV) operator quantization of constrained systems [213, 46, 214, 215, 216, 217, 218, 219, 220] stemming from the BRST method<sup>5</sup> [211, 212].

## 7.4. Dirac quantization from Batalin-Fradkin-Vilkovisky formalism

Here we give a brief overview of the BRST/BFV formalism in gauge systems with first class constraints [211, 212, 213, 46, 214, 215, 217, 218, 219, 220]. Within the technique of the relativistic phase space of gauge and ghost fields [213, 46] we build the operator of the unitary evolution in their representation space and formulate the BRST invariant physical states. A special Batalin-Marnelius (BM) procedure of gauging out the BRST symmetry in the subspace of these states [221, 222, 223] leads to a well-defined physical inner product in the space of BRST singlets [204, 206, 208]. Then, the Dirac quantization scheme turns out to be a special truncation of the BFV formalism [206, 208, 210] – a particular realization of this BM gauge fixing procedure – which allows one to construct a path integral representation for the solution of the Wheeler-DeWitt equations. In the subsequent section these results are explicitly checked in the

---

<sup>5</sup>BRST refers to Becchi, Rouet, Stora, and Tyutin [211, 212].

first one-loop order of semiclassical expansion which, in particular, allows one to formulate a unitary map from the Wheeler-DeWitt (Dirac) wave functions  $\Psi(q)$  to the physical states  $\Psi_{\text{phys}}(g)$  of the reduced phase space quantization.

### *BFV formalism*

We begin by reminding the reader that the ADM canonical formalism incorporates splitting the full configuration space of  $g_{\mu\nu}$  into phase space coordinates  $q^i = g_{ab}(\mathbf{x})$ , and non-dynamical Lagrange multipliers  $N^\mu = (N^\perp(\mathbf{x}), N^a(\mathbf{x}))$  in the canonical action (2.27) or (2.124). The momenta  $p_i$  are conjugated to  $q^i$ , whereas the variables  $N^\mu$  do not have conjugated momenta. In what follows we will use condensed canonical notations when the field labels carry together with discrete indices also spatial coordinates, and contraction of indices  $i$  or  $\mu$  implies spatial (but not time) integration. In open models with asymptotically flat (or other boundaries like horizons, etc.)  $H_0(q, p)$  represents the relevant surface integral specified by boundary conditions which are the part of physical setting for the gravitational system. Below we consider the case of spatially closed cosmology with  $H_0(q, p) = 0$ .

The diffeomorphism invariance of the theory has a manifestation in Poisson brackets algebra of constraints (2.29) with the structure functions  $U_{\mu\nu}^\lambda = U_{\mu\nu}^\lambda(q)$  which depend on phase-space coordinates  $q$ . At the quantum level the classical constraints take the form of equations on physical quantum states (7.47). In the functional coordinate representation of quantum gravity they form the system of the Wheeler-DeWitt equations on the wave function  $\langle q | \Psi \rangle = \Psi(q)$ . Their consistency requires the commutator algebra (7.49) to hold with the operator structure functions  $\hat{U}_{\mu\nu}^\lambda$  standing to the left of  $\hat{H}_\lambda$ .

This BFV quantization generalizes this Dirac quantization scheme by extending the representation space of the original phase space variables  $q^i, p_i$  to that of the relativistic phase space variables. The latter include together with  $q^i, p_i$  the canonically conjugated pairs of Lagrange multipliers and their momenta  $p_\mu$  and canonical pairs of Grassman (fermionic) ghosts  $C^\mu, \mathcal{P}_\mu$  and anti-ghosts  $\bar{C}_\mu, \bar{\mathcal{P}}^\mu$ ,

$$Q^I, P_I = q^i, p_i; N^\mu, p_\mu; C^\mu, \mathcal{P}_\mu; \bar{C}_\mu, \bar{\mathcal{P}}^\mu, \quad (7.55)$$

$$[Q^I, P_J] = i \delta_J^I. \quad (7.56)$$

Here  $[A, B]$  denotes a supercommutator taking into account the Grassman parity  $n$  of  $A$  and  $B$ ,  $[A, B]_\pm = AB - (-1)^{n(A)n(B)}BA$ . In gravity theory with bosonic matter fields  $n(q) = n(N) = 0$  and  $n(C) = n(\mathcal{P}) = n(\bar{C}) = n(\bar{\mathcal{P}}) = 1$ . The canonical commutation relations (7.56) represent the quantum version of classical Poisson superbrackets commutators  $\{Q^I, P_J\} = i \delta_J^I$  (we use units with  $\hbar = 1$ ). Ghost variables have Hermiticity properties compatible with these commutation relations

$$C^{\mu\dagger} = C^\mu, \quad \mathcal{P}_\mu^\dagger = -\mathcal{P}_\mu, \quad \bar{C}_\mu^\dagger = -\bar{C}_\mu, \quad \bar{\mathcal{P}}^{\mu\dagger} = \bar{\mathcal{P}}^\mu. \quad (7.57)$$

In what follows we will regularly omit the hat notation for operators (7.55) and only use it in case when they have to be distinguished from their  $c$ -number eigenvalues. The hat notations will as a rule be used for composite operators. State vectors in the representation space of all relativistic variables will be denoted by double ket notations  $||\Psi\rangle\rangle$  (contrary to vectors  $|\Psi\rangle$  in the representation space of the original operators  $(\hat{q}^i, \hat{p}_i)$ ). The coordinate representation will be introduced as follows:

$$||Q\rangle\rangle \equiv ||q, N, C, \bar{C}\rangle\rangle, \hat{Q}^I ||Q\rangle\rangle = Q^I ||Q\rangle\rangle, \Psi(Q) = \langle\langle Q ||\Psi\rangle\rangle, \quad (7.58)$$

$$\langle\langle \Psi_1 ||\Psi_2\rangle\rangle = \int dQ \Psi_1^*(Q) \Psi_2(Q), \quad (7.59)$$

where the BRST inner product is defined in the sense of Berezin integration over  $Q$ . To finish description of notations for BFV formalism we mention that we will also need the momentum representation in the sector of the Lagrangian multipliers with the interchanged roles of  $N^\mu$  and  $p_\mu$ . It will be denoted by tilde, and the corresponding set of variables will look like

$$\tilde{Q}^I, \tilde{P}_I = q^i, p_i; p_\mu, -N^\mu; C^\mu, \mathcal{P}_\mu; \bar{C}_\mu, \bar{\mathcal{P}}^\mu, \tilde{\Psi}(\tilde{Q}) = \langle\langle \tilde{Q} ||\Psi\rangle\rangle. \quad (7.60)$$

The basic object of the BRST/BFV technique is the nilpotent fermionic BRST operator  $\hat{\Omega}$  acting in the space of  $||\Psi\rangle\rangle$  and satisfying the master equation

$$[\hat{\Omega}, \hat{\Omega}] \equiv \hat{\Omega}^2 = 0. \quad (7.61)$$

This equation allows one to look for the solution as an expansion in powers of the ghosts  $C^\mu$  and their momenta  $\mathcal{P}_\mu$  starting with the combination  $\hat{\Omega} = p_\alpha \bar{\mathcal{P}}^\alpha + C^\mu \hat{H}_\mu + O(\mathcal{P}C^2)$ . The coefficients of this expansion  $\hat{H}_\mu, \hat{U}_{\mu\nu}^\lambda, \hat{U}_{\mu\nu\alpha}^{\lambda\sigma}, \dots$  are the structure functions of the gauge algebra of constraints beginning with (7.49) – higher order structure functions follow from applying the Jacobi identity to multiple commutators of (7.49) with  $\hat{H}_\sigma$  [214, 215, 219]. In non-supersymmetric gravity theory, which we consider here, this sequence terminates at  $\hat{U}_{\mu\nu\alpha}^{\lambda\sigma} = 0$ , and  $\hat{\Omega}$  takes the form

$$\hat{\Omega} = p_\alpha \bar{\mathcal{P}}^\alpha + C^\mu \hat{H}_\mu + \frac{1}{2} C^\nu C^\mu \hat{U}_{\mu\nu}^\lambda \mathcal{P}_\lambda, \quad \hat{\Omega}^\dagger = \hat{\Omega}. \quad (7.62)$$

It is Hermitian in the BRST inner product (7.59) in accordance with the Hermiticity properties of ghost variables (7.57), provided the quantum Dirac constraints have the anti-Hermitian part<sup>6</sup>

$$\hat{H}_\mu - \hat{H}_\mu^\dagger = i\hat{U}_{\mu\lambda}^\lambda. \quad (7.63)$$

---

<sup>6</sup>In higher rank gauge theories with nonvanishing higher order structure functions this Hermiticity properties are modified by their higher order contributions.

In models with  $H_0(q, p) \neq 0$  this BRST operator determines also its BRST extension  $\hat{\mathcal{H}} = \hat{H}_0 + O(\mathcal{CP})$  by the equation  $[\hat{\Omega}, \hat{\mathcal{H}}_0] = 0$  and the so-called *unitarizing* Hamiltonian

$$\hat{\mathcal{H}}_\Phi = \hat{\mathcal{H}}_0 + \frac{1}{i}[\hat{\Phi}, \hat{\Omega}]. \quad (7.64)$$

It explicitly depends on the gauge fermion  $\Phi$ ,  $n(\hat{\Phi}) = 1$ , which provides gauge fixing in the BRST/BFV formalism. In parametrization invariant closed cosmology  $\hat{\mathcal{H}}_0 = 0$ , and the unitarizing Hamiltonian reduces to the commutator of the BRST operator and gauge fermion.

The unitary evolution operator  $\hat{U}_\Phi(t, t_-)$  acting in the space of  $\|\Psi\rangle\rangle$  is a solution of the following Cauchy problem:

$$i\hbar \frac{\partial}{\partial t} \hat{U}_\Phi(t, t_-) = \hat{\mathcal{H}}_\Phi \hat{U}_\Phi(t, t_-), \quad \hat{U}_\Phi(t_-, t_-) = \mathbb{I}. \quad (7.65)$$

It is obvious that from  $[\hat{\Omega}, [\hat{\Phi}, \hat{\Omega}]] \equiv 0$  and  $[\hat{\Omega}, \hat{\mathcal{H}}_\Phi] = 0$  the BRST operator is a constant of motion in this evolution,

$$[\hat{\Omega}, \hat{U}_\Phi(t, t_-)] = 0, \quad (7.66)$$

so that it plays the role of the conserved BRST charge and serves as a generator of BRST transformations in the relativistic phase space.

In the coordinate representation the kernel of the unitary evolution has a representation of the canonical path integral

$$\begin{aligned} \mathbf{U}_\Phi(t_+, Q_+ | t_-, Q_-) &\equiv \langle\langle Q_+ | \hat{U}_\Phi(t_+, t_-) | Q_- \rangle\rangle \\ &= \int_{Q(t_\pm)=Q_\pm} D[Q, P] \exp \left\{ i \int_{t_-}^{t_+} dt \left( P_I \dot{Q}^I - \mathcal{H}_\Phi(Q, P) \right) \right\}, \end{aligned} \quad (7.67)$$

where  $\mathcal{H}_\Phi(Q, P)$  is the  $QP$ -symbol of the unitarizing Hamiltonian given by the Poisson superbracket of  $c$ -number symbols  $\Phi$  and  $\Omega$  of the operator gauge fermion and BRST charge  $\mathcal{H}_\Phi(Q, P) = \{\Phi, \Omega\}$  (remember that  $H_0 = 0$ ). Also,  $D[Q, P]$  is a Liouville integration measure in the full boson-fermion phase space of  $c$ -number histories:

$$D[Q, P] = \prod_t dQ(t) \prod_{t^*} dP(t^*). \quad (7.68)$$

The difference between the set of points  $t = (t_N, \dots, t_1)$  and  $t^* = (t_{N+1}^*, \dots, t_1^*)$ ,  $N \rightarrow \infty$ , over which the product of integration measure factors is taken, reflects the typical slicing of the path integral into a sequence of multiple integrals in the decomposition of the full time segment  $[t_+, t_-]$  into infinitesimal pieces. This decomposition,  $t_+ > t_N > t_{N-1} > \dots > t_1 > t_-$ ,  $t_+ > t_{N+1}^* > t_N > t_N^* > \dots > t_1 > t_1^* > t_-$ , implies that the points  $t_i^*$ , at which the integrated

momenta are taken, are associated with “interiors” of the segments  $[t_{i+1}, t_i]$  whose boundaries carry the integrated coordinates – so that the number of momentum integrations is by one larger than those of coordinate ones.

In the momentum representation for Lagrangian multipliers  $\tilde{Q}^I = q^i, \pi^\mu, C^\mu, \tilde{C}_\mu$ , cf. (7.60), the unitary evolution kernel has a similar path integral representation:

$$\begin{aligned} \tilde{U}_\Phi(t_+, \tilde{Q}_+ | t_-, \tilde{Q}_-) &\equiv \langle \langle \tilde{Q}_+ | | \hat{U}_\Phi(t_+, t_-) | | \tilde{Q}_- \rangle \rangle \\ &= \int_{\tilde{Q}(t_\pm) = \tilde{Q}_\pm} \tilde{D}[Q, P] \exp \left\{ i \int_{t_-}^{t_+} dt \left( \tilde{P}_I \dot{\tilde{Q}}^I - \mathcal{H}_\Phi(Q, P) \right) \right\}, \end{aligned} \quad (7.69)$$

$$\tilde{D}[Q, P] = \prod_t d\tilde{Q}(t) \prod_{t^*} d\tilde{P}(t^*), \quad (7.70)$$

and is of course related by the Fourier transform to the kernel (7.67)

$$\tilde{U}_\Phi(t_+, \tilde{Q}_+ | t_-, \tilde{Q}_-) = \int dN_+ dN_- e^{-ip_+ N_+} \mathbf{U}_\Phi(t_+, Q_+ | t_-, Q_-) e^{ip_- N_-} \quad (7.71)$$

in full accordance with the fact that two symplectic forms in the integrands of path integrals on the left and right hand sides here are related by

$$\int_{t_-}^{t_+} dt \tilde{P} \dot{\tilde{Q}} = \int_{t_-}^{t_+} dt P \dot{Q} - p_+ N_+ + p_- N_-, \quad p_\pm \equiv \pi(t_\pm), \quad N_\pm \equiv N(t_\pm). \quad (7.72)$$

The principal theorem of the BFV quantization is that the matrix elements of the unitary evolution operator  $\hat{U}_\Phi(t, t_-)$  between the BRST-invariant physical states annihilated by  $\hat{\Omega}$  are independent of the choice of the gauge fermion [219]:

$$\hat{\Omega} | | \Psi_{1,2} \rangle \rangle = 0 \quad \implies \quad \delta_\Phi \langle \langle \Psi_1 | | \hat{U}_\Phi(t_+, t_-) | | \Psi_2 \rangle \rangle = 0. \quad (7.73)$$

The logic of the above BRST/BFV construction is based on the observation that relativistic gauge conditions, involving time derivatives of Lagrange multipliers, make the latter propagating and having nonvanishing canonical momenta  $p_\mu$  which are absent in the original action. To compensate the contribution of these artificially introduced degrees of freedom and the degrees of freedom which have to be excluded by first class constraints one introduces dynamical ghosts and antighosts of the statistics opposite to those of  $\hat{H}_\mu$ . Due to statistics they effectively subtract in quantum loops the contribution of these gauge degrees of freedom. However, a similar subtraction should be done in external lines of Feynman diagrams, which means that not all quantum states in BRST space are physical. Physical states form a subspace belonging to the kernel of the BRST operator. In this subspace due to the above theorem the transition amplitudes and quantum averages are independent of the choice of gauge fixing procedure – the corner stone of quantizing the gauge invariant systems.

*Batalin-Marnelius gauge fixing and the physical inner product*

Physical states should be BRST invariant  $\hat{\Omega} ||\Psi\rangle\rangle = 0$ . This equation does not uniquely select its solution because in view of the nilpotent nature of  $\hat{\Omega}$  the BRST transformed state  $||\Psi\rangle\rangle'$ ,

$$||\Psi\rangle\rangle' = ||\Psi\rangle\rangle + \hat{\Omega} ||\Phi\rangle\rangle, \quad (7.74)$$

with an arbitrary  $||\Phi\rangle\rangle$  also satisfies the BRST equation. This invariance results in the problem of constructing (or regulating) the physical inner product.

Problem is that the original inner product (7.59) for physical states represents the  $0 \times \infty$ -indeterminacy. The essence of this indeterminacy can be qualitatively explained by the fact that squaring of the physical state  $\Psi(Q) \sim \delta(\hat{\Omega})$  in (7.59) gives a divergent factor whereas the integration over Grassman variables multiplies it by zero. This inner product can be regulated by transforming the BRST-invariant state  $||\Phi\rangle\rangle$  to a special gauge as it was suggested by Batalin and Marnelius in [221, 222] (see also [223]):

$$||\Psi\rangle\rangle \rightarrow ||\Psi_{BM}\rangle\rangle : \hat{\mathcal{P}}_\mu ||\Psi_{BM}\rangle\rangle = 0, \hat{N}^\mu ||\Psi_{BM}\rangle\rangle = 0, \quad (7.75)$$

$$\hat{\mathcal{P}}^\mu ||\Psi_{BM}\rangle\rangle = 0 \quad (7.76)$$

(the last condition is in fact a corollary of the second one,  $[\hat{\Omega}, N^\mu] ||\Psi_{BM}\rangle\rangle = 0$ ). The consistency of this gauge with the BRST-invariance of  $||\Psi_{BM}\rangle\rangle$  implies that it also satisfies the quantum Dirac constraints

$$0 = \frac{1}{i} [\hat{\Omega}, \hat{\mathcal{P}}_\mu] ||\Psi_{BM}\rangle\rangle = (\hat{H}_\mu + C^\nu \hat{U}_{\nu\mu}^\lambda \hat{\mathcal{P}}_\lambda) ||\Psi_{BM}\rangle\rangle = \hat{H}_\mu ||\Psi_{BM}\rangle\rangle, \quad (7.77)$$

and its wave function is independent of ghost variables,

$$\frac{\partial}{\partial C^\mu} \Psi_{BM}(Q) = 0, \quad \frac{\partial}{\partial \bar{C}_\mu} \Psi_{BM}(Q) = 0. \quad (7.78)$$

Together with (7.75) this means that  $\Psi_{BM}(Q)$  has the form

$$\langle\langle Q ||\Psi_{BM}\rangle\rangle = \langle q | \Psi \rangle \delta(N) \equiv \Psi(q) \delta(N), \quad (7.79)$$

where the ‘‘matter’’ part  $\Psi(q)$  satisfies quantum Dirac constraints in the coordinate representation,

$$\hat{H}_\mu \Psi(q) = 0. \quad (7.80)$$

According to [221, 222] the physical inner product of wave functions  $||\Psi_{BM}\rangle\rangle$  can be regularized by a special operator-valued measure which is explicitly built with the aid of the gauge fermion  $\hat{\Phi}$

$$\langle\langle \Psi' ||\Psi \rangle\rangle_{\text{phys}} = \langle\langle \Psi'_{BM} || e^{[\hat{\Phi}, \hat{\Omega}]} ||\Psi_{BM}\rangle\rangle. \quad (7.81)$$

The choice of this fermion is immaterial because

$$\delta_{\hat{\Phi}} \langle \langle \Psi' | \Psi \rangle \rangle_{\text{phys}} = \int_0^1 ds \langle \langle \tilde{\Psi}'_{BM} | e^{(1-s)[\hat{\Phi}, \hat{\Omega}]} [\delta \hat{\Phi}, \hat{\Omega}] e^{s[\hat{\Phi}, \hat{\Omega}]} | \Psi_{BM} \rangle \rangle = 0, \quad (7.82)$$

since  $[\hat{\Omega}, [\hat{\Phi}, \hat{\Omega}]] = 0$  and  $\hat{\Omega} | \Psi_{BM} \rangle = 0$ . However, with a special choice of this fermion the physical inner product (7.81) resolves the  $0 \times \infty$  uncertainty and becomes well defined [221].

This is easy to show if this fermion is constructed with the aid of gauge conditions functions  $\hat{\chi}^\mu = \chi^\mu(\hat{q}, \hat{p})$  which commute with themselves,  $[\hat{\chi}^\mu, \hat{\chi}^\nu] = 0$ , and with the structure functions operators,  $[\hat{\chi}^\mu, \hat{U}_{\alpha\beta}^\lambda] = 0$  (in gravity theory this is coordinate gauge conditions  $\chi^\mu(q)$  commuting with  $U_{\alpha\beta}^\lambda = U_{\alpha\beta}^\lambda(q)$ ). For the fermion  $\hat{\Phi}_{BM}$  in the form  $\hat{\Phi}_{BM} = \hat{C}_\mu \hat{\chi}^\mu$  we have

$$[\hat{\Phi}_{BM}, \hat{\Omega}] = ip_\mu \hat{\chi}^\mu + i\bar{C}_\mu \hat{J}_\nu^\mu C^\nu, \quad \hat{J}_\nu^\mu = \frac{1}{i} [\hat{\chi}^\mu, \hat{H}_\nu]. \quad (7.83)$$

Then the physical inner product for Batalin-Marnelius wave functions (7.79) in the coordinate representation with  $\hat{p}_\mu = \partial/i\partial N^\mu$  takes the form

$$\begin{aligned} \langle \langle \Psi' | \Psi \rangle \rangle_{\text{phys}} &= \int dq dN dC d\bar{C} \Psi'^*(q) \delta(N) e^{-i\bar{C}_\mu \hat{J}_\nu^\mu C^\nu + \hat{\chi}^\mu \frac{\partial}{\partial N^\mu}} \delta(N) \Psi(q) \\ &= \langle \Psi' | \int d\pi dC d\bar{C} e^{-i\bar{C}_\mu \hat{J}_\nu^\mu C^\nu + ip_\mu \hat{\chi}^\mu} | \Psi \rangle. \end{aligned} \quad (7.84)$$

This leads to the physical inner product as a special operator valued measure  $\hat{M}$  acting in the space of Dirac wave functions  $|\Psi'\rangle$  and  $|\Psi\rangle$  endowed with the auxiliary  $L^2$  inner product (7.53),

$$\langle \Psi' | \Psi \rangle_{\text{phys}} \equiv \langle \langle \Psi' | \Psi \rangle \rangle_{\text{phys}} = \langle \Psi' | \hat{M} | \Psi \rangle, \quad (7.85)$$

$$\hat{M} = \int d\pi dC d\bar{C} e^{-i\bar{C}_\mu \hat{J}_\nu^\mu C^\nu + ip_\mu \hat{\chi}^\mu} = \delta(\hat{\chi}) \det \hat{J}_\nu^\mu (1 + O([\hat{\chi}, \hat{J}])). \quad (7.86)$$

This measure is known in quadratures as an explicit integral over ghost fields and Lagrangian multipliers momenta. In the leading semiclassical order in  $[\hat{\chi}, \hat{J}] = O(\hbar)$ , this integral allows one to disentangle the delta function of gauge conditions,  $\delta(\hat{\chi}) = \prod_\mu \delta(\hat{\chi}^\mu)$ , which is well defined in view of their commutativity.<sup>7</sup>

The expression (7.86) is the analogue of the time-local measure in the canonical Faddeev-Popov path integral [202] with the operator (7.83) semiclassically represented by the Poisson bracket  $\hat{J}_\nu^\mu = \{\chi^\mu, H_\nu\}$ . As we will show below by direct calculations in the one-loop (subleading in  $\hbar$ ) order this inner product [205, 206, 208, 209]

$$\langle \Psi' | \Psi \rangle_{\text{phys}} = \int dq \Psi'^*(q) \delta(\chi(q)) \det \hat{J}_\nu^\mu \Psi(q) + O(\hbar) \quad (7.87)$$

---

<sup>7</sup>This, however, does not save us from extra corrections, because  $J_\nu^\mu(\hat{q}, \hat{p})$  depends on the momentum  $\hat{p}_i$  and is a differential operator acting in the space of  $q$ .

for semiclassical wave functions  $\Psi(q)$  and  $\Psi'(q)$  is independent of the choice of  $\chi^\mu(q)$  and is consistent with the Hermiticity property<sup>8</sup> (7.63) [209].

### *Path integral representation in Dirac quantization scheme*

Transition to the BM gauge can in fact be obtained by a simple procedure of truncation of BRST invariant wave functions to the sector of “matter” variables, that was suggested in [206, 208]. Introduce a wave function  $\Psi(q)$  in the representation space of  $\hat{q}^i, \hat{p}_i$  which can be obtained from the solution  $|\Psi\rangle$  of the BRST equation  $\hat{\Omega}|\Psi\rangle = 0$  by

$$\Psi(q) = \int dN \Psi(q, N, C, \bar{C})|_{C=0}. \quad (7.88)$$

As we show below, from the BRST equation it follows that this function satisfies quantum Dirac constraints and is independent of the antighost variable  $\bar{C}$  (that is why the argument  $\bar{C}$  of the right-hand side is omitted on the left-hand side of this definition):

$$\hat{H}_\mu \Psi(q) = 0, \quad \frac{\partial}{\partial \bar{C}_\mu} \Psi(q) = 0. \quad (7.89)$$

Both properties follow from the BRST equation which in the coordinate representation reads as

$$\hat{\Omega} \Psi(Q) = \left( \frac{\partial}{\partial N^\mu} \frac{\partial}{\partial \bar{C}_\mu} + C^\mu \hat{H}_\mu + \frac{i}{2} C^\nu C^\mu \hat{U}_{\mu\nu}^\lambda \frac{\partial}{\partial C^\lambda} \right) \Psi(Q) = 0. \quad (7.90)$$

Integrating this equation over the Lagrange multipliers  $N$  in infinite limits and assuming that  $\Psi(Q)$  falls off sufficiently rapidly at  $N \rightarrow \pm\infty$ , one finds that the first term of (7.90) vanishes. Then one can differentiate the result with respect to the ghost field  $C^\mu$  and subsequently put  $C = 0$ . This proves the first of relations (7.89). The second relation follows from multiplying Eq. (7.90) by  $N^\mu$ , putting  $C = 0$  and integrating over  $N$  by parts in the remaining first term of this equation

$$0 = \int dN N^\mu \hat{\Omega} \Psi(Q)|_{C=0} = \int dN N^\mu \frac{\partial}{\partial N^\nu} \frac{\partial}{\partial \bar{C}_\nu} \Psi(Q)|_{C=0}. \quad (7.91)$$

This truncation of the BRST quantization scheme to the Dirac quantization suggested in [206, 208] serves in fact as a realization of the Batalin-Marnelius gauge fixing (7.75) of the BRST symmetry (7.74). To put the generic BRST state into the Batalin-Marnelius gauge it is enough to take the bosonic “body”

---

<sup>8</sup>Even though this Hermiticity property holds with respect to another – auxiliary – inner product (7.53) different from (7.87).

of its wave function, integrate it over the Lagrange multipliers argument  $N$  and multiply by  $\delta(N)$ :

$$||\Psi\rangle\rangle \rightarrow ||\Psi_{BM}\rangle\rangle : \quad \Psi_{BM}(Q) = \delta(N) \int dN' \Psi(q, N', C, \bar{C}) \Big|_{C=0}. \quad (7.92)$$

Truncation similar to (7.88) for the kernel of the unitary evolution (7.67) reads

$$U(q_+, q_-) = \int dN_+ dN_- \mathbf{U}_\Phi(t_+, Q_+ | t_-, Q_-) \Big|_{C_\pm=0}. \quad (7.93)$$

This object can be represented as a matrix element of  $\hat{U}_\Phi(t_+, t_-)$  between the following two states  $||\Psi_\pm\rangle\rangle$  which are both zero vectors of the Lagrangian multiplier momentum and trivially satisfy the BRST equation:

$$U(q_+, q_-) \equiv \langle\langle \Psi_+ | \hat{U}_\Phi(t_+, t_-) | \Psi_- \rangle\rangle, \quad (7.94)$$

$$\Psi_\pm(q, N, C, \bar{C}) = \delta(q - q_\pm) \delta(C) \delta(\bar{C}), \quad (7.95)$$

$$\hat{p}_\alpha ||\Psi_\pm\rangle\rangle = 0, \quad \hat{\Omega} ||\Psi_\pm\rangle\rangle = 0. \quad (7.96)$$

Therefore, in virtue of the main theorem of BRST quantization it is independent of the choice of the gauge fermion  $\Phi$  in  $\hat{U}_\Phi(t_+, t_-)$ :

$$\delta_\Phi U(q_+, q_-) = 0, \quad (7.97)$$

which guarantees the uniqueness of its definition. The second important property is that the kernel (7.93) is independent of  $t_\pm$  in parametrization invariant theory with  $H_0 = 0$  because

$$i \frac{\partial}{\partial t_+} U(q_+, q_-) = \langle\langle \Psi_+ | \left[ \hat{\Phi}, \hat{\Omega} \right] \hat{U}_\Phi(t_+, t_-) | \Psi_- \rangle\rangle = 0 \quad (7.98)$$

in view of the Schrödinger equation (7.65) for  $\hat{U}_\Phi(t_+, t_-)$ . This allows one to omit  $\Phi$  and  $t_\pm$  labels in the left-hand side of the definition (7.93).

Finally, applying the same derivations as in (7.90)-(7.91) to the main BRST equation (7.66) for  $\hat{U}_\Phi(t_+, t_-)$ , one proves that this kernel is independent of antighosts  $\bar{C}_\mu^\pm$  and satisfies quantum Dirac constraints with respect to both arguments

$$\hat{H}_\mu U(q, q') = 0, \quad U(q, q') \overleftarrow{H}_\mu^\dagger = 0. \quad (7.99)$$

Integration over  $N_\pm$  in (7.93) implies that this kernel can be interpreted as the unitary evolution kernel in the momentum representation of Lagrange multipliers (7.71) at zero values of  $p_\pm$ ,

$$\begin{aligned} U(q, q') &= \int dN_+ dN_- e^{-ip_+ N_+} \mathbf{U}_\Phi(t_+, Q_+ | t_-, Q_-) e^{ip_- N_-} \Big|_{p_\pm = C_\pm = 0} \\ &= \tilde{U}_\Phi(t_+, \tilde{Q}_+ | t_-, \tilde{Q}_-) \Big|_{p_\pm = C_\pm = 0}. \end{aligned} \quad (7.100)$$

Therefore, it has the path integral representation (7.69) with the symbol of the unitarizing Hamiltonian – the Poisson superbracket of the gauge fermion  $\Phi$  and BRST charge  $\Omega$ ,

$$U(q_+, q_-) = \int_{\tilde{Q}(t_{\pm})=\tilde{Q}_{\pm}} \tilde{D}[Q, P] \exp \left[ i \int_{t_-}^{t_+} dt (\tilde{P}_I \dot{\tilde{Q}}^I - \{\Phi, \Omega\}) \right] \Big|_{p_{\pm}=C_{\pm}=0} \quad (7.101)$$

There exists a special choice of the gauge fermion,  $\Phi = \mathcal{P}_{\mu} N^{\mu} + \bar{C}_{\mu} \chi^{\mu}(q)$ , which generates in the Lagrangian formalism (after integrating out the phase space momenta) calculationally most useful relativistic gauge conditions of the form  $\dot{N}^{\mu} - \chi^{\mu}(q) = 0$  [213, 46, 217, 218]. With this fermion (note that it differs from the BM gauge fermion  $\Phi = \bar{C}_{\mu} \chi^{\mu}(q)$ ) the path integral becomes

$$U(q_+, q_-) = \int \tilde{D}[Q, P] \exp \left[ i \int_{t_-}^{t_+} dt (\tilde{P}_I \dot{\tilde{Q}}^I - N^{\mu} H_{\mu} - p_{\mu} \chi^{\mu} - \bar{C}_{\mu} J_{\nu}^{\mu} C^{\nu} - \mathcal{P}_{\alpha} (\bar{\mathcal{P}}^{\alpha} + U_{\mu\nu}^{\alpha} N^{\mu} C^{\nu})) \right] \Big|_{\tilde{Q}(t_{\pm})=\tilde{Q}_{\pm} p_{\pm}=C_{\pm}=0}. \quad (7.102)$$

A standard procedure of transition to the unitary gauge  $\chi^{\mu}(q) = 0$  consists in rescaling the gauge function  $\chi^{\mu}$  by a small numerical parameter  $\varepsilon$ ,  $\chi^{\mu} \rightarrow \chi^{\mu}/\varepsilon$  and making the change of integration variables  $p_{\mu}$  and  $\bar{C}_{\mu}$  (with a unit Jacobian)

$$p_{\mu} \rightarrow \varepsilon p_{\mu}, \quad \bar{C}_{\mu} \rightarrow \varepsilon \bar{C}_{\mu}. \quad (7.103)$$

Note that boundary conditions at  $t = t_{\pm}$  admit this change of variables, and all this does not affect the answer for  $U(q, q')$  in view of its gauge independence. Then in the limit  $\varepsilon \rightarrow 0$  the kinetic terms  $-N^{\mu} \dot{p}_{\mu}$  and  $\bar{\mathcal{P}}^{\alpha} \dot{\bar{C}}_{\alpha}$  disappear from the integrand of (7.102), Gaussian integration over ghost momenta does not contribute any field-dependent measure, and the projector to Dirac states takes the form of the usual canonical Faddeev-Popov path integral

$$\begin{aligned} U(q_+, q_-) &= \int D[q, p] DN D\pi DC D\bar{C} \times \\ &\times \exp \left\{ i \int_{t_-}^{t_+} dt (p_i \dot{q}^i - N^{\mu} H_{\mu} - p_{\mu} \chi^{\mu} - \bar{C}_{\mu} J_{\nu}^{\mu} C^{\nu}) \right\} \Big|_{q(t_{\pm})=q_{\pm}, p_{\pm}=C_{\pm}=0} \\ &= \int_{q(t_{\pm})=q_{\pm}} D[q, p] DN \left( \prod_{t \neq t_{\pm}} \delta(\chi) \det J_{\nu}^{\mu} \right) e^{i \int_{t_-}^{t_+} dt (p_i \dot{q}^i - N^{\mu} H_{\mu})}, \quad (7.104) \end{aligned}$$

in which, however, the gauge fixing factor  $\delta(\chi) \det J_{\nu}^{\mu}$  is absent at the both end points  $t_{\pm}$ . This completely specifies the expression in the right-hand side (7.11) formulated above and proves all its properties – projection on the subspace of Dirac constraints (7.99), gauge independence (7.97) and independence of the choice of  $t_{\pm}$  (7.98). Its relation to the physical sector evolution operator standing in the left-hand side of (7.11) is discussed below.

## 7.5. Semiclassical approximation

The formalism of the above type should stand verification by a viable calculational scheme. Such a scheme applicable to theories of a general type is the semiclassical loop expansion in powers of  $\hbar$ . So here we reinstate the  $\hbar$  parameter (thus far chosen to be 1 in universal units) and develop this expansion in the first (one-loop) order linear in the Planck constant.

### *Operator realization of constraints*

The first thing to do is to find the operator realization of constraints  $\hat{H}_\mu$  and structure functions  $\hat{U}_{\mu\nu}^\lambda$  that would satisfy the quantum algebra (7.49) with the operators of structure functions standing to the left of the constraints (with the Planck constant reinstated the right-hand side of these algebraic relations should be multiplied by  $\hbar$ ). Remarkably, in the approximation linear in  $\hbar$  there exists in a closed form the following solution to this problem – the operators are given by the Weyl (symmetric in  $\hat{q}$  and  $\hat{p}$ ) ordering of the following expressions [206, 209]:

$$\hat{H}_\mu = N_W \left\{ H_\mu(\hat{q}, \hat{p}) + \frac{i\hbar}{2} U_{\mu\nu}^\nu(\hat{q}, \hat{p}) + O(\hbar^2) \right\}, \quad (7.105)$$

$$\hat{U}_{\mu\nu}^\lambda = N_W \left\{ U_{\mu\nu}^\lambda(\hat{q}, \hat{p}) - \frac{i\hbar}{2} U_{\mu\nu\sigma}^{\lambda\sigma}(\hat{q}, \hat{p}) + O(\hbar^2) \right\}. \quad (7.106)$$

Here  $N_W$  is a symbol of Weyl ordering, and these algorithms are true for a generic theory subject to first class constraints having at the classical level the hierarchy of structure functions of the Poisson bracket algebra beginning with  $H_\mu(q, p)$ ,  $U_{\mu\nu}^\lambda(q, p)$ ,  $U_{\mu\nu\omega}^{\lambda\sigma}(q, p)$ , ..., as mentioned above higher order structure functions following from the Jacobi identities applied to multiple commutators of (7.49) with  $\hat{H}_\sigma$  [214, 215, 219]. In Einstein gravity theory, of course,  $U_{\mu\nu}^\lambda$  is  $p$ -independent and  $U_{\mu\nu\omega}^{\lambda\sigma} \equiv 0$ . Note a nontrivial anti-Hermitian part of the constraint operator generated by the trace of the structure constant, which is compatible with the Hermiticity properties of the BfV operators (7.63) relative to the *auxiliary* inner product (7.53) in the space of  $\Psi(q)$ .

A similar algorithm holds for the physical observables – functions  $\mathcal{O}_I(q, p)$  on the phase space which are classically gauge invariant in a weak sense, that is commuting with classical constraints modulo the constraints themselves,  $\{\mathcal{O}_I, H_\mu\} = V_{I\mu}^\nu H_\nu$ . They read [209]

$$\hat{\mathcal{O}}_I = N_W \left\{ \mathcal{O}_I + \frac{i\hbar}{2} V_{I\lambda}^\lambda + O(\hbar^2) \right\}, \quad (7.107)$$

$$\hat{V}_{I\mu}^\nu = N_W \left\{ V_{I\mu}^\nu - \frac{i\hbar}{2} V_{I\mu\sigma}^{\nu\sigma} + O(\hbar^2) \right\}. \quad (7.108)$$

This operator realization applies, in particular, to  $\hat{H}_0$  and  $\hat{V}_\mu^\nu \equiv \hat{V}_{0\mu}^\nu$  of Eq.(7.49) in theories with a nonvanishing Hamiltonian  $H_0(q, p)$ .<sup>9</sup>

### *Semiclassical physical states*

Semiclassical wave functions are characterized by the Hamilton-Jacobi function  $S(q)$  and pre-exponential factor  $P(q)$  expandable in  $\hbar$ -series beginning with the one-loop order  $O(\hbar^0)$ :

$$\Psi(q) = P(q) \exp \left[ \frac{i}{\hbar} S(q) \right]. \quad (7.109)$$

The general semiclassical solution of quantum constraints (7.99) with operators (7.105) was found in [205, 206, 208] in the form of the two-point kernel  $U(q, q')$  “propagating” the initial data from  $q'$  throughout the space of  $q$

$$U(q, q') = P(q, q') \exp \left[ \frac{i}{\hbar} S(q, q') \right]. \quad (7.110)$$

In both expressions (7.109) and (7.110) the phase in the exponential satisfies the Hamilton-Jacobi equation (2.44), while the one-loop pre-exponential factor is subject to continuity type equation originating from the full quantum constraint in the approximation linear in  $\hbar$  [205, 206, 209]:

$$\frac{\partial}{\partial q^i} (\nabla_\mu^i P^2) = U_{\mu\lambda}^\lambda P^2, \quad (7.111)$$

$$\nabla_\mu^i \equiv \left. \frac{\partial H_\mu}{\partial p_i} \right|_p = \partial S / \partial q. \quad (7.112)$$

Note a nontrivial right-hand side in the “continuity” equation, generated by the anti-Hermitian part of  $\hat{H}_\mu$ .

For a two-point kernel the Hamilton-Jacobi function coincides with the principal Hamilton function  $S(q, q')$  (action on the extremal joining points  $q$  and  $q'$ ) and the solution of the continuity equation can be found as a generalization of the Pauli-Van Vleck-Morette ansatz for the one-loop pre-exponential factor [224, 208] of the Schrödinger propagator. This generalization is nothing but a Faddeev-Popov gauge-fixing [225] procedure for a matrix of mixed second-order derivatives of the principal Hamilton function:

$$S_{ik'} = \frac{\partial^2 S(q, q')}{\partial q^i \partial q^{k'}}, \quad (7.113)$$

which is degenerate in virtue of the Hamilton-Jacobi equations (2.44) giving rise to the left zero-value eigenvectors (7.112) and analogous right zero-vectors

---

<sup>9</sup>If the observables satisfy the Poisson algebra  $\{\mathcal{O}_I, \mathcal{O}_J\} = U_{IJ}^L \mathcal{O}_L + U_{IJ}^\lambda H_\lambda$ ,  $U_{IJ}^L = \text{const}$ , the above expressions are also modified by the trace of structure constants  $U_{IL}^L$  [209].

[205, 206]:

$$\nabla_{\mu}^i S_{ik'} = 0, \quad S_{ik'} \nabla_{\nu}^{k'} = 0, \quad \nabla_{\nu}^{k'} \equiv \left. \frac{\partial H_{\nu}(q', p')}{\partial p'_{k'}} \right|_{p'} = -\partial S / \partial q'. \quad (7.114)$$

The pre-exponential factor reads

$$P = \left[ \frac{\det F_{ik'}}{J(q)J(q') \det c_{\mu\nu}} \right]^{1/2}, \quad (7.115)$$

where  $F_{ik'}$  is a nondegenerate matrix of the initial action Hessian (7.113) supplied with a gauge-breaking term

$$F_{ik'} = S_{ik'} + \chi_i^{\mu} c_{\mu\nu} \chi_{k'}^{\nu}, \quad (7.116)$$

and  $J(q)$  and  $J(q')$  are the Faddeev-Popov “ghost” determinants [225] compensating for the inclusion of this term. They are constructed with the aid of two sets of arbitrary covectors  $(\chi_i^{\mu}, \chi_{k'}^{\nu})$  (“gauge” conditions) satisfying the only requirement of the nondegeneracy of their ghost operators [205, 208, 206]:

$$\begin{aligned} J_{\nu}^{\mu}(q) &= \chi_i^{\mu} \nabla_{\nu}^i, & J(q) &\equiv \det J_{\nu}^{\mu}(q) \neq 0, \\ J_{\nu}^{\mu}(q') &= \chi_{i'}^{\mu} \nabla_{\nu}^{i'}, & J(q') &\equiv \det J_{\nu}^{\mu}(q') \neq 0. \end{aligned} \quad (7.117)$$

The invertible gauge-fixing matrix  $c_{\mu\nu}$  and its determinant are the last ingredients of the generalized Pauli-Van Vleck-Morette ansatz (7.115).

### *Reduction to the physical sector*

The interpretation of the semiclassical state (7.110), (7.115) is rather transparent in the physical sector of the theory [205, 206, 208]. The sector explicitly arises after the reduction to physical variables by disentangling them from the original phase space of  $(q^i, p_i)$  in a unitary gauge. Such a reduction in [205, 206, 208] was given for a special type of gauge conditions imposed only on phase space coordinates  $q^i$ ,  $\chi^{\mu}(q) = 0$ . These *coordinate* gauge conditions determine the embedding of the  $(n - m)$ -dimensional space  $\Sigma$  of physical *coordinates* directly into the space of original coordinates  $q^i$  – superspace. This fact strongly simplifies the reduction of the semiclassical kernel (7.110), (7.115) to the physical sector, because this reduction in the main boils down to the embedding of the arguments of  $U(q, q')$  into the physical subspace  $\Sigma$ . The geometry of this embedding, considered in much detail in [208], can be better described in special coordinates on superspace  $\bar{q}^i = (\xi^A, \theta^{\mu})$ , in which  $g^A$ ,  $A = 1, \dots, n - m$ , serve as intrinsic coordinates on  $\Sigma$  (physical configuration coordinates), and  $\theta^{\mu}$  is determined by gauge conditions:

$$q^i \rightarrow \bar{q}^i = (g^A, \theta^{\mu}), \quad q^i = e^i(g^A, \theta^{\mu}), \quad \theta^{\mu} = \chi^{\mu}(q). \quad (7.118)$$

The equation of the surface  $\Sigma$  in the new coordinates is  $\theta^\mu = 0$ , so that its embedding equations coincide with the above reparametrization equations at  $\theta^\mu = 0$ ,  $e^i(g) = e^i(g, 0)$

$$\Sigma : q^i = e^i(g), \quad \chi^\mu(e^i(g)) \equiv 0. \quad (7.119)$$

The relation between the integration measures on superspace  $dq = d^n q$  and on  $\Sigma$ ,  $dg = d^{n-m} g$ ,

$$dg = dq \delta(\chi) \mathcal{M}, \quad \delta(\chi) = \prod_{\mu} \delta(\chi^\mu(q)), \quad \mathcal{M} = (\det[e_A^i, e_\mu^i])^{-1}, \quad (7.120)$$

involves the Jacobian of this reparametrization, built of the basis of vectors tangential and normal to  $\Sigma$ :

$$e_A^i = \frac{\partial e^i}{\partial g^A}, \quad e_\mu^i = \frac{\partial e^i}{\partial \theta^\mu}. \quad (7.121)$$

Note that  $m$  covectors normal to the surface can be chosen as gradients of gauge conditions

$$\chi_i^\mu = \frac{\partial \chi^\mu}{\partial q^i}, \quad \chi_i^\mu e_\nu^i = \delta_\nu^\mu, \quad (7.122)$$

that can be identified with auxiliary covectors participating in the algorithm for the pre-exponential factor (7.115). With this identification the Faddeev-Popov operator  $J_\nu^\mu(q, \partial S / \partial q)$  coincides with the operator  $J_\nu^\mu(q)$  of this algorithm (which explains the use of the same notation).

On the same footing with  $(e_A^i, e_\mu^i)$  as a full local basis one can also choose the set  $(e_A^i, \nabla_\mu^i)$  with vectors  $\nabla_\mu^i$  transversal to  $\Sigma$  given by eq.(7.112). The normal vectors of the first basis when expanded in the new basis,  $e_\mu^i = J^{-1}{}^\nu{}_\mu \nabla_\nu^i + \Omega_\mu^A e_A^i$ , have one expansion coefficient always determined by the inverse of the Faddeev-Popov matrix  $J^{-1}{}^\nu{}_\mu$  and, thus, independent of the particular parametrization of  $\Sigma$  by internal coordinates. The second coefficient is less universal and depends on a particular choice of this parametrization. Missing information about  $\Omega_\mu^A$  does not prevent, however, from finding the relation between the determinants of matrices of the old and new bases:

$$\det[e_A^i, \nabla_\mu^i] = \frac{J}{\mathcal{M}}. \quad (7.123)$$

The reduction to physical sector in coordinate gauges follows after identifying  $g^A$  with the physical coordinates. The corresponding conjugated momenta  $p_A$  can be found from the transformation of the symplectic form restricted to the physical subspace (7.119)

$$\int dt p_i \dot{q}^i = \int dt \left( p_i e_A^i \dot{g}^A + p_i \frac{\partial e^i(g, t)}{\partial t} \right), \quad (7.124)$$

$$p_A = p_i e_A^i, \quad (7.125)$$

as projections of the original momentum to the tangential components of the basis (7.121). The normal projections of  $p_i$  should be found from the constraints  $H_\mu(q, p) = 0$ , the local uniqueness of their solution being granted by the nondegeneracy of the Faddeev-Popov determinant. Together with (7.119) this solution yields all the original phase space variables  $(q^i, p_i)$  as known functions of the physical degrees of freedom  $(g^A, p_A)$ . The original action (2.124) reduced to physical sector (that is to the subspace of constraints and gauge conditions) acquires the usual canonical form with the physical Hamiltonian contributed by the second term of (7.124) and  $H_0(q, p)$  when the latter is nonvanishing. Note that generically, especially for systems with  $H_0(q, p) = 0$ , the canonical gauge conditions should explicitly depend on time,  $\chi^\mu(q) = \chi^\mu(q, t)$ , in order to generate the dynamical evolution in reduced phase space theory [208, 206]. Therefore, the reduced symplectic form generates a nontrivial contribution to the physical Hamiltonian, proportional to the time derivative of the embedding functions (7.119) explicitly depending on  $t$ ,  $q^i = e^i(g^A, t)$ . The total physical Hamiltonian then takes the form

$$H_{\text{phys}}(g, \pi) = \left[ H_0(q, p) - p_i \frac{\partial e^i(g, t)}{\partial t} \right]_{q=q(g, \pi), p=p(g, \pi)}, \quad (7.126)$$

where we retain for the sake of generality also the nonzero Hamiltonian  $H_0(q, p)$ .

Canonical quantization of such a classical system runs as usual along the lines of a particular representation and operator realization in the Hilbert space of the theory. In the one-loop (linear in  $\hbar$ ) approximation with the Weyl ordering of the above Hamiltonian this quantization is basically exhausted by the unitary evolution kernel  $U(t, g|t', g')$  of the Schrödinger equation. In the coordinate representation it is given by the well-known Pauli-Van Vleck-Morette ansatz [224]

$$U_{\text{phys}}(t, g|t', g') \equiv \left[ \det \frac{i}{2\pi\hbar} \frac{\partial^2 S(t, g|t', g')}{\partial g^A \partial g^{B'}} \right]^{1/2} e^{\frac{i}{\hbar} S(t, g|t', g')}, \quad (7.127)$$

where the principal Hamilton function of the physical variables  $S(t, g|t', g')$  is a classical action evaluated at the classical extremal passing the points  $g'$  and  $g$  respectively at initial  $t'$  and final  $t$  moments of time. The pre-exponential factor here is built of Van Vleck determinant and guarantees in the approximation linear in  $\hbar$  the unitarity of the Schrödinger evolution of the physical states  $\Psi(t, g)$

$$\Psi_{\text{phys}}(t, g) = \int dg' U_{\text{phys}}(t, g|t', g') \Psi_{\text{phys}}(t', g') \quad (7.128)$$

in the Hilbert space with a simple  $L^2$  inner product (denoted by round brackets)

$$(\Psi'_{\text{phys}} | \Psi_{\text{phys}}) \equiv \int dg \Psi'^*_{\text{phys}}(g) \Psi_{\text{phys}}(g). \quad (7.129)$$

The unitary map between the reduced phase space quantization of the above type and the Dirac quantization of Sects. 7.2.–7.4. consists in a special relation between the two-point kernel (7.110) with prefactor (7.115) and the Schrödinger evolution operator (7.127) [205, 206, 208]. This relation is based on the equality of the principal Hamilton functions in the original constrained theory and the reduced one and its corollary – the relation between their Van Vleck matrices

$$S(t, g|t', g') = S(q, q') \Big|_{q=e(g,t), q'=e(g',t')}, \quad (7.130)$$

$$S_{ik'} e_A^i e_{B'}^{k'} = \frac{\partial^2 S(t, g|t', g')}{\partial g^A \partial g^{B'}}. \quad (7.131)$$

Decomposing the gauge-fixed matrix (7.116) in the basis of vectors ( $e_A^i, \nabla_\mu^i$ ) and using (7.123) one then easily finds the needed relation [205, 206, 208]

$$\begin{aligned} U_{\text{phys}}(t, g|t', g') \\ = \left( \frac{J}{\mathcal{M}} \right)^{1/2} U(q, q') \left( \frac{J'}{\mathcal{M}'} \right)^{1/2} \Big|_{q=e(g,t), q'=e(g',t')} + O(\hbar). \end{aligned} \quad (7.132)$$

This is the semiclassical (one-loop) implementation of the relation (7.11) formulated above.

This relation implies that the kernel  $U(q, q')$  similarly to the Schrödinger propagator  $U_{\text{phys}}(t, g|t', g')$  can be regarded as a propagator of the Dirac wave function  $\Psi(q)$  in superspace. Indeed, introducing the following map between  $\Psi(q)$  and  $\Psi_{\text{phys}}(t, g)$

$$\Psi_{\text{phys}}(g, t) = \left( \frac{J}{\mathcal{M}} \right)^{1/2} \Psi(q) \Big|_{q=e(g,t)} + O(\hbar) \quad (7.133)$$

and taking into account the relation (7.120) between the integration measures on superspace and the physical space  $\Sigma$ , one finds that the propagation law (7.128) in  $g$ -space can be regarded as a projection onto  $\Sigma(t)$  of the following propagation of the Dirac wave function  $\Psi(q)$  from the initial Cauchy surface  $\Sigma(t')$  in  $q$ -space to the entire superspace:

$$\Psi(q) = \int dq' U(q, q') \delta(\chi(q', t')) J\left(q', -\frac{\partial S(q, q')}{\partial q'}\right) \Psi(q') + O(\hbar). \quad (7.134)$$

Here the actual integration runs over the initial physical space  $\Sigma(t')$ . However, the integration measure involves not just local quantities at this surface, but also the normal derivatives of the kernel (or the wave function  $\Psi(q')$  itself) arising in the one-loop approximation as a Hamilton-Jacobi argument  $p = -\partial S(q, q')/\partial q'$  of  $J(q, p)$ .

*Semiclassical physical inner product: Hermiticity and gauge independence*

The unitary map from the Dirac quantization to the reduced phase space quantization (7.133) suggests that the simple inner product (7.129) of the former quantization scheme induces a correct physical product in the latter one

$$\langle \Psi' | \Psi \rangle_{\text{phys}} = (\Psi'_{\text{phys}} | \Psi_{\text{phys}}) \quad (7.135)$$

on account of the relation (7.133) and the change of integration variables (7.122). The result for semiclassical states of the form (7.109) looks exactly like (7.81) [204, 205, 206], where the delta-function of gauge conditions determines the  $(n-m)$ -dimensional physical subspace  $\Sigma$  embedded in superspace and  $J(q, \partial S / \partial q)$  is a corresponding Faddeev-Popov determinant with the Hamilton-Jacobi value of the momentum<sup>10</sup>.

Thus, in the one-loop approximation we have a complete agreement between the reduced phase space (ADM) quantization and the BfV induced Dirac quantization method of Sect.7.4.. The same can be done for the semiclassical solution of quantum Dirac constraints. While here the two-point prefactor (7.115) was obtained by directly solving the quantum Dirac constraint, the same can be done by calculating in the one-loop approximation the path integral (7.102) in the relativistic gauge of the form  $\dot{N}^\mu - \chi^\mu(q, p) = 0$ . This was explicitly done in [226, 227] with the same result (7.115).

It remains to discuss the Hermiticity and gauge dependence properties of relevant observable matrix elements relative to this physical inner product. When it is rewritten in terms of the auxiliary inner product with the delta-function type *operator* measure

$$\langle \Psi' | \Psi \rangle_{\text{phys}} = \langle \Psi' | \hat{J} \delta(\hat{\chi}) | \Psi \rangle + O(\hbar), \quad (7.136)$$

the Hermiticity of the observable (7.107) with a real  $\mathcal{O}_I(q, p)$  immediately follows from the relation [209]  $\langle \Psi' | [\hat{\mathcal{O}}_I, \hat{J} \delta(\hat{\chi})] - i \hbar V_{I\lambda}^\lambda | \Psi \rangle = O(\hbar^2)$ . Thus,  $\hat{\mathcal{O}}_I$  is Hermitian with respect to  $\langle \dots | \dots \rangle_{\text{phys}}$ , even though it has an anti-Hermitian part,  $\hat{\mathcal{O}}_I^\dagger = \hat{\mathcal{O}}_I - i \hbar V_{I\lambda}^\lambda + O(\hbar^2)$  with respect to  $\langle \dots | \dots \rangle$ .

The gauge independence of the physical inner product is based, as shown in [206, 207], on the fact that it can be rewritten as the integral over  $(n-m)$ -dimensional surface  $\Sigma$  of a certain  $(n-m)$ -form which is closed in virtue of the

---

<sup>10</sup>In eq.(7.81) the product of two semiclassical wave functions  $\Psi'^* \Psi_1 = P'^* P \exp[i(S' - S)/\hbar]$  involves two different Hamilton-Jacobi functions, so that it seems ambiguous on which Lagrangian manifold ( $p = \partial S / \partial q$  or  $p = \partial S' / \partial q$ ) the relevant momentum argument of  $J(q, p)$  should be constructed. One should remember, however, that in semiclassical expansion the integral is calculated by the stationary phase method in which a dominant contribution comes from the stationary point satisfying  $\partial S_1 / \partial q = \partial S_2 / \partial q$ . This makes these Lagrangian surfaces to coincide in the leading order, their difference being treated perturbatively in  $\hbar$ .

Dirac constraints on physical states

$$\langle \Psi' | \Psi \rangle_{\text{phys}} = \int_{\Sigma} \omega^{(n-m)}, \quad d\omega^{(n-m)} = 0, \quad (7.137)$$

$$\omega^{(n-m)} = \frac{dq^{i_1} \wedge \dots \wedge dq^{i_{n-m}}}{(n-m)!} \epsilon_{i_1 \dots i_n} \Psi'^* \nabla_1^{i_{n-m+1}} \dots \nabla_m^{i_n} \Psi. \quad (7.138)$$

It follows then from the Stokes theorem that this integral is independent of the choice of  $\Sigma$  in the class of surfaces the cobordism between which is enforced by some regular  $(n-m+1)$ -dimensional subspace of the  $q$ -space.

The closure of  $\omega^{(n-m)}$  is a corollary of the approximate continuity equation for  $\Psi'^* \Psi$ ,  $\partial(\nabla_{\mu}^i \Psi'^* \Psi) / \partial q^i = U_{\mu\lambda}^{\lambda} \Psi'^* \Psi + O(\hbar)$ , where the correction  $O(\hbar)$  is due to the difference between the momentum arguments related to different Hamilton-Jacobi functions  $p' = \partial S' / \partial q$  and  $p = \partial S / \partial q$ , which goes beyond the one-loop approximation (see the footnote 9).

A similar gauge independence property holds for the matrix elements of physical observables, which only differ from (7.137) by the presence of  $\mathcal{O}_I(q, \partial S / \partial q)$  in the integrand, and follows from gauge invariance of  $\hat{\mathcal{O}}_I$ . Thus, as it should have been expected from the theory of gauge fields and BFV formalism the gauge independence of the physical matrix elements or expectation values of observables follows from the gauge invariance of the latter. This is true not only at the formal path-integral quantization level, but also in the operator Dirac quantization scheme.

The exterior form representation (7.137)-(7.138) of the physical inner product gives a correct understanding of the Wheeler-DeWitt conserved current (7.54) – this is a flux of the  $(n-m)$ -form through the  $(n-m)$ -dimensional surface  $\Sigma$  in  $q$ -space, defined by (7.119). In fact, this surface is a coordinate physical subspace embedded into the coordinate configuration space of  $q$ . The codimension of this surface is the range of the index  $\mu$  enumerating the constraints –  $m$ , therefore the flux conservation through such surface requires not one, but rather  $m$  continuity type equations. The analogy with the Klei-Gordon type current becomes stronger if we notice that each  $\nabla_{\mu}^i$  factor in the definition (7.138) of  $\omega^{(n-m)}$  can be represented as a result of action of the Wronskian type operator

$$\frac{1}{2} \left( \frac{\partial H_{\mu}}{\partial p_i} \Big|_{p=\vec{\hbar}\partial/i\partial q} - \frac{\partial H_{\mu}}{\partial p_i} \Big|_{p=\overleftarrow{\hbar}\partial/i\partial q} \right) + O(\hbar) \quad (7.139)$$

on the wave functions  $\Psi$  and  $\Psi'^*$ , with  $\hbar \overrightarrow{\partial} / i \partial q$  acting to the right on the factor  $\Psi$  and  $\hbar \overleftarrow{\partial} / i \partial q$  – to the left on the factor  $\Psi'^*$  [207]. For the Hamiltonian constraint  $H_{\mu}$  with  $\mu = (\perp, \mathbf{x})$  this is a first order differential Wronskian operator,

$$\frac{\hbar}{i} \left( G^{abcd}(\mathbf{x}) \frac{\overrightarrow{\partial}}{\delta g_{cd}(\mathbf{x})} - \frac{\overleftarrow{\partial}}{\delta g_{cd}(\mathbf{x})} G^{abcd}(\mathbf{x}) \right). \quad (7.140)$$

The product of  $m = 4 \times \infty^3$  such operator factors in (7.138) is what distinguishes Einstein field theory from the relativistic quantum mechanics of a single Klein-

Gordon equation corresponding to  $m = 1$ . Unfortunately, this analogy has not yet been extended beyond linear order in  $\hbar$  to get *exact* conservation law for a flux of many Klein-Gordon type equations subject to a commutator involution of the Dirac constraints type.

In a spatially closed gravitational system with  $H_0 = 0$  a time evolution arises due to the choice of explicitly time dependent gauge conditions, which according to (7.119) implies the motion of this physical subspace in super-space of 3-metrics  $q$ . The one-parameter family of such surfaces  $\Sigma(t)$  induces the nonvanishing values of lapse and shift functions  $N^\mu = -J_\nu^{-1\mu} \partial \chi^\mu / \partial t$  (cf. Eq.(3.15)), which according to (2.63) determine the “velocity” of the motion of the 3-dimensional spatial section in spacetime. This is the way how a “frozen” formalism becomes evolutionary due to the imposed time variation of gauge conditions. Such time evolution enters the construction of quantum averages via explicit time dependence of the operator measure in the physical inner product,  $\hat{M}_t = \delta(\chi(q, t)) \hat{J}_t + O(\hbar)$ ,  $\hat{J}_t \equiv \det[\chi^\mu(q, t), H_\nu(q, \hat{p})] / i\hbar$ . As a result, quantum averages of generic operators become  $t$ -dependent even though the Wheeler-DeWitt wave function  $\Psi(q)$  is constant in time. The nontrivial conservation in time of such averages for matrix elements between different Dirac states is the manifestation of unitarity that was checked above both at the reduced phase space quantization level and in the Dirac-Wheeler-DeWitt formalism.

## 7.6. Problems and prospects of quantum gravity and cosmology

Thus, the reduced phase space (ADM) quantization and quantum geometrodynamics – Dirac-Wheeler-DeWitt quantization, the latter generated by the truncation of the operator BFV formalism, form equivalent theories. The picture of this equivalence still suffers from a number of inconsistencies and unsolved issues. To begin with, unitary equivalence of these two schemes (7.133) is likely to be restricted to the one-loop approximation for semiclassical states. Apparently the most consistent and universal technique is the operator BFV/BRST quantization. Due to embedding of the theory into extended relativistic phase space of gauge and ghost fields it has the most efficient tools and a lot of gauge fixing flexibility for the analysis of gauge invariance properties, gauge dependence issues, etc. In this regard, the efficiency of the Dirac quantum geometrodynamical approach is much lower, and for the ADM quantization it is nearly absent at all in view of the necessity to solve explicitly complicated nonlinear constraint equations and to deal with spacetime nonlocality.

The problem of the choice of a good gauge-fixing procedure is another fundamental issue in quantum geometrodynamics. Note that, similarly to the Wronskian inner product of the Klein-Gordon equation, the flux of the current form (7.137)-(7.138) is not positive definite. At the classical level this is equivalent to the statement that the determinant factor  $J(q, p) = \det J_\nu^\mu(q, p)$  generally is

not positive definite either. But the degeneration of the Faddeev-Popov functional matrix  $J_V^\mu(q, p)$  means the breakdown of the gauge fixing procedure (the analogue of the Gribov copies problem in Yang-Mills gauge theories [228, 229]). The canonical gauges  $\chi^\mu(q, p)$  that would avoid this problem are, to the best of our knowledge, not known (the discussion of Gribov copies in cosmological context can be found in [230]). This would motivate in analogy with the second quantization of Klein-Gordon equation and secondary quantized string field theory the so called *third quantization* of gravity targeting once very popular speculations on the physics of multiple baby universes [231].

Another difficulty in quantum geometrodynamics is lacking manifest space-time covariance which is critical for the studies of ultraviolet divergences. The reader might have noticed that the superspace dimensionality and the number of spacetime diffeomorphisms was silently denoted by  $n$  and  $m$  which in pure 4-dimensional gravity are divergent quantities  $n = 6 \times \infty^3$  and  $m = 4 \times \infty^3$ , so that majority of constructions underlying the Wheeler-DeWitt equation are purely formal. Commutators of phase space coordinates and momenta in local quantum constraint are divergent  $\sim \delta^{(3)}(0)$ , traces of structure functions also represented pure divergences  $U_{\mu\nu}^\nu \sim \delta^{(3)}(0)$ , etc. All these expressions require regularization and renormalization, which is hard to render covariant in manifestly noncovariant formalism.

The path integral method which is a central point of the BFV quantization has an efficient means to overcome this difficulty – integration over phase space momenta converts the path integral to the Lagrangian form [213, 46] which in a special class of background covariant gauge conditions allows one to enjoy all the advantages of the manifestly covariant treatment. A covariant regularization allows one to minimize the violation of classical symmetries of the theory at the quantum level. In this respect, the path integral representation is what one needs – calculating this integral by the technique of covariantly regularized Feynman graphs is a right alternative to the attempts of directly solving the Wheeler-DeWitt equation. This really denies disparaging remarks on this equation, mentioned in Preface, as an absolutely useless tool in quantum gravity and explains its first principle founding nature. Like Schrödinger equation in quantum field theory, it lies at the foundation of the theory, though in concrete applications it gives way to a more efficient but a derivative tool – path integral method.

The realization of this strategy for the two point kernel (7.102) gives it as a projector onto the space of solutions of Wheeler-DeWitt equations, acting in superspace of 3-metrics and matter fields  $q^i = (g_{ab}(\mathbf{x}), \phi(\mathbf{x}))$ .

$$U[g_{ab}^+(\mathbf{x}), \varphi^+(\mathbf{x}) | g_{ab}^-(\mathbf{x}), \varphi^-(\mathbf{x})] = \int D[g_{\mu\nu}, \phi] e^{iS[g_{\mu\nu}, \phi]}. \quad (7.141)$$

Here  $S[g_{\mu\nu}, \phi]$  is the covariant action of the gravitational and matter fields, the details of Faddeev-Popov's gauge fixing procedure are hidden in the measure  $D[g_{\mu\nu}, \phi]$  and the integration runs over gravitational and matter spacetime histories  $q^i(t), N^\mu(t) = (g_{\mu\nu}(x), \phi(x))$ ,  $x = (t, \mathbf{x})$ , interpolating between config-

urations at initial and final spatial hypersurfaces (labeled by  $t_{\pm}$ ),

$$g_{ab}(\mathbf{x}, t_{\pm}) = g_{ab}^{\pm}(\mathbf{x}), \quad \phi(\mathbf{x}, t_{\pm}) = \varphi^{\pm}(\mathbf{x}). \quad (7.142)$$

Depending on the choice of these configurations and possible analytic continuation of the time integration contour between  $t_{-}$  and  $t_{+}$  into the complex plane (Euclidean time or “Euclidean quantum gravity”) this path integral generates various prescriptions for a cosmological quantum state. These prescriptions include, in particular, the Hartle-Hawking no-boundary [232, 233] and the tunneling [234, 235, 236, 237] quantum states of the Universe, which are usually associated with creation of the Universe from “Nothing”. The last two decades, with the advent of precision cosmology, are characterized by a growing interest in cosmological inflation theory that is widely recognized as underlying the observable large scale structure of the Universe. This explains the interest in these two quantum states as they, with this or that level of success, describe the origin of inflationary Universe.

Even more appealing sounds a recently suggested construction of the initial cosmological state as a microcanonical density matrix [238, 239]. This density matrix is proposed as a projector on all possible solutions of the Wheeler-DeWitt equation (7.141). The motivation for such a proposal is the principle of Occam razor – minimum set of assumptions, because this is an ultimate equipartition in the full set of states of the theory — “*Sum over Everything*” [239]. Creation of the Universe from “Everything” is conceptually more appealing than creation from “Nothing”, because the democracy of the equipartition better fits the principle of Occam’s razor than the selection of a concrete state.

The statistical sum of this microcanonical state is the trace of  $\hat{U}$  over the *physical* configuration space, which has a representation of the covariant path integral over periodic in time histories of gravitational and matter fields [210]

$$Z = \text{tr}_{\text{phys}} \hat{U} \equiv \int dq \hat{M} U(q, q') \Big|_{q'=q} = \int_{\text{periodic}} D[g_{\mu\nu}, \phi] e^{iS[g_{\mu\nu}, \phi]}. \quad (7.143)$$

Semiclassical calculation of this integral by the saddle point method runs in a manifestly covariant formalism which admits covariant UV renormalization underlying nontrivial quantum scaling behavior which in its turn is generated by a conformal anomaly of matter fields [238, 239].

This leads to numerous new physical effects in the theory of very early quantum Universe, which are impossible within the no-boundary and tunneling prescriptions. These effects culminate in the prediction of a new type of inflationary scenario with a bounded sub-Planckian energy scale [238, 241], which incorporates models of the Starobinsky  $R^2$ -inflation and Higgs inflation [240]. These models give, as is widely recognized, the best explanation of the observable cosmic microwave background (CMB) data and even establish intriguing relation between the basic parameters of this data and the mass value of the recently discovered Higgs boson [242, 243]. There is a lot more in store for us, which is ultimately based on the Dirac-Wheeler-DeWitt quantum geometrodynamics!

# Appendices

## A1. Geometry of manifolds

This appendix contains a brief summary of definitions of the basic concepts of the differential geometry of manifolds used in the main text. The details and proofs of facts collected here can be found in [2, 3, 4, 24].

### *Manifold*

Let  $M$  be a topological Hausdorff space with a countable base (the Hausdorff property means the separability of points in the space  $M$  in the following sense: for any  $p, q \in M$ , there are open neighbourhoods  $U_p \ni p$  and  $V_q \ni q$ , such that  $U_p \cap V_q = \emptyset$ ). The space  $M$  is called an  $n$ -dimensional smooth manifold, if it is locally homeomorphic to  $R^n$ , i.e., 1) for each open set  $U_i$  (the collection of which covers the whole space  $\cup_i U_i = M$ ), a homeomorphism  $\varphi_i : U_i \rightarrow R^n$  to a region of  $R^n$  is defined, which means the introduction of the local coordinates  $x^1(p), \dots, x^n(p)$  for the points  $p \in U_i$ ; 2) in the intersections  $U_i \cap U_j \neq \emptyset$ , the composition of maps  $\varphi_j \circ \varphi_i^{-1} : R^n \rightarrow R^n$  is defined which is a smooth function of its arguments. For simplicity, the smoothness is understood everywhere as an infinite differentiability. The pair  $(U_i, \varphi_i)$  is called a chart, and the complete set of consistent charts (in the sense of the smoothness of transition functions in the intersections) is called an atlas of the manifold  $M$ .

The local properties of manifolds are clarified when studying the tangent spaces. One can define the tangent vector  $X_p$  at the point  $p \in M$  as the linear differential operator acting on the set of smooth functions defined on the

manifold  $M$  in the neighbourhood of the point  $p$ , that satisfies the condition  $X(fg) = X(f)g(p) + f(p)X(g), \forall f, g$ . The set  $T_pM$  of all tangent vectors at the point  $p \in M$  constitutes an  $n$ -dimensional linear vector space and is called a tangent space to  $M$  at a point  $p$ . The operators  $(\partial_\mu)_p = (\partial/\partial x^\mu)_p$  [where  $x^\mu, \mu = 1, \dots, n$ , are the coordinates in the local chart and  $(\partial_\mu)_p f = (\partial_\mu f)_p$ ] constitute the basis in  $T_pM$ , which is called the coordinate basis. Any vector  $X_p$  is represented by the linear combination  $X = X^\mu(p)\partial_\mu$ .

Let  $M_1$  and  $M_2$  be the smooth manifolds. The mapping  $f : M_1 \rightarrow M_2$  is called a differentiable mapping, if in any local charts  $(U_i, \varphi_i)$  in  $M_1$  and  $(V_j, \psi_j)$  in  $M_2$  the composition  $\psi_j \circ f \circ \varphi_i^{-1}$  is a differentiable function. The differentiable mapping  $f$  determines the mapping of the respective tangent spaces: a tangent vector  $X_p$  at the point  $p \in M_1$  is mapped to the vector  $Y_p = f_*(X_p)$  at the point  $q = f(p) \in M_2$  in accordance with the rule  $f_*(X)(g) = X(g \circ f)$  for any smooth function  $g : M_2 \rightarrow R$ . The map  $f_*$  is called a differential of  $f$ ; this map is also often denoted as  $df$ .

We say that the smooth vector field  $X$  is defined on  $M$ , if each point  $p \in M$  is mapped to the vector  $X_p \in T_pM$  and for any smooth function  $f$  on  $M$   $X_p(f) = (Xf)(p)$  is a smooth function.

The space of linear forms  $T_p^*M$  dual to  $T_pM$  is called a cotangent space at a point  $p \in M$ . In the local coordinates  $\{x^\mu\}$  in the neighbourhood of  $p$ , the basis in  $T_p^*M$  can be defined by the set of differentials  $(dx^1, dx^2, \dots, dx^n)$ , the latter basis is dual to the coordinate frame basis  $\{\partial_\mu\}$ .

Using  $T_pM$  and  $T_p^*M$ , one can consider arbitrary rank tensors as the elements of the tensor products  $T_pM \otimes \dots \otimes T_pM \otimes T_p^*M \otimes \dots \otimes T_p^*M$ .

In the local chart  $\{x^\mu\}$ , any smooth tensor field  $T$  covariant of the rank  $k$  and contravariant of the rank  $l$  is defined by its components  $T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_k}(x)$  – the smooth functions of coordinates

$$T = T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_k} \partial_{\mu_1} \otimes \dots \otimes \partial_{\mu_k} \otimes dx^{\nu_1} \otimes \dots \otimes dx^{\nu_l}.$$

When changing the coordinates  $x^\mu \rightarrow y^\mu(x)$ , the tensor components are transformed as

$$T_{\nu_1 \dots \nu_l}^{\mu_1 \dots \mu_k} = T_{\beta_1 \dots \beta_l}^{\alpha_1 \dots \alpha_k} \frac{\partial y^{\mu_1}}{\partial x^{\alpha_1}} \dots \frac{\partial y^{\mu_k}}{\partial x^{\alpha_k}} \frac{\partial x^{\beta_1}}{\partial y^{\nu_1}} \dots \frac{\partial x^{\beta_l}}{\partial y^{\nu_l}},$$

in accordance with the transformations of bases of the tangent and cotangent spaces

$$\frac{\partial}{\partial y^\mu} = \frac{\partial x^\nu}{\partial y^\mu} \frac{\partial}{\partial x^\nu}, \quad dy^\mu = \frac{\partial y^\mu}{\partial x^\nu} dx^\nu.$$

### Differential forms

The elements of the cotangent space  $\omega \in T_p^*M$ , defined at each point  $p \in M$ , are called a smooth differential 1-form, if in the local coordinates  $\{x^\mu\}$  the decomposition coefficients of the form  $\omega$  with respect to the basis of  $T_p^*M$ ,  $\omega = \omega_\mu(x) dx^\mu$  are differentiable functions.

The exterior product of spaces  $T_p^*M$  is defined as a totally antisymmetrized tensor product

$$T_p^*M \wedge \cdots \wedge T_p^*M = \text{altern}(T_p^*M \otimes \cdots \otimes T_p^*M).$$

The elements of space  $\Lambda^k(T_p^*M)$  – of  $k$ -th exterior power  $T_p^*M$  – are the completely antisymmetric covariant tensors

$$\omega = \frac{1}{k!} \omega_{\mu_1 \dots \mu_k}(x) dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_k}, \quad \omega_{\mu_1 \dots \mu_k} = \omega_{[\mu_1 \dots \mu_k]}.$$

If at each point  $p \in M$ , the element from  $\Lambda^k(T_p^*M)$  is defined such that the coefficients  $\omega_{\mu_1 \dots \mu_k}(x)$  are smooth functions, we say that an exterior differential form of rank  $k$  is defined on the manifold  $M$ . The space of all  $k$ -forms on  $M_n$  is denoted by  $\Lambda^k$  (it is clear that  $\Lambda^k = \emptyset$ , if  $k > n = \dim M$ ). The direct sum  $\Lambda^*(M) = \Lambda^0 \oplus \Lambda^1 \oplus \dots \oplus \Lambda^n$  is a linear space (with an obvious definition of the sum and multiplication) and the introduction of the operation of exterior product in it converts  $\Lambda^*(M)$  into an algebra. In the local coordinates  $\{x^\mu\}$ , the exterior product of a  $k$ -form

$$\omega = \frac{1}{k!} \omega_{\mu_1 \dots \mu_k} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_k}$$

and an  $l$ -form

$$\varphi = \frac{1}{l!} \varphi_{\mu_1 \dots \mu_l} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_l}$$

is defined as a  $(k+l)$ -form  $\sigma = \omega \wedge \varphi$ ,

$$\sigma = \frac{1}{(k+l)!} \sigma_{\mu_1 \dots \mu_{k+l}} dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_{k+l}}$$

where

$$\sigma_{\mu_1 \dots \mu_{k+l}} = \frac{(k+l)!}{k!l!} \omega_{[\mu_1 \dots \mu_k} \varphi_{\mu_{k+1} \dots \mu_{k+l}]}$$

The exterior product is non-commutative:

$$\binom{(k)}{\omega} \wedge \binom{(l)}{\varphi} = (-1)^{kl} \binom{(l)}{\varphi} \wedge \binom{(k)}{\omega}.$$

For a differentiable mapping  $f : M_1 \rightarrow M_2$  along with its differential  $f_* : TM_1 \rightarrow TM_2$ , the dual mapping of forms (which called a pull-back map) is always determined: the form  $\varphi$  on  $M_2$  is mapped to the form  $\omega = f^*\varphi$  on  $M_1$ , so that  $f^*\varphi(X_1, \dots, X_k) = \varphi(f_*X_1, \dots, f_*X_k)$ , where  $X_1, \dots, X_k \in T_pM_1$ .

The operation of exterior differentiation is most important notion of the exterior algebra. The mapping  $d : \Lambda^k \rightarrow \Lambda^{k+1}$  is called the exterior differential, if it has the following properties: the Leibniz rule  $d(\omega^l \wedge \varphi) = d\omega \wedge \varphi + (-1)^l \omega \wedge d\varphi$  with  $\omega \in \Lambda^l$ ; for 0-forms (functions on  $M$ )  $d$  coincides with the differential  $df$ ; and  $d$  is nilpotent,  $d^2 = 0$ . In the local coordinates

$$d \binom{(k)}{\omega} = \frac{1}{k!} \partial_{[\mu_1} \omega_{\mu_2 \dots \mu_{k+1}]} dx^{\mu_1} \wedge dx^{\mu_2} \wedge \cdots \wedge dx^{\mu_{k+1}}.$$

If  $M$  is a Riemannian space with the metric  $g_{\mu\nu}(x)$  of the signature  $p$ , then the operation of dualization is defined on the exterior algebra, or the Hodge (“star”) operator  $*$ :  $\Lambda^k \rightarrow \Lambda^{n-k}$ . In the local coordinates, this mapping has the following form:

$$* \omega = \frac{1}{(n-k)!k!} \varepsilon_{\mu_1 \dots \mu_{n-k} \nu_1 \dots \nu_k} \omega^{\nu_1 \dots \nu_k} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{n-k}},$$

where  $\varepsilon_{\mu_1 \dots \mu_n}$  is totally antisymmetric Levi-Civita tensor density. From the definition, we have a property

$$** \omega = (-1)^{k(n-k)+p} \omega.$$

The theory of integration of exterior forms on manifolds can be studied in [3, 4]. Here, we will confine ourselves to the local definition only. Every smooth  $k$ -form  $\omega$  on the manifold  $M_n$  defines on any  $k$ -dimensional hypersurface  $N_k \subset M_n$  a  $k$ -form  $\gamma^* \omega$ , where  $\gamma: N_k \rightarrow M_n$  is the relevant embedding map. If the local coordinates  $(y^1 \dots y^k)$  are chosen in a region  $U \subset N_k$ , the integral of the form  $\omega$  over  $U$  is defined as the number  $\int_U (\gamma^* \omega)_{12 \dots k} dy^1 \dots dy^k$ . This number is denoted as  $\int_U \omega$ . The invariant integral over the entire  $k$ -hypersurface  $\int_N \omega$  is then determined using the natural matching of the integrals over the charts covering  $N_k$ .

## Bundles

The notion of a bundle space is very important in physics, because it formalizes the idea of an “internal space” of states in which a physical system can be in each point of the spacetime.

Let  $V_m$  be an  $m$ -dimensional vector space, and the action of some group  $G: G \times V^m \rightarrow V^m$  is defined on it. Then, the trivial vector bundle over  $M$  is called a direct product  $V^m \times M = E$ , where the action of the group  $G: G \times E \rightarrow E$  is naturally defined:  $u = (v, x) \in E$  is mapped into  $gu = (gv, x)$ . The manifold  $M$  is called a base of bundle  $E$ ,  $V^m$  is a typical fiber of a bundle, and  $G$  is a structural group. The generalization of this structure is a locally trivial bundle  $E(M, V, G, \pi)$ . By definition, this is an  $(m+n)$ -dimensional smooth manifold  $E$  where the projection  $\pi: E \rightarrow M$  is defined, and in addition there is a smooth atlas,  $\cup_i U_i = M$ , such that for any point  $x \in M$  there is a neighbourhood  $U_i \ni x$ , the preimage of which  $\pi^{-1}(U_i)$  is diffeomorphic to  $V \times U_i$  (the local triviality), and the corresponding diffeomorphisms  $\varphi_i: \pi^{-1}(U_i) \rightarrow V \times U_i$  are identical, i.e., for  $(v, x) \in V \times U_i$  one has  $\pi \circ \varphi_i^{-1}(v, x) = x$ . In addition, in the intersection regions  $U_i \cap U_j \neq \emptyset$ , the diffeomorphisms are compatible in the sense that  $\varphi_i^{-1} \circ \varphi_j = g \in G$  determines an automorphism of the fiber  $V$  under the action of the structural group  $G$ . If the smooth manifold  $V$  is a space of a linear representation of the group  $G$ , i.e., it is an  $m$ -dimensional vector space, the bundle  $E$  is called vector bundle.

An example of the vector bundle is the  $2n$ -dimensional manifold defined as  $\bigcup_{x \in M} T_x M = TM$  which is called a tangent bundle.

When the fiber coincides with the structural group  $V = G$  acting on itself by right shifts, the bundle is called a principal one. It denoted by  $P(M, G)$ . where  $P$  is a bundle space,  $M$  is its base. In a certain sense, the notion of a principal bundle generalizes the notion of the Lie group, since formally it is nothing but the collection of the copies of the group  $G$  “numbered” by the points from  $M$ . Its properties are described in details in [3]. Here we mention just one important mathematical fact. It is known that the Lie algebra  $\mathcal{G}$  generates an action on the Lie group  $G$ : namely, every element  $a \in \mathcal{G}$  generates a one-parametric subgroup in  $G$  by means of the exponential map  $\exp a$ . As a result, in the principal bundle  $P(M, G)$  the right action of  $G$  on  $P$  induces a natural homomorphism of the Lie algebra  $\mathcal{G}$  into the Lie algebra of the vector fields on  $P$ : for  $a \in \mathcal{G}$  the corresponding induced vector field  $a^*$  on  $P$  is called a fundamental vector field. By construction,  $a^*$  is tangent to the fiber, i.e., it is a vertical vector field,  $\pi_*(a^*) = 0$ .

Let the action of the group  $G$  be defined on a smooth manifold  $V$ . Then, for the principal bundle  $P(M, G)$  one can introduce an associated bundle with  $V$  as the typical fiber:  $E(M, V, G, P) = (P \times V)/G$ , where the factorization is considered with respect to the equivalence relation defined on  $P \times V$  by the natural right action  $G$ , namely,  $(p, v)g = (R_g p, g^{-1}v)$ ,  $\forall p \in P, v \in V, g \in G$ .

The example of the principal bundle is the bundle of linear frames  $L(M)$  with a structural group  $GL(4, R)$  (see Sec. 5.2.), and the tangent bundle can be considered as an associated one with it.

The smooth mapping  $\sigma : M \rightarrow E(M, V, G)$  such that  $\pi \circ \sigma = id_M$  is called the cross-section of the bundle  $E$ . The cross-section of the principal bundle exists only if  $P = M \times G$  is the trivial bundle [3, 4].

Let  $H$  be a subgroup of the group  $G$  and  $\tilde{P}(M, H)$  is a principal bundle over  $M$  with a structural group  $H$ . The mapping  $f : \tilde{P}(M, H) \rightarrow P(M, G)$  is called the reduction of the structural group  $G$  to the subgroup  $H$ , if the induced mapping  $f : M \rightarrow M$  is identical, and  $f : H \rightarrow G$  is a monomorphism. Then the subbundle  $\tilde{P}(M, H)$  is called the reduced bundle. One can prove that the necessary and sufficient condition for the reduction of the structural group  $G$  of the principal bundle  $P(M, G)$  to a closed subgroup  $H$  is the existence of a cross-section of the bundle  $E(M, G/H, G, P)$  associated with  $P$  with the typical fiber  $G/H$ , [3].

### *Connection*

The gauge field in the geometrical approach in the field theory is identified with a connection in the bundle space. Let  $P(M, G)$  be a principal bundle. The connection in  $P$  is a smooth mapping  $\rho_p$  of the tangent space to the base into the tangent space to the bundle,  $\rho_p : T_x M \rightarrow T_p P$  (with  $\pi(p) = x$ ) such that the horizontal subspace  $\mathcal{H}_p = \rho_p(T_x M)$  is invariant with respect to the action

of the group  $G$  on  $P$  and  $\pi_* \circ \rho_p = id_{T_x M}$ . The smoothness is understood as a smooth dependence of  $\rho_p$  on  $p \in P$ . Thus introduced connection defines the unique partition of the tangent space  $T_p P = \mathcal{H}_p \oplus V_p$  into the direct sum of the horizontal  $\mathcal{H}_p$  and the vertical  $V_p$  subspaces. The latter subspace  $V_p$  is formed by the tangent vectors to the fiber, and therefore, it is isomorphic to the Lie algebra of the group  $G$ . Observing this, one can give an equivalent definition of a connection. We say that the connection is defined in the bundle  $P(M, G)$  if a 1-form of  $\omega$  with the values in the Lie algebra of the structural group  $G$  is given on  $P$ , such that  $\omega(a^*) = a$  and  $R_g^* \omega = g^{-1} \omega g$  for all  $a \in \mathcal{G}$  and  $g \in G$ . One can show that definition of the connection 1-form  $\omega$  in  $P$  is equivalent to the existence of a set of 1-forms  $\omega_i$  on the base space  $M$ , which are determined for any  $i$  in the corresponding chart  $U_i$  (where  $\{U_i\}$  is an open covering of  $M$ ) by the local cross-sections  $\sigma_i : U_i \rightarrow P$ , so that  $\omega_i = \sigma_i^* \omega$ . Other properties of the connection are discussed in detail in [3, 4].

The gauge field strength coincides in the local coordinates with the components of the curvature of connection in the principal bundle. The form of the curvature  $R$  of a connection  $\omega$  on  $P$  is 2-form with the values in the Lie algebra of the structural group  $G$ , which is determined as exterior covariant differential from  $\omega$ :

$$R(X, Y) = D\omega(X, Y) = d\omega(hX, hY), \quad X, Y \in T_p P,$$

where  $h$  is a projection to the horizontal subspace  $\mathcal{H}_p$ . The curvature satisfies the structure equations [3, 4]

$$d\omega(X, Y) = -\frac{1}{2} [\omega(X), \omega(Y)] + R(X, Y), \quad X, Y \in T_p P.$$

Here the square bracket  $[ , ]$  denotes the Lie algebra commutator.

The definition of a connection  $\omega$  determines the parallel transport of the vectors in the bundle associated with  $P$ , as well as introduces the covariant derivative in  $E(M, V, G, P)$ , [3].

## A2. Spinor analysis on an arbitrary manifold

Let us consider the spinor bundle, the typical fiber of which is a two-dimensional complex vector space  $S_2$ , the structural group is  $GL(2, C)$ , and the base is a spacetime  $M_4$ . It is more convenient to work in the local chart, where the main object of our investigation – the contravariant spinor of the 1st rank is a two-component complex field

$$\xi^A = \begin{pmatrix} \xi^1(x) \\ \xi^2(x) \end{pmatrix}.$$

The spinor indices run the two values,  $A, B, C, \dots = 1, 2$ . When changing the local basis, the spinor is transformed as

$$\xi^{A'} = L^{A'}_A \xi^A, \quad L^{A'}_A \in GL(2, C).$$

As usual, the bilinear non-singular interior product ( $S_2 \times S_2 \rightarrow C$ ) is introduced in the space  $S_2$ , which in local basis is set by antisymmetric matrix [81, 82]

$$\varepsilon_{AB} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The elements of the dual spinor space – the covariant spinors ( $\xi_A$  in local basis) are transformed under the action of  $L^{-1} \in GL(2, C)$ . Raising and lowering of the indices can be performed with the help of  $\varepsilon_{AB}$  and  $\varepsilon^{AB}$ , where  $\varepsilon^{BA}\varepsilon_{BC} = \delta^A_C$ .

In addition, we define the bundle with typical fiber  $\overline{S}_2$  over  $M_4$ , which is called the conjugate spinor space. The elements of the fiber  $(\overline{S}_2)_x$  over the point  $x$  in the local chart are the complex-conjugate quantities to spinors from  $S_2$ . They are denoted by  $\Psi^{\dot{A}} = (\Psi^A)^*$ , and they are transformed under the action of the complex-conjugate matrix  $L^{\dot{A}}_{\dot{B}} = (L^A_B)^*$ . In  $\overline{S}_2$ , the bilinear non-singular antisymmetric interior product is also introduced in local basis by the matrix  $\varepsilon_{\dot{A}\dot{B}} = (\varepsilon_{AB})^*$ .

Spinors of the higher ranks are defined as the elements of the tensor products  $S_2 \otimes \dots \otimes S_2 \otimes \dots \overline{S}_2 \otimes \dots \overline{S}_2$  and are transformed in an obvious way as the products of the 2-spinors of the first rank.

The linear connection in the spinor bundle is defined by the corresponding 1-form with the values in the Lie algebra  $gl(2, C)$ . For a local cross-section – the spinor field  $\xi(x)$  – this introduces the coefficients of the local connection  $\omega_\mu(x)$  which under the change of the local basis are transformed according to  $\omega_\mu \rightarrow \omega'_\mu = L\omega_\mu L^{-1} + L\partial_\mu L^{-1}$ . Accordingly, the covariant derivative of spinor fields is defined by

$$D_\mu \xi^A_{\dot{B}} = \partial_\mu \xi^A_{\dot{B}} + \omega^A_{C\mu} \xi^C_{\dot{B}} - \omega^{\dot{D}}_{\dot{B}\mu} \xi^A_{\dot{D}}.$$

Such a general  $gl(2, C)$ -connection in the spinor bundle is not consistent with the interior spinor product in the sense that

$$D_\mu \varepsilon_{AC} = 2\omega_{[AC]\mu} = \varphi_\mu \varepsilon_{AC},$$

$\varphi_\mu = \omega^A{}_{A\mu}$  is the trace of the coefficients of the spinor connection.

Until now, the properties of the spinor bundle were considered separately from the geometry of the base space, since its own structure (the interior product, connection, etc.) can be defined quite independently. However, developing a consistent gauge approach to the gravitational field, we now will establish a relation of the spinor structure with geometrical spacetime structures. This issue is the most important one in the construction of the spinor analysis on an arbitrary manifold.

Let us make use of the fact that there exists a fundamental isomorphism of the space  $(S_2 \otimes \bar{S}_2)_x$  (the index  $H$  means the Hermitization) at a point  $x \in M_4$  and the tangent space  $T_x M_4$ . For the typical fibers  $S_2 \otimes \bar{S}_2$  and  $R^4$ , this isomorphism determines an arbitrary 4-vector as a linear combination of pairs  $\xi^A \eta^{\dot{B}}$ .

In the local chart, the aforementioned isomorphism is established with the help of the fundamental spin-tensor objects (the generalized Pauli matrices)  $g^a{}_{A\dot{B}} = -(g^a{}_{A\dot{B}})^\dagger$  (the Latin indices  $a, b, c, \dots = 0, 1, 2, 3$  refer to an arbitrary Lorentz frame in  $T_x M$ ). The matrices  $g^a{}_{A\dot{B}}$  satisfy the postulate

$$g^{aA\dot{B}} g^b{}_{C\dot{B}} = \eta^{ab} \delta_C^A + \frac{i}{2} \varepsilon^{abcd} g_c{}^{A\dot{B}} g_{dC\dot{B}}, \quad (\text{A2.1})$$

and are arbitrary in other respects. The generalized Pauli matrices can be chosen at one's convenience. We will use a particular representation:

$$g^{0A\dot{B}} = -i \mathbf{1}, \quad g^{kA\dot{B}} = -i \sigma^k, \quad k = 1, 2, 3,$$

where  $\mathbf{1}$  is the  $2 \times 2$  unit matrix, and  $\sigma^k$  are the standard  $2 \times 2$  Pauli matrices.

The important consequence of (A2.1) is the relation

$$g^{(a|A\dot{B}} |g^{b)}{}_{C\dot{B}} = \eta^{ab} \delta_C^A, \quad (\text{A2.2})$$

which provides a link between the world metric and the spinor interior product. As usual, their consistency is achieved with the help of the translational gauge gravitational field – the tetrads  $h_\mu^a$ , which introduce the world Pauli matrices  $g^\mu{}_{A\dot{B}} = h_\mu^a g^a{}_{A\dot{B}}$ , and hence obviously modifies (A2.1), (A2.2).

One can make the connections  $(\Gamma^a{}_{b\mu}$  and  $\omega^A{}_{B\mu})$  compatible by imposing a natural requirement of coincidence of the covariant derivative of a 4-vector with respect to the connection  $\Gamma$  on  $M_4$  and the covariant derivative with respect to  $\omega$  of its spinor image under the canonical isomorphism. Then we postulate

$$\nabla_\mu g_\nu{}^{A\dot{B}} = \frac{1}{2} K_{\mu\nu}{}^\lambda g_\lambda{}^{A\dot{B}}, \quad (\text{A2.3})$$

which is consistent with (A2.2).

The equation (A2.3) is easily solved with respect to the spinor connection by multiplying it by  $g^\nu{}_{C\dot{B}}$  and using the property (A2.1):

$$\omega^A{}_{B\mu} = \frac{1}{4} \Gamma_{[ab]\mu}{}^{(-)} S^{abA}{}_{B}, \quad (\text{A2.4})$$

where  $S^{abA}{}_{B}{}^{(-)} := g^{[a|A\dot{C}}|g^{b]}{}_{B\dot{C}}$ . The spin-tensor is symmetric  $S^{ab}{}_{AB} = S^{ab}{}_{BA}$  and anti-self-dual,  $*S^{ab}{}_{AB} = -iS^{ab}{}_{BA}$ . In a similar formula for  $\omega^{\dot{A}}{}_{B\mu}$ , instead of  $S^{(-)}$  one has the self-dual spin-tensor  $S^{ab\dot{A}}{}_{\dot{B}}{}^{(+)} := g^{[a|C\dot{A}}|g^{b]}{}_{C\dot{B}}$ .

The connection (A2.4) is determined up to a term  $i\Delta_{\mu}\delta_{\dot{B}}^{\dot{A}}$ , where  $\Delta_{\mu}$  is an arbitrary real vector, usually related to electromagnetic field, which was assumed to vanish.

Computing the commutator of the covariant spinor derivatives, one can obtain the expression for the spinor curvature

$$R^A{}_{B\mu\nu} = \partial_{\mu}\omega^A{}_{B\nu} - \partial_{\nu}\omega^A{}_{B\mu} + \omega^A{}_{C\mu}\omega^C{}_{B\nu} - \omega^A{}_{C\nu}\omega^C{}_{B\mu}$$

in the form

$$R^A{}_{B\mu\nu} = \frac{1}{4}R_{ab\mu\nu}(\Gamma)S^{(-)abA}{}_{B},$$

and similarly for  $R^{\dot{A}}{}_{\dot{B}\mu\nu}$ .

The equation (A2.4) shows that the Lorentz (the Riemann-Cartan) part of the connection  $\Gamma_{[ab]\mu}$  interacts with the spinors in a conventional minimal coupling manner, while the non-metricity is represented only by the Weyl vector  $K_{\mu}$ . This agrees with the fact that the structural group of the spinor bundle  $GL(2, C)$  is isomorphic to the Weyl group (which is the direct product of the Poincaré group  $\mathcal{P}_{10}$  by the 1-dimensional Abelian group of dilations  $R^1$ ), multiplied by the group of the phase transformations.

The transition from the 2-spinors  $\xi^A$  to the bispinors (or the 4-spinors of Dirac) is straightforward:

$$\Psi = \begin{pmatrix} \xi^A \\ \eta_{\dot{A}} \end{pmatrix}.$$

The  $4 \times 4$  Dirac matrices then read

$$\gamma^{\mu} = \begin{pmatrix} 0 & g^{\mu A\dot{B}} \\ g^{\mu}{}_{A\dot{B}} & 0 \end{pmatrix}.$$

With our choice of the generalized Pauli matrices, these Dirac matrices reproduce Weinberg's [192] representation. The Dirac matrices satisfy

$$\gamma^{\mu}\gamma^{\nu} = g^{\mu\nu}I + \frac{i}{2}\varepsilon^{\mu\nu\alpha\beta}\gamma_{\alpha}\gamma_{\beta}\gamma_5, \quad (\text{A2.5})$$

where  $\gamma_5 = \frac{-i}{4!}\varepsilon_{\alpha\beta\mu\nu}\gamma^{\alpha}\gamma^{\beta}\gamma^{\mu}\gamma^{\nu} = -i\hat{\gamma}^0\hat{\gamma}^1\hat{\gamma}^2\hat{\gamma}^3$ , and  $I$  is a  $4 \times 4$  unit matrix. From these fundamental relations, one straightforwardly derives

$$\begin{aligned} \gamma^{[\alpha}\gamma^{\beta]}\gamma^{\mu} - \gamma^{\mu}\gamma^{[\alpha}\gamma^{\beta]} &= 2(\gamma^{\alpha}g^{\beta\mu} - \gamma^{\beta}g^{\alpha\mu}), \\ \gamma^{[\alpha}\gamma^{\beta]}\gamma^{\mu} + \gamma^{\mu}\gamma^{[\alpha}\gamma^{\beta]} &= 2i\varepsilon^{\alpha\beta\mu\nu}\gamma_{\nu}\gamma_5. \end{aligned}$$

Finally, we define the covariant derivative of 4-spinors:  $D_\mu \Psi = \partial_\mu \Psi + \omega_\mu \Psi$ . Similarly to (A2.3), we impose the compatibility condition

$$D_\mu \gamma^\nu = \partial_\mu \gamma^\nu + \Gamma^\nu_{\lambda\mu} \gamma^\lambda + \omega_\mu \gamma^\nu - \gamma^\nu \omega_\mu = -\frac{1}{2} K_{\mu\lambda}{}^\nu \gamma^\lambda,$$

which is consistent with the symmetrized relation  $\gamma^{(\mu}\gamma^{\nu)} = g^{\mu\nu} I$ , see (A2.5). Note that  $\nabla_\lambda g^{\mu\nu} = -g^{\mu\alpha} g^{\nu\beta} K_{\lambda\alpha\beta} = -K_{\lambda}{}^{\mu\nu}$ . The solution of the compatibility condition is straightforward and represents a 4-spinor generalization of the connection (A2.4)

$$\omega_\mu = \frac{1}{4} \Gamma_{[ab]\mu} \gamma^{[a}\gamma^{b]}. \quad (\text{A2.6})$$

Hence, the covariant derivative of 4-spinor reads

$$D_\mu \Psi = \partial_\mu \Psi + \omega_\mu \Psi = \partial_\mu \Psi + \frac{1}{4} \Gamma_{[ab]\mu} \gamma^{[a}\gamma^{b]} \Psi. \quad (\text{A2.7})$$

The covariant derivative of the Dirac conjugated spinor  $\bar{\Psi} = \Psi^\dagger \beta$  has the following form:

$$D_\mu \bar{\Psi} = \partial_\mu \bar{\Psi} - \bar{\Psi} \omega_\mu = \partial_\mu \bar{\Psi} - \frac{1}{4} \Gamma_{[ab]\mu} \bar{\Psi} \gamma^{[a}\gamma^{b]}. \quad (\text{A2.8})$$

It is worthwhile to note that for the metric with the signature  $(-, +, +, +)$  the Dirac conjugation is defined by the matrix  $\beta = i\gamma^{\hat{0}}$ , and the Dirac matrices are anti-Hermitian in the sense [192]

$$\beta \gamma^{a\dagger} \beta = -\gamma^a.$$

In Weinberg's representation we explicitly have

$$\beta = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix},$$

and hence  $\beta^2 = I$ .

## A3. Basic notations

We summarize here the basic notations and conventions used in the book.

### *Indices*

Greek indices  $\alpha, \beta, \dots = 0, 1, 2, 3$  refer to the local world coordinates on the spacetime manifold. Latin indices  $a, b, c, \dots = 0, 1, 2, 3$  label the local Lorenz (tetrad) components, except for the Chapters 2, 3, and 7, where the indices  $a, b, c, \dots = 1, 2, 3$  refer to the 3-dimensional (spatial) coordinates. Separate local Lorentz components are denoted with hats over the index, to distinguish them from the world components. E.g.,  $v^{\hat{0}}$  is the 0-th component of a vector with respect to the orthonormal tetrad, in contrast to the 0-th component  $v^0$  in the coordinate basis. The upper case Latin indices  $A, B, C, \dots = 1, 2$  label the spinor components.

### *Metric*

The spacetime metric in 4 dimensions has the signature  $(-, +, +, +)$ , unless the other signature is assumed. Its determinant is  $g \equiv |^4g| = -\det g_{\mu\nu}$ . In Chapters 2, 3, and 7, the determinant of the 3-dimensional (spatial) metric is denoted as  $g^{1/2} = (\det g_{ab})^{1/2}$ . The totally antisymmetric Levi-Civita tensor density is  $\varepsilon_{\mu\nu\alpha\beta} = \sqrt{g}\eta_{\mu\nu\alpha\beta}$ , where the completely antisymmetric symbol has the only nontrivial component  $\eta_{0123} = +1$ . The scalar product for the spinors is defined by  $\varepsilon_{AB} = -\varepsilon_{BA}$ .

### *Derivatives*

The partial derivatives with respect to the spacetime coordinates are denoted  $\frac{\partial}{\partial x^\mu} = \partial_\mu = ,_\mu$ . The Riemannian covariant derivatives (defined by the Christoffel symbols) are denoted as  $;\mu$  or  $\overset{\{ \}}{\nabla}_\mu$ . The general covariant derivatives with respect to an arbitrary linear connection  $\Gamma$  are denoted as  $\nabla, \nabla^\Gamma, D^\Gamma$ . For the non-Riemannian objects, constructed from  $\Gamma$ , the functional dependence is always explicitly specified, for example  $R(\Gamma)$ .

## A4. Comments on the literature and general remarks

### Chapter 1

The equations of the general relativity theory of gravitation (GR) in the Riemannian spacetime were postulated by A. Einstein [22] in 1915, and simultaneously they were derived by D. Hilbert [23] on the basis of the principle of least action. In 1922, É. Cartan [20] introduced the torsion into the differential geometry. His studies of the manifolds with torsion were motivated by the interest in GR. Apparently, being under the influence of the results of the Cosserats brothers [25] on a continuous medium as a manifold, Cartan proposed [19] to couple the torsion tensor to an internal angular momentum of the matter. Later, the ideas of Cosserats-Cartan formed the basis of the continuous theory of dislocations [29].

In the gravity theory, these notions received a substantiation only after the gauge approach was formulated (H. Weyl [5], C.N. Yang and R. Mills [30], R. Utiyama [31]). The physical ideas that underlie the gauge theory about the existence of the fundamental matter fields and the fields mediating interactions became the basis of R. Utiyama's attempt to consider the gravitational field as a gauge one for the Lorentz group  $SO(3,1)$  (see also [32]). In his work, the Lorentz connection for the first time appeared as the corresponding gauge potential. However, the weak point in the approach of R. Utiyama was the introduction of the tetrads (metric) beyond the gauge scheme and an ad hoc requirement of the absence of the torsion. Eventually, these shortcomings were recognized. D. Sciama [33] dropped the condition of the vanishing torsion tensor. K. Hayashi [34] and B. De Witt [35] pointed out that the metric could be associated with a group of the spacetime translations  $T_4$  (more precisely, with the general coordinate transformations, understood as translations), see also [13, 37]. T. Kibble [36] considered the Poincaré group  $P_{10} = SO(3, 1) \rtimes T_4$  as a fundamental spacetime symmetry, which allowed him to give a gauge interpretation of the connection and the metric, cf. also [38]. Taking the Hilbert-Einstein linear in the curvature Lagrangian as a function of the independent metric and connection, D. Sciama and T. Kibble derived the field equations of the gauge  $P_{10}$ -theory of gravity, later named by A. Trautman [6] as the Einstein-Cartan theory (ECT). The thorough analysis of this theory is presented in the papers of F. Hehl [7] and A. Trautman [6] (see also [15, 16]).

### Chapter 2

The canonical formalism of systems with singular Lagrangians was constructed by P. Dirac [40]. He was also the first who applied this formalism to the gravitational field and reduced the Hilbert-Einstein gravitational action of the canonical form [40]. The Hamiltonian formalism of the gravitational field was thoroughly studied in the papers of R. Arnowitt, S. Deser, and C. Misner [41]. The same program was realized by K. Kuchař [42, 43, 44] on the invariant

language making use of the system of arbitrary space-like hypersurfaces, also extending the canonical formalism to the case of arbitrary matter field minimally interacting with the gravitation. In our presentation of the canonical formalism of gravitating systems, we followed the works of this author.

The proof of the degeneracy of the Lagrangian for the action invariant with respect to the local group, was borrowed from the lecture of V. P. Frolov at the Lebedev Physical Institute of the Russian Academy of Sciences.

The relation of the group of invariance of the action in the configuration space with the corresponding system of canonical transformations in the phase space, forming the “pseudogroup”, was studied in the papers of E. S. Fradkin, G. A. Vilkovisky [45, 46], and I. A. Batalin [47]. The same was done in the paper of C. Teitelboim [48] who discussed the general coordinate invariance group for the action of the physical fields in the curved spacetime.

### Chapter 3

The systematic method of the choice of dynamically independent degrees of freedom of the gravitational field was constructed by R. Arnowitt, S. Deser, and C. Misner [41], and subsequently generalized by K. Kuchař [44, 52] to the case of arbitrary gravitating systems with the matter sources.

An important difference of the dynamics of the asymptotically flat open world from the closed worlds was emphasized by De Witt [53], who attracted attention to the significance of the surface term in the Hamiltonian of the asymptotically flat gravitating world. The role of this surface integral in the Einstein theory was also discussed by T. Regge and C. Teitelboim [51], who connected it with the generator of the time translations at a spatial infinity.

The dynamics of the linearized gravitational field presented here has been considered by ADM [41], which identified the two transverse traceless gravitational wave modes and constructed the Hamiltonian of the gravitational radiation.

### Chapter 4

É. Cartan [79] was the founder of the spinor theory. B. Wan der Waerden and L. Infeld [80] introduced the 2-spinors. H. Weyl [84], V. A. Fock, and D. D. Ivanenko [85] developed a framework for the spinor analysis in the Riemannian space  $V_4$ . The expressions obtained by them for the spin connections entered the literature as the Fock-Ivanenko coefficients. The Dirac equation in the Riemann-Cartan spacetime  $U_4$  was first considered by H. Weyl [83]. V. I. Rodichev [55] demonstrated that the Dirac equation (with the zero mass) in the Minkowski space  $\mathcal{M}_4$  with torsion becomes a non-linear spinor equation of Ivanenko-Heisenberg [56] type (that underlies the unified theory of matter developed by W. Heisenberg [57]). This result was generalized by T. Kibble [36] to the spaces  $U_4$ . The spinor analysis in the space  $G_4$  was constructed in [18]. Detailed presentations of the spinor formalism can be found in [81, 82].

The question, whether the torsion field interacts with vector fields is still open in the modern literature. A. Trautman [6] and F. Hehl [7] made a convincing observation that the massless vector field should not interact with the torsion since such a coupling would violate the gauge invariance (see also [89]). Nevertheless, in the book we decided to demonstrate that the opposite point of view has some interesting consequences, caused by this interaction. In particular, the non-linear nature of the vector field equations, arising due to the ( $A$ - $Q$ )-interaction, leads to the existence of the soliton solutions [63].

The conformal transformations introduced in Sec. 4.2., were also considered independently in [90, 91].

The first non-singular cosmological solutions in the ECT were obtained in [88, 123, 174]. The analysis of different types of spatially homogeneous cosmological models with the Weyssenhoff fluid is given in [175]. The analogy between the torsion and the cosmological  $\Lambda$ -term was indicated in [176, 177].

The first correct formulation of the problem of the particle production was developed for cosmology in GR by A. A. Grib, S. G. Mamaev, V. M. Mostepanenko [61], L. Parker [76], Ya. B. Zeldovich, A. A. Starobinsky [77]. We follow [61] in the presentation of the production of the scalar particles in the  $U_4$  space-time.

## Chapter 5

The work of T. Kibble [36] is one of main contributions in the framework of so-called physical approach, where the direct localization of the global transformation of some group makes it necessary to “compensate” the corresponding non-covariance of the ordinary derivative by the introduction of the gauge field. Later works in this direction essentially were focused on the finding of the “true” gauge group for the theory of gravity, leaving the main structure of the physical approach unchanged (in the work [94], one can find an incomplete list of the groups and space-time structures). The criticism of this approach is contained in [96]. Along with this, the geometrical approach was developed on the basis of an idea of identifying the potential of the gauge field with a connection on some bundle over the spacetime. For the Yang-Mills theory, this fact was first noticed in [95] and later it was fully developed in investigations of A. Trautman [97]. A somewhat intermediate position between the physical and the geometrical approaches is held by the direction, developed by the group of F. Hehl [7, 98], which is based not so much on the gauge concept but on the fundamental link of the local invariance principle with the conservation laws. The latter studies established a remarkable fact (see also [38]): the source of the gauge field of the group  $G$  is the canonical Noether current corresponding to  $G$ -invariance of the theory.

The current status of the gauge theory of gravity is characterized by the application of the modern differential-geometric and topological methods for the study of their structure. The connection structure in the bundle of affine frames

presented in this book was first proposed in [99, 100], as well as independently developed in several different formulations in [101]-[105].

The idea to relate the existence of the gravitational field to the spontaneous breaking of a spacetime symmetry was in different forms expressed by many authors [97], [106]-[111]. The nonlinear realizations were introduced earlier in the study of the phenomenological Lagrangians in the theory of chiral symmetries [112, 113, 114]. The analysis of the dynamics of the non-linear gauge fields was considered in the paper of Y.M. Cho [116], and the authors followed this work in the description of spontaneous breaking in the Higgs mechanism.

## Chapter 6

The Lagrangians and the theories based on them considered in Sec. 6.1. were studied in the following papers: (b) – in [128] (and more consistently in [129]); (d) see also [129]; (e) – in [130]. The general Lagrangians constructed as a sum of all possible terms quadratic in the curvature and the torsion (with arbitrary coefficients) were studied in [131, 132, 133, 138]. Some authors considered the possibility to restrict the choice of the coupling coefficients by attracting additional physical and geometrical ideas such as the requirement of the absence of the “ghost” and tachyon poles in the tree-level propagator [134, 135], the validity of Birkhoff’s theorem [136], the correspondence with GR [131], etc.

In Sec. 6.3., when studying the correspondence of the  $\mathcal{P}_{10}$ -theory with GR, we used the method applied in [137] to the Lagrangian of a particular form. For a review of the studies of the issue of the cosmological constant in the gravity and supergravity, see in [139].

The gravitational instantons (such as Euclidean Schwarzschild, Taub-NUT,  $CP_2$ ) in the Riemannian gravity theory are studied in [142, 143, 144].

## Chapter 7

The method of canonical quantization of the degenerate systems and, in particular, of the gravitational field was pioneered by P. A. M. Dirac [40], B. S. DeWitt [53] L. D. Fadeev and V. N. Popov [154], E. S. Fradkin and G. A. Vilkovisky [45]. The path integral technique was introduced into the quantum theory in the works of Feynman [162]. The method of the covariant quantization of the gravitational field is studied in detail in [45, 46, 154, 157, 158, 159]. Many papers are dedicated to the quantization of the cosmological systems in the ADM formalism. We mention just a few of them, which more clearly characterize the main features of this approach [162, 163, 165, 178]. In the majority of works, the study of the quantum dynamics was conducted in the framework of the model assumptions about cosmology. For example, it was common to assume homogeneity, or a certain symmetry, to take the majority of the modes of the physical fields as frozen, to neglect them on one or another stage of evolution, etc.

A. Peres [166] introduced the equations of the Einstein-Hamilton-Jacobi theory as a link between the classical GR and the quantum geometrodynamics.

Quantum geometrodynamics was developed in the works of P. A. M. Dirac [40], B. S. DeWitt [53], and J. A. Wheeler [167]. The attempt to construct the path integral in the quantum geometrodynamics was made by H. Leutwyler [169], who, in particular, was the first to propose the correct local measure for the quantum gravity. The method of the path integration in the quantum geometrodynamics was constructed in [168, 204]. Old application of the quantum geometrodynamical method in special models can be found in very many works of which we mention only a few: [165, 173, 179]-[182]. A special mention deserve the attempts of solving the Wheeler-DeWitt equation nonperturbatively [117]-[119] in the approximation of the so-called ultralocal state in the quantum gravitodynamics [117].

The Batalin-Fradkin-Vilkovisky (BFV) path integral and operator quantization of constrained systems was developed in [214, 215, 216, 217, 218, 219, 220] starting with the introduction of the relativistic phase space and the nilpotent BRST operator acting in their representation space [213, 46]. The Batalin-Marnelius procedure of gauging out the BRST symmetry in the physical subspace of this space was built in [221, 222, 223] and used for the construction of the inner product in quantum geometrodynamics in [204, 206, 208, 207, 210]. These works also convey the consistency between the reduced (physical) phase space quantization and the Dirac quantization in generic constrained systems and gravity theory, in particular. This consistency is demonstrated explicitly in the one-loop order of the semiclassical expansion in [205, 206, 207, 208, 209], where also the operator realization of quantum Dirac constraints is derived in the subleading (one-loop) order in the Planck constant.

Widely acclaimed prescriptions for the quantum state of the Universe in the form of the no-boundary or tunneling wavefunction can be found in [232, 233, 234, 235, 236, 237]. Recent applications of the path integral method in quantum geometrodynamics are the construction of the microcanonical density matrix of the early Universe [239], the calculation of its statistical sum for models dominated by conformal quantum fields, which suppresses to zero in the ensemble the contribution of the Hartle-Hawking no-boundary state [238] and imposes an upper subplanckian bound on the energy scale of the new type of (hill-top) inflationary scenario [240, 241].

### Further reading

Two books by M. Blagojević and F. Hehl [200, 201] on the gauge gravity were published since the first Russian edition of our monograph. They give a broad overview of the history and of the recent developments, providing a complementary discussion of both mathematical and physical aspects of the subject. In an attempt to fill all the gaps and omissions that reflect our selection of the topics for the current book, we compiled an essentially complete bibliography on the gauge approach in gravity and related issues. The alphabetically ordered list can be found at the end of this book on page 201.

# References

- [1] *Quantum Theory of Gauge fields*, ed. N. P. Konopleva (Mir Publ.: Moscow, 1977) 436 p., in Russian.
- [2] Dubrovin B.A., Novikov S.P., and Fomenko A.T., *Modern Geometry – Methods and Applications* (Springer: New York, 1985-1992).
- [3] Kobayashi S. and Nomizu K., *Foundations of Differential Geometry*, Vol. 1 (Interscience Publishers: New York, 1963).
- [4] Lichnerowicz A., *Global theory of connections and holonomy groups* (Springer: Netherlands, 1976).
- [5] Weyl H., *Raum-Zeit-Matterie* (Springer: Berlin, 1920).
- [6] Trautman A., *On the structure on the Einstein-Cartan equation*, *Symposia Mathematica* **12** (1973) 139-160.
- [7] Hehl F.W., von der Heyde P., Kerlick G.D., and Nester J.M., *General relativity with spin and torsion: foundation and prospects*, *Revs. Mod. Phys.* **48** (1976) 393-416.
- [8] Rodichev V.I., *Gravitation Theory in the Orthogonal Frame* (Nauka: Moscow, 1974) in Russian.
- [9] Tonnelat M. A., *Les théories unitaires de l'électromagnétisme et de la gravitation* (Gauthier-Villars: Paris, 1965).

- [10] Courant R. and Hilbert D., *Methods of Mathematical Physics*, Vol. 1 (Interscience Publishers: New York, 1953).
- [11] Ponomarev V.N. and Tseytlin A.A., *On variational principle in general relativity*, *Vestnik. Mosk. Univ., ser. fiz., astr.*, **20**, n. 5 (1979) 68-80 (in Russian).
- [12] Minkevich A.V. and Karakura F.J., *On the relativistic dynamics of spinning matter in space-time with curvature torsion*, *J. Phys.* **A16** (1983) 1409-1418.
- [13] Cho Y.M., *Einstein Lagrangian as the translational Yang-Mills Lagrangian*, *Phys. Rev.* **D14** (1976) 2521-2525.
- [14] Gibbons G. W. and Hawking S. W., *Action integrals and partition functions in quantum gravity*, *Phys. Rev.* **D15** (1977) 2752-2756.
- [15] Kopczyński W., *The Palatini principle with constraints*, *Bull. Acad. Polon. Sci., sér. sci. math., astr., phys.*, **23** (1975) 467-473.
- [16] Ponomarev V.N. and Tseytlin A.A., *On the correct usage of the Palatini principle in the gravitational theories*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 6 (1978) 57-59 (in Russian).
- [17] Safko J.L. and Elston F., *Lagrange multipliers and gravitational theory*, *J. Math. Phys.* **17** (1976) 1531-1537.
- [18] Ponomarev V.N. and Obukhov Yu.N., *The generalized Einstein-Maxwell theory of gravitation*, *Gen. Relat. Grav.* **14** (1982) 309-330.
- [19] Cartan É., *Sur les variétés à connexion affine et la théorie de la relativité généralisée, 2 partie*, *Ann. Éc. Norm.* **42** (1925) 17-88.
- [20] Cartan É., *Sur une généralisation de la notion de courbure de Riemann et les espaces à torsion*, *C.R. Acad. Sci. (Paris)* **174** (1922) 593-595; English translation: *On a generalization of the notion of Riemann curvature and spaces with torsion*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979) /Eds. P.G.Bergmann and V.De Sabbata (Plenum: New York, 1980) 489-491.
- [21] Arkuszewski W., Kopczyński W., and Ponomarev V.N., *Matching conditions in the Einstein-Cartan theory of gravitation*, *Commun. Math. Phys.* **45** (1975) 183-191.
- [22] Einstein A., *Zur allgemeinen Relativitätstheorie*, *Königlich Preußische Akademie der Wissenschaften (Berlin). Sitzungsberichte* (1915) 778-786; *Zur allgemeinen Relativitätstheorie (Nachtrag)*, *Königlich Preußische Akademie der Wissenschaften (Berlin). Sitzungsberichte* (1915) 799-801.

- [23] Hilbert D., *Die Grundlagen der Physik (Erste Mitteilung)*, Nachrichten. Königlich Gesellschaft Wiss. Gottingen, Math.-phys. Klasse, Heft 3 (1915) 395-405.
- [24] Schouten J.A., *Tensor analysis for physicists* (Clarendon Press: Oxford, 1951).
- [25] Cosserat E. and Cosserat F., *Theorie des corps deformables* (Hermann: Paris, 1909).
- [26] Gambini R. and Herrera L., *Einstein-Cartan theory in the spin formalism*, *J. Math. Phys.* **21** (1980) 1449-1454.
- [27] Landau L.D. and Lifshitz E.M., *The classical theory of fields. Course of Theoretical Physics*, Vol. 2, 4th Engl. ed. (Butterworth-Heinemann: Amsterdam, 1980).
- [28] Palatini A., *Deduzione invariantiva delle equazioni gravitazionali del principio di Hamilton*, *Rend. Circ. Mat. Palermo* **43** (1919) 203-212; English translation: *Invariant deduction of the gravitational equations from the principle of Hamilton*, in: "Proc. 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979) /Eds. P.G.Bergmann and V.De Sabbata (Plenum: New York, 1980) 477-488.
- [29] Kröner E., *Algemeine Kontinuums theorie der Versetzungen und Eigenspannungen*, *Arch. Rat. Mech. Anal.* **4** (1960) 273-334.
- [30] Yang C. N. and Mills R. L., *Conservation of isotopic spin and isotopic gauge invariance*, *Phys. Rev.* **96** (1954) 191-195.
- [31] Utiyama R., *Invariant theoretical interpretation of interaction*, *Phys. Rev.* **101** (1956) 1597-1607.
- [32] Brodskii A.M., Ivanenko D.D., and Sokolik H.A., *New treatment of the gravitational field*, *ZhETF* **41** (1962) 1307-1309. (in Russian).
- [33] Sciama D.W., *The analogy between charge and spin in general relativity*, in: "Recent Developments in General Relativity" (Pergamon Press and PWN: Oxford and Warsaw, 1962) 415-439.
- [34] Hayashi K., *The gauge theory of the translation group and underlying geometry*, *Phys. Lett.* **B69** (1977) 441-444.
- [35] De Witt B.S., *Dynamical theory of groups and fields* (Gordon and Breach: New York, 1965) 248 p.
- [36] Kibble T.W.B., *Lorentz invariance and the gravitational field*, *J. Math. Phys.* **2** (1961) 212-221.

- [37] Minkevich A.V., *Gravitational field and the local invariance principle*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 4 (1966) 117-118 (in Russian).
- [38] Frolov B.N., *Principle of local invariance and Noether theorem*, *Vestnik Mosk. Univ., Ser. fiz., astron.*, n. 6 (1963) 48-58 (in Russian).
- [39] Einstein A., *Geometrie und Erfahrung*, *Preußische Akademie der Wissenschaften (Berlin). Sitzungsberichte* (1921) 123-130.
- [40] Dirac P. A. M., *Lectures on Quantum Mechanics* (Dover: Minneola, 2001).
- [41] Arnowitt R., Deser S., and Misner C. *The dynamics of general relativity*, in: "Gravitation, an Introduction to current research", Ed. L. Witten (J. Wiley and Sons: New York, 1963) 227-265.
- [42] Kuchař K., *Geometry of hyperspace. I*, *J. Math. Phys.* **17** (1976) 777-791.
- [43] Kuchař K., *Kinematics of tensor fields in hyperspace. II*, *J. Math. Phys.* **17** (1976) 792-800.
- [44] Kuchař K., *Dynamics of tensor fields in hyperspace. III*, *J. Math. Phys.* **17** (1976) 801-820.
- [45] Fradkin E. S. and Vilkovisky G. A., *S Matrix for Gravitational Field. II. Local Measure; General Relations; Elements of Renormalization Theory*, *Phys. Rev.* **D8** (1973) 4241-4285.
- [46] Fradkin E. S. and Vilkovisky G. A., *Quantization of relativistic systems with constraints. Equivalence of canonical and covariant formalism in quantum theory of gravitational field*. Preprint CERN No. Th-2332 (1977. Geneva) 53 pp.
- [47] Batalin I.A., *Quasigroup Structure and Constraints of I Kind*, Preprint FIAN USSR No. 157 (Moscow, 1979).
- [48] Teitelboim C., *How commutators of constraints reflect the spacetime structure*, *Ann. Phys.* **79** (1973) 542-557.
- [49] Denisov V.I. and Logunov A.A., *The inertial mass defined in the general theory of relativity has no physical meaning*, *Theor. Math. Phys.* **15** (1982) 421-426.
- [50] Klotz F.S., *Simplification of the constraints in the Dirac formalism for general relativity*, *Int. J. Theor. Phys.* **6** (1972) 251-257.
- [51] Regge T. and Teitelboim C., *Role of surface integrals in the Hamiltonian formulation of general relativity*, *Ann. Phys.* **88** (1974) 286-318.
- [52] Kuchar K., *A bubbletime canonical formalism for geometrodynamics*, *J. Math. Phys.* **13** (1972) 768-781.

- [53] De Witt B.S., *Quantum theory of gravity. I. The canonical theory*, *Phys. Rev.* **160** (1967) 1113-1148.
- [54] Weyssenhoff J. and Raabe A., *Relativistic dynamics of spin-fluids and spin-particle*, *Acta Phys. Pol.* **9** (1947) 7-18.
- [55] Rodichev V.I., *Space with torsion and nonlinear field equations*, *ZhETF* **40** (1961) 1469-1472 (in Russian) [*Sov. Phys. JETP* **13** (1961) 1029-1031].
- [56] Ivanenko D., *Bemerkungen zur theorie der wechselwirkung*, *Physikalische Zeitschrift der Sowjetunion* **13** (1938) 141-150.
- [57] Heisenberg W., *Introduction to the Unified Field Theory of Elementary Particle* (Interscience Publishers: New York, 1966) 192 p.
- [58] Krechet V.G. and Ponomarev V.N., *Nonlinearity and torsion*, *Theor. Math. Phys.* **24** (1975) 1036-1038 [*Teor. Mat. Fiz.* **24** (1975) 141-144 (in Russian)].
- [59] Isham C.J., Salam A., and Strathdee J., *f-Dominance of gravity*, *Phys. Rev.* **D3** (1971) 867-897.
- [60] Ponomarev V.N. and Pronin P.I., *Spontaneous symmetry breaking by the torsion field and nonlinear spinor theorems*, *Sov. J. Phys.* **22** (1979) 1324-1325 [*Izvestiya VUZov, Fizika*, n. 12 (1979) 81-82 (in Russian)].
- [61] Grib A.A., Mamayev S.G., and Mostepanenko V.M., *Quantum Effects in Strong External Fields* (Atomizdat: Moscow, 1980) 295 pp. In Russian.
- [62] Ponomarev V.N. and Smetanin E.V., *New version of nonlinear electrodynamics. I.*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 5 (1978) 29-37 (in Russian).
- [63] Ponomarev V.N. and Smetanin E.V., *New version of nonlinear electrodynamics. II.*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 6 (1978) 3-8 (in Russian).
- [64] Glashow S. L. and Gell-Mann M., *Gauge theories of vector particles*, *Ann, Phys.* **15** (1961) 437-460.
- [65] Urbakh V.Y., *Generalized theory of vector fields*, *Dokl. Akad. Nauk SSSR* **101** (1955) 1043-1046 (in Russian).
- [66] Markov M.A., *Elementary particles of maximally large masses (quarks and maxions)*, *Sov. Phys. JETP* **24** (1967) 584-592 [*ZhETF* **51** (1966) 878-890 (in Russian)].
- [67] Tunyak V.N., *Central-symmetric static field of a charged mass in an affine space with the Riemannian metric and torsion*, *Izvestiya AN BSSR, ser. fiz.-mat. nauk*, n. 2 (1972) 114-117 (in Russian).

- [68] Dirac P. A. M., *Long range forces and broken symmetries*, Proc. Roy. Soc. (London) **A333** (1973) 403-418.
- [69] Obukhov Yu.N., *Conformal invariance and space-time torsion*, Phys. Lett. **A90** (1982) 13-16.
- [70] Bergmann P., *On Einstein's  $\lambda$ -transformations*, Phys. Rev. **103** (1956) 780-781.
- [71] Chernikov N. A. and Tagirov E. A., *Quantum theory of scalar field in de Sitter space-time*, Ann. Inst. H. Poincaré **A9** (1968) 109-141.
- [72] Callan C. G., Coleman S., and Jackiw R., *A new improved energy-momentum tensor*, Ann. Phys. **59** (1970) 42-73.
- [73] Freund P. G. O., *Local scale invariance and gravitation*, Ann. Phys. **74** (1974) 440-454.
- [74] Ponomarev V.N. and Tseytlin A.A., *Microscopic gravity as a Yang-Mills field*, Phys. Lett. **A70** (1979) 164-166.
- [75] Rumyantseva E.N., *Static model of the spherical Friedman world*, in: "Problems of the Theory of Gravitation and Elementary Particles" Ed. K.P. Stanyukovich (Atomizdat: Moscow, 1978) issue 9, p. 29-33.
- [76] Parker L., *Particle creation in expanding universes*, Phys. Rev. Lett. **21** (1968) 562-564.
- [77] Zeldovich Ya.B. and Starobinsky A.A., *Particle production and vacuum polarization in an anisotropic gravitational field*, Sov. Phys. JETP **34** (1972) 1159-1166 [*ZhETF* **61** (1971) 2161-2175 (in Russian)].
- [78] Ponomarev V.N. and Pronin P.I., *Creation of massless scalar particles by the torsion field*, Theor. Math. Phys. **39** (1979) 559-561 [*Teor. Mat. Fiz.* **39** (1979) 425-428 (in Russian)].
- [79] Cartan E., *The Theory of Spinors* (MIT Press: Cambridge, MA, 1966).
- [80] Infeld L. and van der Waerden B.L., *Die Wellengleichungen des Elektrons in der allgemeinen Relativitätstheorie*, Sitzungsber. preuss. Akad. Wiss. Berlin (1933) 380-401.
- [81] Corson E. H., *Introduction to tensors, spinors and relativistic wave equations* (Blackie: London, 1953).
- [82] Schmutzer E., *Relativistische Physik* (Akademie: Berlin, 1968).
- [83] Weyl H., *A Remark on the coupling of gravitation and electron*, Phys. Rev. **77** (1950) 699-701.

- [84] Weyl H., *Elektron und Gravitation*, *Z. Phys.* **56** (1929) 330-352.
- [85] Fock V. and Ivanenko D., *Über eine mögliche geometrische Deutung der relativistischen Quantentheorie*, *Z. Phys.* **54** (1929) 798-802.
- [86] Hawking S.W. and Ellis G.F.R., *The Large Scale Structure of Space-Time* (Cambridge Univ. Press: Cambridge, 1973).
- [87] Gurovich V.Ts., *Quantum Effects and Regular Models of the Universe*, Preprint of Institute of Physics of Acad. Sci. KirgSSR (Frunze, 1980) 112 pp.
- [88] Ponomarev V.N., *Singularities and torsion*, in: "Relativity Theory and Gravitation", Ed. V.I. Rodichev (Nauka: Moscow, 1976) 167-176.
- [89] Benn I.M., Dereli T., and Tucker R.W., *Gauge field interactions in spaces with arbitrary torsion*, *Phys. Lett.* **B96** (1980) 100-104.
- [90] Dereli T. and Tucker R.W., *Weyl scaling and spinor matter interactions in scalar-tensor theories of gravitation*, *Phys. Lett.* **B110** (1982) 206-210.
- [91] Nieh H.T., *A spontaneously broken conformal gauge theory of gravitation*, *Phys. Lett.* **A88** (1982) 388-390.
- [92] Isham C.J., *Quantum gravity - an overview*, in: "Quantum Gravity II: A second Oxford symposium", eds C.J. Isham, R. Penrose and D.W. Sciama, (Clarendon Press: Oxford, 1981) 1-62.
- [93] Van Nieuwenhuizen P., *Supergravity*, *Phys. Repts.* **68** (1981) 189-398.
- [94] Trautman A., *A classification of space-time structures*, *Repts Math. Phys.* **10** (1976) 297-310.
- [95] Lubkin E., *Geometric definition of gauge invariance*, *Ann. Phys. (USA)* **23** (1963) 233-283.
- [96] Benn I.M., Dereli T., and Tucker R.W., *A critical analysis of some fundamental differences in gauge approach to gravitation*, *J. Phys.* **A15** (1982) 849-866.
- [97] Trautman A., *Fiber bundles, gauge fields and gravitation*, in: "General Relativity and Gravitation: One Hundred Years after the Birth of Albert Einstein", Ed. A. Held (Plenum: New York, 1980) vol. **1**, 287-308.
- [98] Hehl F.W., Nitsch J., and von der Heyde P., *Gravitation and the Poincaré gauge field theory with quadratic Lagrangians*, in: "General Relativity and Gravitation: One Hundred Years after the Birth of Albert Einstein", Ed. A. Held (Plenum: New York, 1980) vol. **1**, 329-355.

- [99] Ponomarev V.N. and Tseytlin A.A., *Gauge approach in gravity*, in: “Gravitation and General Relativity” /Ed. V.R.Kaigorodov (Kazan Univ. Publ. House: Kazan, 1983) **19**, 124-133 (in Russian).
- [100] Tseytlin A.A., *Poincaré and de Sitter gauge theories of gravity with propagating torsion*, *Phys. Rev.* **D26** (1982) 3327-3341.
- [101] Stelle K.S. and West P.C., *Spontaneously broken de Sitter symmetry and the gravitational holonomy group*, *Phys. Rev.* **D21** (1980) 1466-1481.
- [102] Pilch K.A., *Geometrical meaning of the Poincaré group gauge theory*, *Lett. Math. Phys.* **4** (1980) 49-51.
- [103] Norris L.K., Fulp R.O., and Davis W.R., *Underlying fiber bundle structure of  $A(4)$  gauge theories*, *Phys. Lett.* **A79** (1980) 278-282.
- [104] Luehr C.P. and Rosenbaum M., *Gravitation as an internal gauge theory of the Poincaré group*, *J. Math. Phys.* **21** (1980) 1432-1438.
- [105] Giachetti R., Ricci R., and Sorace E., *Tetrad fields and metric tensors in gauge theory of gravitation*, *Lett. Math. Phys.* **5** (1981) 85-91.
- [106] Misner C. W., *Harmonic maps as models for physical theories*, *Phys. Rev.* **D18** (1978) 4510-4524.
- [107] Ivanenko D. and Sardanashevily G., *The gauge treatment of gravity*, *Phys. Repts.* **94** (1983) 3-45.
- [108] Ne’eman Y. and Šijački Dj., *Unified affine gauge theory of gravity and strong interactions with finite and infinite  $GL(4, R)$  spinor fields*, *Ann. Phys. (USA)* **120** (1979) 292-315.
- [109] Isham C. J., Salam A., and Strathdee J., *Nonlinear realizations of space-time symmetries. Scalar and tensor gravity*, *Ann. Phys.* **62** (1971) 98-119.
- [110] Borisov A.B. and Ogievetsky V.I., *Theory of dynamical affine and conformal symmetries as the theory of the gravitational fields*, *Theor. Math. Phys.* **21** (1974) 1179-1188 [*Teor. Mat. Fiz.* **21** (1974) 329-342 (in Russian)].
- [111] Volkov M.K. and Pervushin V.N., *Essentially Non-Linear Quantum Theories, Dynamical Symmetries and Meson Physics* (Atomizdat: Moscow, 1978) 240 p. (in Russian).
- [112] Callan C.G., Jr., Coleman S., Wess J., and Zumino B., *Structure of phenomenological Lagrangians. II*, *Phys. Rev.* **177** (1969) 2247-2258.
- [113] Salam A. and Strathdee J., *Nonlinear Realizations. I. The Role of Goldstone Bosons*. *Phys. Rev.* **184** (1969) 1750-1759.

- [114] Volkov D.V., *Phenomenological Lagrangians, Physics of Elementary Particles and Atomic Nuclei* **4** (1973) 3-41 (in Russian).
- [115] Goldstone J., *Field theories with "Superconductor" solutions*, *Nuovo cimento* **19** (1961) 154-164.
- [116] Cho Y. M., *Nonlinear realization and unified interaction*, *Phys. Rev.* **D18** (1978) 2810-2812.
- [117] Henneaux M., *Zero Hamiltonian signature spacetime*, *Bull. Soc. Math. Belgique* **31** (1979) 47-63.
- [118] Teitelboim C., *Quantum mechanics of the gravitational field*, *Phys. Rev.* **D25** (1982) 3159-3179.
- [119] Pilati M., *Strong-coupling quantum gravity. I. Solution in a particular gauge*, *Phys. Rev.* **D26** (1982) 2645-2663.
- [120] Tolman R., *Relativity, Thermodynamics and Cosmology* (Clarendon: Oxford, 1969).
- [121] Stanyukovich K.P. and Melnikov V.N., *Hydrodynamics, Fields and Constants in the Theory of Gravitation* (Energoizdat: Moscow, 1982) 323 p. (in Russian).
- [122] Markov M.A., *Can the gravitational field prove essential for the theory of elementary particles?*, *Progr. Theor. Physics Suppl., Extra Number* (1965) 85-95.
- [123] Kopczyński W., *An anisotropic universe with torsion*, *Phys. Lett.* **A43** (1973) 63-64.
- [124] Lifshitz E.M., *On gravitational stability of expanding world*, *ZhETF* **16** (1946) 587-593.
- [125] Nurgaliev I. S. and Ponomarev V., *The earliest evolution stages of the Universe and the space-time torsion*, in: "Contr. papers to the 10th Intern. Conf. on Gen. Relat. Grav.", Padova, 4-9 July 1983, ed. B. Bertotti, F. De Felice, and A. Pascolini (Consiglio Nazionale Delle Ricerche: Roma, 1983) vol. 2, p. 771-773.
- [126] Kerlick G.D., *Cosmology and particle pair production via gravitational spin-spin interaction in the Einstein-Cartan-Sciama-Kibble theory of gravity*, *Phys. Rev.* **D12** (1975) 3004-3006.
- [127] Pronin P. I., *Renormalization of selfinteracting fields in a mean field expansion in curved spacetime*, in: "Contr. papers to the 10th Intern. Conf. on Gen. Relat. Grav.", Padova, 4-9 July 1983, ed. B. Bertotti, F. De Felice, and A. Pascolini (Consiglio Nazionale Delle Ricerche: Roma, 1983) vol. 2, p. 1151-1153.

- [128] Yang C.N., *Integral formalism for gauge fields*, *Phys. Rev. Lett.* **33** (1974) 445-447.
- [129] Fairchild E.E., Jr., *Gauge theory of gravitation*, *Phys. Rev.* **D14** (1976) 384-391; (E) 3439.
- [130] Lopez C., *Gravitational energy from a quadratic Lagrangian with torsion*, *Int. J. Theor. Phys.* **16** (1977) 167-176.
- [131] Frolov B.N., *Tetrad Palatini formalism and quadratic Lagrangians in the gravitational field theory*, *Acta Phys. Pol.* **B9** (1978) 823-829.
- [132] Nikolaenko V.M., *Second order Lagrangians of curvature in the theory of gravity*, *Acta Phys. Pol.* **B7** (1976) 681-692.
- [133] Minkevich A.V., *Generalized cosmological Friedmann equations without gravitational singularity*, *Phys. Lett.* **A80** (1980) 232-234.
- [134] Neville D.E., *Spin-2 propagating torsion*, *Phys. Rev.* **D23** (1981) 1244-1249.
- [135] Sezgin E., and van Nieuwenhuizen P., *New ghost-free gravity Lagrangians with propagating torsion*, *Phys. Rev.* **D21** (1980) 3269-3280.
- [136] Rauch R.T., and Shaw S.J., and Nieh H.T., *Birkhoff's theorem for ghost-free tachyon-free  $R + R^2 + Q^2$  theories with torsion*, *Gen. Relat. Grav.* **14** (1982) 331-354.
- [137] Debney G., Fairchild E.E., Jr., and Siklos S.T.C., *Equivalence of vacuum Yang-Mills gravitation and vacuum Einstein gravitation*, *Gen. Relat. Grav.* **9** (1978) 879-887.
- [138] Christensen S.M., *Second and fourth order invariants on curved manifold with torsion*, *J. Phys.* **A13** (1980) 3001-3009.
- [139] Christensen S.M. and Duff M.J., *Quantizing gravity with a cosmological constant*, *Nucl. Phys.* **B170[FS1]** (1980) 480-505.
- [140] Mansouri F., and Chang L.N., *Gravitation as a gauge theory*, *Phys. Rev.* **D13** (1976) 3192-3200.
- [141] Camenzind M., *Internal geometrical structures on Minkowski space-time and Yang-Mills symmetries*, *Intern. J. Theor. Phys.* **10** (1974) 197-216.
- [142] Hawking S. W., *Gravitational instantons*, *Phys. Lett.* **A60** (1977) 81-83.
- [143] Belavin A. A. and Burlankov D. M., *The renormalisable theory of gravitation and the Einstein equations*, *Phys. Lett.* **A58** (1976) 7-8.
- [144] Eguchi Y. and Freund P. G. O., *Quantum gravity and world topology*, *Phys. Rev. Lett.* **37** (1976) 1251-1254.

- [145] Hanson A. and Regge T., *Torsion and quantum gravity*, *Lect. Notes Phys.* **94** (1979) 354-361.
- [146] Barker B.M., Gupta S.N., and Haracz R.D., *One-graviton exchange interaction of elementary particles*, *Phys. Rev.* **149** (1966) 1027-1032.
- [147] Hehl F.W., Lord E.A., and Ne'eman Y., *Hypermomentum in hadron dynamics and in gravitation*, *Phys. Rev.* **D17** (1978) 428-433.
- [148] Collins P.D.B. and Squires E.J., *Regge poles in particle physics* (Springer: Berlin, 1968).
- [149] Jackiw R., *Introducing scale symmetry*, *Phys. Today* **25** (1972) 23-27.
- [150] Loos H.G., *The range of gauge fields*, *Nucl. Phys.* **72** (1965) 677-691.
- [151] Ikeda M. and Miyachy Y., *On the static and spherically symmetric solutions of the Yang-Mills Field*, *Prog. Theor. Phys.* **27** (1962) 474-482.
- [152] Ponomarev V.N. and Tseytlin A. A., *Spherically symmetric solutions of Yang-Mills equations in Riemann spaces*, *Yad. Fiz.* **29** (1979) 539-541 (in Russian).
- [153] Goldstone J., Salam A., and Weinberg S., *Broken symmetries*, *Phys. Rev.* **127** (1962) 965-970.
- [154] Fadeev L.D. and Popov V.N., *Covariant quantization of the gravitational field*, *Sov. Phys. Uspekhi* **16** (1974) 777-788 [*Usp. Fiz. Nauk* **111** (1973) 427-450, in Russian].
- [155] Batalin I. A. and Vilkovsky G. ., *Relativistic S-matrix of dynamical systems with boson and fermion constraints*, *Phys. Lett.* **B69** (1977) 309-312.
- [156] 't Hooft G., *Renormalization of massless Yang-Mills fields*, *Nucl. Phys.* **B33** (1971) 173-199.
- [157] De Witt B.S., *Quantum theory of gravity. II. The manifestly covariant theory*, *Phys. Rev.* **162** (1967) 1195-1239.
- [158] De Witt B.S., *Quantum theory of gravity. III. Applications of the covariant theory*, *Phys. Rev.* **162** (1967) 1239-1256.
- [159] Fradkin E. S. and Tjutin I. V., *S matrix for Yang-Mills and gravitational fields*, *Phys. Rev.* **D2** (1970) 2841-2857.
- [160] Hawking S., *Particle creation by black holes*, *Comm. Math. Phys.* **43** (1975) 199-220.

- [161] Frolov V. P. and Vilkovisky G. A., *Quantum gravity removes classical singularities and shortens life of black holes*, in: *Proc. of the 2nd Marcel Grossmann Meeting on General Relativity, Trieste, Italy, 1979*, Ed. R. Ruffini (North-Holland: Amsterdam, 1982) p. 455-469.
- [162] Feynman R.P., *Space-time approach to non-relativistic quantum mechanics*, *Rev. Mod. Phys.* **20** (1948) 367-387.
- [163] Lapchinsky V.G. and Rubakov V.A., In: “*Problems of the theory of gravitation and elementary particles*”, ed. K.P. Stanyukovich (Atomizdat: Moscow, 1978) issue 9, p. 33-40.
- [164] Barvinsky A.O. and Ponomarev V.N., *Global effects of the production of scalar matter in quantum cosmology*, *Theor. Math. Phys.* **37** (1978) 983-991 [*Teor. Mat. Fiz.* **37** (1978) 224-236 (in Russian)].
- [165] Misner C.W., *A minisuperspace example: The Gowdy  $T^3$  cosmology*, *Phys. Rev.* **D8** (1973) 3271-3285.
- [166] Peres A., *On Cauchy’s problem in general relativity. II*, *Nuovo cimento* **26** (1962) 53-62.
- [167] Wheeler J.A., *Einsteins Vision* (Springer: Berlin, 1968).
- [168] Barvinsky A.O., *Methods of the Quantum Theory of Gravitation and Some of Their Applications*, PhD thesis (Moscow State University: Moscow, 1981).
- [169] Leutwyler H., *Gravitational field: Equivalence of Feynman quantization and canonical quantization*, *Phys. Rev.* **134** (1964) B1155-B1182.
- [170] Dirac P.A.M., *On the analogy between classical and quantum mechanics*, *Revs Mod. Phys.* **17** (1945) 195-199.
- [171] Vilkovisky G.A., *On the quantization of a particle in an external gravitational field*, *Theor. Math. Phys.* **8** (1971) 889-895 [*Teor. Mat. Fiz.* **8** (1971) 359-368 (in Russian)].
- [172] Cheng K.S., *Quantization of a general dynamical system by Feynman’s Path integration formulation*, *J. Math. Phys.* **13** (1972) 1723-1726.
- [173] Kuchař K., *Geometrodynamics regained: A Lagrangian approach*, *J. Math. Phys.* **15** (1974) 708-715.
- [174] Kopczyński W., *A non-singular universe with torsion*, *Phys. Lett.* **A39** (1972) 219-220.
- [175] Tafel J., *A class of cosmological models with torsion and spin*, *Acta Phys. Pol.* **B6** (1975) 537-554.

- [176] Kuchowicz B., *A static universe filled with spinning matter and magnetic field*, *Current Sci. (India)* **44** (1975) 537-538.
- [177] Nurgaliev I.S. and Ponomariev V.N., *Torsion of space-time and cosmology*, *Sov. J. Phys.* **25** (1982) 929-931 [*Izvestia VUZov, Fizika*, n. 10 (1982) 63-65 (in Russian)].
- [178] Kaup D. J. and Vitello A. P., *Solvable quantum cosmological model and the importance of quantizing in a special canonical frame*, *Phys. Rev.* **D9** (1974) 1648-1655.
- [179] Tarski J., *Model of linearized deviations in the quantum theory of gravity*, *Ann. Inst. H. Poincaré* **A20** (1974) 95-111.
- [180] Christodoulou D., *The chronos principle*, *Nuovo cimento* **B26** (1975) 67-93.
- [181] Kuchař K., *Canonical quantization of cylindrical gravitational waves*, *Phys. Rev.* **D4** (1971) 955-986.
- [182] Liang E.P.T., *Quantum models for the lowest-order velocity-dominated solutions of irrotational dust cosmologies*, *Phys. Rev.* **D5** (1972) 2458-2466.
- [183] Brill D. and Goudi R., *Quantization of general relativity*, *Rep. Prog. Phys.* **33** (1970) 413-488.
- [184] De Witt B.S., *The Everett-Wheeler interpretation of quantum mechanics*, in: “*Battelle Rencontres: 1967 Lectures in Mathematics and Physics*”, eds C.M. DeWitt and J.A. Wheeler (W. A. Benjamin, Inc.: New York, 1968) 318-332.
- [185] Everett H., III, “*Relative state*” *formulation of quantum mechanics*, *Revs Mod. Phys.* **29** (1957) 454-462.
- [186] Wheeler J.A., *Assessment of Everett’s “Relative state” formulation of quantum theory*, *Revs Mod. Phys.* **29** (1957) 463-465.
- [187] Ponomariev V. and Zhytnikov V., *On the equivalence of the vacuum Einstein equation and the equations of the gauge theory of gravity*, in: “*Contr. papers to the 10th Intern. Conf. on Gen. Relat. Grav.*”, Padova, 4-9 July 1983, ed. B. Bertotti, F. De Felice, and A. Pascolini (Consiglio Nazionale Delle Ricerche: Roma, 1983) vol. 1, p. 330-331.
- [188] Hehl F.W., *On the energy tensor of spinning massive matter in classical field theory and general relativity*, *Repts Math. Phys.* **9** (1976) 55-82.
- [189] Ni W.-T., *Yang’s gravitational field equations*, *Phys. Rev. Lett.* **35** (1975) 319-320.

- [190] Batalin I.A. and Fradkin E.S., *Operator quantization of relativistic dynamical systems subject to first class constraints*, P.N. Lebedev Phys. Inst. preprint No. 51, (Moscow, 1983) 26 p.
- [191] Obukhov Yu.N. and Korotky V.A., *The Weyssenhoff fluid in Einstein-Cartan theory*, *Class. Quantum Grav.* **4** (1987) 1633-1657.
- [192] Weinberg S., *The quantum theory of fields. Volume 1: Foundations* (Cambridge University Press: Cambridge, 2005).
- [193] Obukhov Yu.N., Ponomarev V.N., and Zhytnikov V.V., *Quadratic Poincaré gauge theory of gravity: a comparison with the general relativity theory*, *Gen. Relat. Grav.* **21** (1989) 1107-1142.
- [194] Nakamichi A., Sugamoto A., and Oda I., *Topological four-dimensional self-dual gravity*, *Phys. Rev. D* **44** (1991) 3835-3842.
- [195] Gupta S.N., *Particle-particle and particle-antiparticle interactions*, *Nucl. Phys.* **57** (1964) 19-28.
- [196] Obukhov Yu.N. and Yakushin I.V., *Experimental bounds on parameters of spin-spin interaction in gauge theory of gravitation*, *Theor. Math. Phys.* **90** (1992) 209-213.
- [197] Obukhov Yu.N. and Yakushin I.V., *Experimental restrictions on spin-spin interactions in gauge gravity*, *Int. J. Theor. Phys.* **31** (1992) 1993-2001.
- [198] Yakushin I.V., *Contribution of torsion to the hyperfine splitting of the hydrogen atom*, *Sov. J. Nucl. Phys.* **55** (1992) 418-420.
- [199] Hehl F.W., McCrea J.D., Mielke E.W., and Ne'eman Y., *Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilaton invariance*, *Phys. Repts.* **258** (1995) 1-171.
- [200] Blagojević M., *Gravitation and gauge symmetries* (Institute of Physics Publishing: Bristol, 2002) 534 p.
- [201] Blagojević M., and Hehl F.W., *Gauge theories of gravitation: A reader with commentaries* (Imperial College Press: London, 2013) 635 pp.
- [202] Faddeev F.D., *Feynman integral for singular Lagrangians*, *Theor. Math. Phys.* **1** (1969) 1-13.
- [203] Popov V.N. and Faddeev L.D., *Perturbation theory for gauge invariant fields*, preprint FERMILAB-PUB-72-057-T, NAL-THY-57 (1972) 21 pp.
- [204] Barvinsky A.O., and Ponomarev V.N., *Quantum Geometrodynamics: The Path Integral and the Initial Value Problem for the Wave Function of the Universe*, *Phys. Lett.* **B167** (1986) 289-294.

- [205] Barvinsky A.O., *The General semiclassical solution of the Wheeler-DeWitt equations and the issue of unitarity in quantum cosmology*, *Phys. Lett.* **B241** (1990) 201-206.
- [206] Barvinsky A.O. and Krykhtin V., *Dirac and BFV Quantization Methods in the One-loop Approximation: Closure of the Quantum Constraint Algebra and the Conserved Inner Product*, *Class. Quantum Grav.* **10** (1993) 1957-1984.
- [207] Barvinsky A.O., *Operator Ordering in Theories Subject to Constraints of the Gravitational Type*, *Class. Quantum Gravity* **10** (1993) 1985-1999.
- [208] Barvinsky A.O., *Unitarity approach to quantum cosmology*, *Phys. Rep.* **230** (1993) 237-367.
- [209] Barvinsky A.O., *Geometry of the Dirac Quantization of Constrained Systems*, e-Print: gr-qc/9612003, 20 pp.
- [210] Barvinsky A.O., *BRST technique for the cosmological density matrix*, *JHEP* **1310** (2013) 051, 17 pp.
- [211] Becchi C., Rouet A., and Stora R., *Renormalization of the Abelian Higgs-Kibble Model*, *Commun. Math. Phys.* **42** (1975) 127-162.
- [212] Tyutin I.V., *Gauge Invariance in Field Theory and Statistical Physics in Operator Formalism*, Preprint LEBEDEV-75-39 (1975) 62 pp. (in Russian); e-Print: arXiv:0812.0580.
- [213] Fradkin E.S. and Vilkovisky G.A., *Quantization Of Relativistic Systems With Constraint*, *Phys. Lett.* **B55** (1975) 224-226.
- [214] Batalin I.A. and Vilkovisky G.A., *Relativistic S Matrix of Dynamical Systems with Boson and Fermion Constraints*, *Phys. Lett.* **B69** (1977) 309-312.
- [215] Fradkin E.S. and Fradkina T.E., *Quantization of Relativistic Systems with Boson and Fermion First and Second Class Constraints*, *Phys. Lett.* **B72** (1978) 343-348.
- [216] Batalin I.A. and Vilkovisky G.A., *Existence Theorem for Gauge Algebra*, *J. Math. Phys.* **26** (1985) 172-184.
- [217] Batalin I.A. and E.S.Fradkin, *Operator Quantization and Abelization of Dynamical Systems Subject to First Class Constraints*, *Riv. Nuovo Cim.* **9** (1986) 1-48.
- [218] Batalin I.A. and E.S.Fradkin, *Operatorial quantizaion of dynamical systems subject to constraints. A Further study of the construction*, *Ann. Inst. Henri Poincare* **49** (1988) 145-214.

- [219] Batalin I.A. and Fradkin E.S., *Operator Quantization of Dynamical Systems With Irreducible First and Second Class Constraints*, *Phys. Lett.* **B180** (1986) 157-162, Erratum: *Phys. Lett.* **B236** (1990) 528.
- [220] Henneaux M. and Teitelboim C., *Quantization of gauge systems*, Princeton University Press, Princeton, 1992.
- [221] Marnelius R., *Simple BRST quantization of general gauge models*, *Nucl. Phys.* **B395** (1993) 647-660.
- [222] Batalin I.A. and Marnelius R., *Solving general gauge theories on inner product spaces*, *Nucl. Phys.* **B442** (1995) 669-696.
- [223] Ferraro R., Henneaux M., and Puchin M., *Path integral and solutions of the constraint equations: The case of reducible gauge theories*, *Phys. Lett.* **B333** (1994) 380-385.
- [224] De Witt-Morette C., *The Semiclassical Expansion*, *Annals Phys.* **97** (1976) 367-399, Erratum: *Annals Phys.* **101** (1976) 682.
- [225] Faddeev L.D. and Popov V.N., *Feynman Diagrams for the Yang-Mills Field*, *Phys.Lett.* **B25** (1967) 29-30.
- [226] Barvinsky A. O., *Quantum Dirac constraints, Ward identities and path integral in relativistic gauge*, *Phys. Lett.* **B428** (1998) 322-328
- [227] Barvinsky A. O., *Solution of quantum Dirac constraints via path integral*, *Nucl. Phys.* **B520** (1998) 533-560.
- [228] Gribov V.N., *Quantization of non-Abelian gauge theories*, *Nucl. Phys.* **B139** (1978) 1-19.
- [229] Vandersickel, N. and Zwanziger D. (2012). *The Gribov problem and QCD dynamics*, *Phys. Rept.* **520** (2012) 175-251.
- [230] Altshuler B.L. and Barvinsky A.O., *Quantum cosmology and physics of transitions with a change of the spacetime signature*, *Physics-Uspekhi* **39** (1996) 429-459.
- [231] Giddings S.B. and Strominger A., *Baby Universes, Third Quantization and the Cosmological Constant*, *Nucl. Phys.* **B321** (1989) 481-508.
- [232] Hartle J.B. and Hawking S.W., *Wave Function of the Universe*, *Phys. Rev.* **D28** (1983) 2960-2975.
- [233] Hawking S.W., *The Quantum State of the Universe*, *Nucl. Phys.* **B239** (1984) 257-276.
- [234] Vilenkin A., *Quantum Creation of Universes*, *Phys. Rev.* **D30** (1984) 509-511.

- [235] Linde A.D., *Quantum Creation of the Inflationary Universe*, *Lett. Nuovo Cim.* **39** (1984) 401-405.
- [236] Rubakov V.A., *Particle Creation In A Tunneling Universe*, *JETP Lett.* **39** (1984) 107-110.
- [237] Zeldovich Ya.B. and Starobinsky A.A., *Quantum creation of a universe in a nontrivial topology*, *Sov. Astron. Lett.* **10** (1984) 135-137.
- [238] Barvinsky A.O. and Kamenshchik A.Yu., *Cosmological landscape from nothing: Some like it hot*, *JCAP* **0609** (2006) 014 (24 pages).
- [239] Barvinsky A.O., *Why there is something rather than nothing (out of everything)?*, *Phys. Rev. Lett.* **99** (2007) 071301 (4 pages).
- [240] Barvinsky A.O., Kamenshchik A. Yu., and Nesterov D.V., *New type of hill-top inflation*, *JCAP* **1601** (2016) 036 (15 pages).
- [241] Barvinsky A.O., *CFT driven cosmology and conformal higher spin fields*, *Phys. Rev.* **D93** (2016) 103530 (13 pages).
- [242] Barvinsky A.O., Kamenshchik A. Yu., and Starobinsky A.A., *Inflation scenario via the Standard Model Higgs boson and LHC*, *JCAP* **0811** (2008) 021 (13 pages).
- [243] Barvinsky A.O., Kamenshchik A. Yu., Kiefer C., Starobinsky A.A., and Steinwachs C.F., *Asymptotic freedom in inflationary cosmology with a non-minimally coupled Higgs field*, *JCAP* **0912** (2009) 003 (27 pages).



# Subject index

- Action
  - of gravitational field
    - in GR, 11, 36
  - of Yang-Mills type, 108, 122, 127
- Affine
  - connection structure, 1
  - frame, 92
  - tangent bundle, 92, 96
  - transformations, 92
- Anholonomy object, 7
- Arnowitt-Deser-Misner energy, 51
- Bianchi identity, 6
- BRST
  - equation, 149
  - inner product, 144
  - operator, 144
- Bundle
  - associated, 92, 96, 167
  - of affine frames, 92
  - of linear frames, 90, 92, 167
  - of orthonormal frames, 6, 101
  - principal, 6, 167
  - spinor, 169
  - tangent, 167
  - vector, 166
- Canonical (soldering) form, 90
- Canonical momentum, 20
  - gravitational, 37
  - in geometrodynamics with matter sources, 41
  - of matter field, 41
- Christoffel symbol, 4
- Commutation relations, 83, 129, 140
- Conformally invariant gravity theory in  $U_4$ , 77
- Connection, 2, 167
  - defect tensor, 4
  - form, 91
  - generalized affine, 94, 96
  - local Lorentz, 7
  - semi-symmetric, 12
  - spinor, 170
- Conservation of total angular momentum, 16, 113
- Constraint

- first kind, 22
  - primary, 20, 23
  - second kind, 22
- Contortion tensor, 4
- Coordinate frame, 6, 164
- Cosmological term, 78, 116
- Cotangent space, 91, 164
- Covariant derivative, 2, 168
  - spinor, 172
- Creation and annihilation
  - operators, 83, 136
- Cross-section, 90
- Curvature
  - form, 91
  - local Lorentz, 7
  - tensor, 3
    - decomposition, 5
    - homothetic, 3, 127
- De Sitter
  - boost, 104
  - group, 103
  - universe, 123
    - isotropic, 54, 75, 123, 134
- Density
  - matrix, 137
  - number of particles, 85, 86
- DeWitt supermetric, 37
- Dilation current, 122, 126, 128
- Dirac
  - brackets, 25
  - matrix, 171
  - quantization, 140
- Einstein's
  - $\lambda$ -transformations, 72
  - equation, 8
  - tensor, 6
  - universe, 78, 123
- Einstein-Cartan theory, 13
- Electromagnetic field, 20, 26
- Energy-momentum tensor
  - canonical, 14, 16
  - conservation law in ECT, 16
  - conservation law in gauge gravity, 113
  - conservation law in GR, 8, 16
  - effective, 16
  - metrical, 8, 14
  - of Dirac spinor, 63
  - of fluid, 61
- Exterior
  - differential form, 164
  - product, 165
- Extrinsic curvature, 32
- Faddeev-Popov
  - ghost fields, 132
  - operator, 131
- Frenkel condition, 58
- Friedman flat universe, 81
- Gauge
  - conditions, 25
  - gravity theory, 87, 106
  - group, 88
  - theory, 87
- Gauss-Bonnet identity, 115
- Gauss-Codazzi identity, 34
- Gauss-Weingarten formula, 32
- General
  - affine group, 92
  - linear group, 90, 121
- Ghost field, 122
- Goldstone field, 89, 97, 99, 102, 125
- Gravitational
  - constraints, 39, 44
  - field Hamiltonian, 38, 50
- Hamilton-Jacobi
  - equation, 28, 138
  - theory, 138
- Higgs
  - field, 99
  - mechanism, 99
- Hilbert action, 8
- Hodge operator, 110, 166
- Hypermomentum, 121
- Hypersurface, 30
- Ideal fluid, 58
- Infinitesimal transformation
  - of coordinates, 15, 18, 30, 88
- Instantons, 117, 118

- Involution relations, 23
- Ivanenko-Heisenberg equation, 64
- Killing vector, 30, 49
- Lapse and shift functions, 37, 53
- Lie derivative, 15, 30, 41
- Local Lorentz
  - (anholonomic) frame, 7
  - connection, 7, 101
- Lorentz group, 4, 8
- Manifold, 1, 163
- Matter with microstructure, 59
- Method of diagonalization
  - of Hamiltonian, 84, 136
- Metric structure, 1
- Metric-affine gravity, 121
- Metricity condition, 3
- Nieh-Yan topological invariant, 118
- Nonlinear gauge field, 98, 105
- Nonmetricity tensor, 3
- Normal basis on the hypersurface,
  - 31
- Palatini
  - equation, 16
  - principle, 11
- Parallel transport, 2, 7, 168
- Particle
  - interpretation, 83, 136
  - relativistic, 19, 26
- Path integral, 130
- Pauli matrix, 170
- Penrose-Hawking energy condition,
  - 75, 126
- Poincaré group, 100, 107
- Pontryagin index, 118
- Regge-Teitelboim theorem, 51
- Ricci identity, 5
- Riemann-Cartan
  - connection, 4
  - space, 7
- Riemannian space, 1
- Scalar field with conformal
  - coupling, 81
- Schrödinger equation, 130, 133
- sin-Gordon equation, 71
- Singular Lagrangian, 19
- Spin density tensor, 14
  - of Dirac spinor, 63
  - of fluid, 61
- Spontaneous
  - breaking of conformal
    - symmetry, 124
  - symmetry breaking, 97, 99
- Superhamiltonian, 41, 135
- Supermomenta, 43
- Superspace, 140
- Tangent
  - bundle, 6, 89, 167
  - space, 6, 90
  - vector, 164
- Tetrad coefficients (tetrads), 6
- Third quantization, 161
- Three-dimensional
  - derivative, 31
  - metric of hypersurface, 31
- Topological Hausdorff space, 163
- Torsion
  - form, 91
  - tensor, 3
  - decomposition, 4
- Translations, 89, 101
- Truncated Lagrangian, 10
- Unitarizing Hamiltonian, 145, 151
- Vector
  - bundle, 166
  - field, 164
  - non-linear, 68
- Weyl
  - group, 6, 171
  - space, 7
  - transformations, 71, 124
  - vector, 4, 128
- Weyssenhoff spinning fluid, 58
- Wheeler-DeWitt equation, 141
- Yang-Mills equation, 122



# Bibliography on gauge gravity theory and related issues

## A

1. Abe S., *Superfield formulation of a spinning particle in Riemann-Cartan space-time*, *Nuovo Cim.* **A89** (1985) 149-156.
2. Abedi H. and Salti M., *Multiple field modified gravity and localized energy in teleparallel framework*, *Gen. Rel. Gravit.* **47** (2015) 93.
3. Abedi H., Wright M., and Abbassi A.A., *Nonminimal coupling in anisotropic teleparallel inflation*, *Phys. Rev.* **D95** (2017) 064020 (11 pages).
4. Acedo L., *Autoparallel vs. geodesic trajectories in a model of torsion gravity*, *Universe* **1** (2015) 422-445.
5. Achucarro A., *Lineal gravity from planar gravity*, *Phys. Rev. Lett.* **70** (1993) 1037-1040.
6. Accioly A. and Blas H., *Metric-scalar gravity with torsion and the measurability of the non-minimal coupling*, *Mod. Phys. Lett.* **A22** (2007) 961.
7. Adak M., *Nonmetricity and torsion induced by dilaton gravity in two dimension*, *Gen. Rel. Grav.* **38** (2006) 971-980.
8. Adak M., *Non-minimally coupled Dirac equation with torsion: Poincaré gauge theory of gravity with even and odd parity terms*, *Class. Quantum Grav.* **29** (2012) 095006.
9. Adak M. and Dereli T., *The quadratic symmetric teleparallel gravity in two dimensions*, *Eur. Phys. Lett.* **82** (2008) 30008.
10. Adak M., Dereli T., and Ryder L.H., *Neutrino oscillations induced by spacetime torsion*, *Class. Quantum Grav.* **18** (2001) 1503-1512.
11. Adak M., Dereli T., and Ryder L.H., *Dirac equation in spacetimes with non-metricity and torsion*, *Int. J. Mod. Phys.* **D12** (2003) 145.
12. Adak M., Dereli T., and Ryder L.H., *Possible effects of space-time nonmetricity on neutrino oscillations*, *Phys. Rev.* **D69** (2004) 123002 (7 pages).

13. Adak M. and Grumiller D., *Poisson-sigma model for 2D gravity with non-metricity*, *Class. Quantum Grav.* **24** (2007) F65.
14. Adamowicz W., *Equivalence between Einstein-Cartan and general relativity theories in linear approximation for a classical model of spin*, *Bull. Acad. Polon. Sci., sér. sci. math., astr., phys.*, **23** (1975) 1203-1205.
15. Adamowicz W., *Plane waves in gauge theories of gravitation*, *Gen. Relat. Grav.* **12** (1980) 677-691.
16. Adamowicz W. and Trautman A., *The principle of equivalence for spin*, *Bull. Acad. Polon. Sci., sér. mat., astr., phys.*, **21** (1975) 339-342.
17. Adler R.J., *Spinors in Weyl geometry*, *J. Math. Phys.* **11** (1970) 1185-1191.
18. Agnese A.G. and Calvini P., *Gauge fields arising from space-time symmetries and gravitational theories, I and II*, *Phys. Rev.* **D12** (1975) 3800-3809.
19. Agricola I. and Friedrich T., *The Casimir operator of a metric connection with skew-symmetric torsion*, *J. Geom. Phys.* **50** (2004) 188-204.
20. Akdeniz K.G., Kizilersü A., and Rizaoglu E., *Instanton and eigen-modes in a two-dimensional theory of gravity with torsion*, *Phys. Lett.* **B215** (1988) 81-83.
21. Akdeniz K.G., Kizilersü A., and Rizaoglu E., *Fermions in two-dimensional theory of gravity with dynamical metric and torsion*, *Lett. Math. Phys.* **17** (1989) 315-320.
22. Akdeniz K.G., Dayi O.F., and Kizilersü A., *Canonical description of a two-dimensional gravity*, *Mod. Phys. Lett.* **A7** (1992) 1757-1764.
23. Akimov A.E. and Tarasenko V.Ya., *Models of polarized states of the physical vacuum and torsion fields*, *Russ. Phys. J.* **35** (1992) 214-222 [*Izvestiya VUZov, Fizika* n. **3** (1992) 13-23 (in Russian)].
24. Akimov A.E., Tarasenko V.Ya., and Shipov G.I., *Torsion fields as cosmophysical factor*, *Biofizika* **40** (1995) 938-943 (in Russian).
25. Albanese C. and De Ritis R., *Conformal equivalence in  $U_4$  - scalar field theories*, *Phys. Lett.* **A148** (1990) 301-305.
26. Aldersley S.J., *Scalar-metric and scalar-metric-torsion gravitational theories*, *Phys. Rev.* **D15** (1977) 3507-3512.
27. Aldersley S.J., *Metric-torsion gravitational theories*, *Gen. Relat. Grav.* **8** (1977) 397-409.
28. Aldersley S.J., *Charge conservation in metric-torsion gravitational theories*, *J. Math. Phys.* **19** (1978) 700-702.
29. Aldersley S.J., *The field equations of metric-affine gravitational theories*, *Z. Naturforsch.* **A33** (1978) 398-401.
30. Aldinger R.R., *Geometrical  $SO(4, 1)$  gauge theory as a basis of extended relativistic objects for hadrons*, *J. Phys. A: Math. Gen.* **A23** (1990) 1885-1907.
31. Aldrovandi R., *Breeding curvature from extended gauge covariance*, *Phys. Lett.* **A155** (1991) 459-463.
32. Aldrovandi R., Barros P.B., and Pereira J.G., *The equivalence principle revisited*, *Found. Phys.* **33** (2003) 545-575.
33. Aldrovandi R. and Pereira J.G., *A natural Poincaré gauge model*, *Phys. Rev.* **D33** (1986) 2788-2795.
34. Aldrovandi R. and Pereira J.G., *On the quantization of the Poincaré and de Sitter gauge models*, *J. Math. Phys.* **29** (1988) 1472-1476.
35. Aldrovandi R. and Pereira J.G., *An introduction to geometrical physics* (World Scientific: Singapore, 1995) 699 p.

36. Aldrovandi R. and Pereira J.G., *Gravitation: in search of the missing torsion*, *Annales de la Fondation L. de Broglie* **32** (2007) 229-251.
37. Aldrovandi R. and Pereira J.G., *Teleparallel gravity: An introduction* (Springer: Dordrecht, 2013) 214 pp.
38. Aldrovandi R., Pereira J.G., and Vu K.H., *Gravitation: global formulation and quantum effects*, *Class. Quantum Grav.* **21** (2004) 51.
39. Aldrovandi R. and Stedile E., *Complete gauge theory for the whole Poincaré group*, *Int. J. Theor. Phys.* **23** (1984) 301-323.
40. Alexander S. and Yunes N., *Chern-Simons modified gravity as a torsion theory and its interaction with fermions*, *Phys. Rev.* **D77** (2008) 124040 (11 pages).
41. Alekseev D.V. and Schapiro I.L., *Renormalization group approach in quantum field theory in curved space with torsion*, *Sov. Phys. J.* **33** (1990) 840-843 [*Izvestia VUZov, Fizika*, **n 10** (1990) 34-38 (in Russian)].
42. Ali S.A., Cafaro C., Capozziello S., and Corda C., *Abelian magnetic monopoles and topologically massive vector bosons in scalar-tensor gravity with torsion potential*, *Int. J. Mod. Phys.* **A23** (2008) 4315.
43. Ali S.A., Cafaro C., Capozziello S., and Corda C., *On the Poincaré gauge theory of gravitation*, *Int. J. Theor. Phys.* **48** (2009) 3426-3448.
44. Alimohammadi M. and Shariati A., *Neutrino oscillation in a space-time with torsion*, *Mod. Phys. Lett.* **A14** (1999) 267-274.
45. Alimohammadi M. and Shariati A., *Quantum reflection of massless neutrinos from a torsion-induced potential*, *Int. J. Mod. Phys.* **A15** (2000) 4099-4106.
46. Alix S.A., Cafaro C., Capozziello S., and Corda C., *A bound quantum particle in a Riemann-Cartan space with topological defects and planar potential*, *Phys. Lett.* **A366** (2007) 315-323.
47. Aman J.E., Fonseca-Neto J.B., MacCallum M.A.H., and Rebouças M.J., *Riemann-Cartan spacetimes of Gödel type*, *Class. Quantum Grav.* **15** (1998) 1089-1101.
48. Alfaro J. and Riquelme M.S., *Bosonic (p1)-forms in Einstein-Cartan theory of gravity*, *Phys. Rev.* **D90** (2014) 104005 (8 pages).
49. Allemandi G., Capone M., Capozziello S., and Francaviglia M., *Conformal aspects of the Palatini approach in extended theories of gravity*, *Gen. Rel. Grav.* **38** (2006) 33-60.
50. Amorim R., *Charged spinning fluids with magnetic dipole moment in the Einstein-Cartan theory*, *Phys. Rev.* **D31** (1985) 3099-3103.
51. Amorim R., *The motion of classical spinning particle in a Riemann-Cartan space-time*, *Gen. Relat. Grav.* **17** (1985) 525-533.
52. Amorim R., *Chromodynamics in Einstein-Cartan theory*, *Phys. Rev.* **D33** (1986) 2796-2800.
53. Anandan J., *Remarks concerning the geometries of gravity and gauge fields*, in: *Directions in General Relativity: Proc. of the Misner-Brill 1993 Int. Symp.*, Maryland (Cambridge: University Press, 1993) Eds. B.L. Hu, M.P. Ryan, Jr., and C.P. Vishveshwara, vol. **1**, 10-20.
54. Anandan J., *Topological and geometrical phases due to gravitational field with curvature and torsion*, *Phys. Lett.* **A195** (1994) 284-292.
55. Anandan J., *Gravitational phase operator and cosmic strings*, *Phys. Rev.* **D53** (1996) 779-786.
56. Anandan J. and Lesche B., *Interferometry in a space-time with torsion*, *Lett. Nuovo Cim.* **37** (1983) 391-397.
57. Anchordoqui L.A., *Wormholes in space-time with torsion*, *Mod. Phys. Lett.* **A13** (1998) 1095-1100.

58. Anchordoqui L.A., *Brans-Dicke wormholes endowed with torsion*, *Nuovo Cim.* **B113** (1998) 1497-1501.
59. Andersson F., and Edgar S.B., *Curvature-free asymmetric metric connections and Lanczos potentials in Kerr-Schild spacetimes*, *J. Math. Phys.* **39** (1998) 2859-2861.
60. Antoniadis I. and Odintsov S.D., *Conformal dynamics of quantum gravity with torsion*, *Mod. Phys. Lett.* **A8** (1993) 979-986.
61. Antonowicz M. and Szczyrba W., *Spinor matter fields in  $SL(2, C)$  gauge theories of gravity: Lagrangian and Hamiltonian approach*, *Phys. Rev.* **D31** (1985) 3104-3129.
62. Antonowicz M. and Szczyrba W., *An  $SU(2)$ -covariant dynamical formulation of the Einstein-Cartan-Dirac theory*, *Class. Quantum Grav.* **2** (1985) 515-533.
63. Antonowicz M. and Szczyrba W., *The dynamical structure of gravitational theories with  $GL(4, R)$  connections*, *J. Math. Phys.* **26** (1985) 1711-1727.
64. Antunes V.F. and Novello M., *Spin-2 fields from torsion: Dark energy and bouncing cosmology*, *Gravitation and Cosmology* **22** (2016) 1-9.
65. Ao X.-C. and Li X.-Z., *de Sitter gauge theory of gravity: an alternative torsion cosmology*, *JCAP* **10** (2011) 039.
66. Ao X.-C. and Li X.-Z., *Torsion cosmology of Poincaré gauge theory and the constraints of its parameters via SNeIa data*, *JCAP* **02** (2012) 003.
67. Ao X.-C., Li X.-Z., and Xi P., *Analytical approach of late-time evolution in a torsion cosmology*, *Phys. Lett.* **B694** (2010) 186-190.
68. Aouane R., Mena M., and Tahiri M., *On torsional observables in topological 4D gravity*, *Class. Quantum Grav.* **24** (2007) 2445.
69. Aprea G., Montani G., and Ruffini R., *Test particles behavior in the framework of a Lagrangian geometric theory with propagating torsion*, *Int. J. Mod. Phys.* **D12** (2003) 1875.
70. Aragone C. and Restuccia A., *Dynamical structure of linearized  $GL(4)$  gravities*, *Gen. Relat. Grav.* **9** (1978) 409-429.
71. Aragone C. and Restuccia A., *A gravitating  $SO(3, 1)$  gauge fields*, *Ann. Phys. (USA)* **113** (1978) 387-399.
72. Arbuzov B.A., and Filippov A.T., *Geometrization of electromagnetic field and violation of  $CP$ -invariance*, *ZhETF* **52** (1967) 1092-1105 (in Russian) [*Sov. Phys. JETP* **25** (1966) 725-734].
73. Arbuzov B.A., *On the unified description of gravitation and electromagnetism*, *ZhETF* **56** (1969) 1046-1056 (in Russian) [*Sov. Phys. JETP* **29** (1969) 562-567].
74. Arcos H.I., De Andrade V.C., and Pereira J.G., *Torsion and gravitation: A new view*, *Int. J. Mod. Phys.* **D13** (2004) 807 (12 pages).
75. Arcos H.I., Gribl Lucas T., and Pereira J.G., *Consistent gravitationally coupled spin-2 field theory*, *Class. Quantum Grav.* **27** (2010) 145007.
76. Arcos H.I., Mayor C.S.O., Otalora G., and Pereira J.G., *Spin-2 fields and helicity*, *Found. Phys.* **42** (2012) 1339-1349.
77. Arcos H.I. and Pereira J.G., *Torsion and the gravitational interaction*, *Class. Quantum Grav.* **21** (2004) 5193.
78. Arcos H.I. and Pereira J.G., *Torsion gravity: A reappraisal*, *Int. J. Mod. Phys.* **D13** (2004) 2193 (48 pages).
79. Aringazin A.K., and Mikhailov A.L., *Matter fields in spacetime with vector non-metricity*, *Class. Quantum Grav.* **8** (1991) 1685-1700.

80. Arkuszewski W., Kopczyński W., and Ponomariev V.N., *On the linearized Einstein-Cartan theory*, *Ann. Inst. H.Poincaré* **A21** (1974) 89-95.
81. Arkuszewski W., Kopczyński W., and Ponomariev V.N., *Matching conditions in the Einstein-Cartan theory of gravitation*, *Commun. Math. Phys.* **45** (1975) 183-191.
82. Aros R. and Contreras M., *Torsion induces gravity*, *Phys. Rev.* **D73** (2006) 087501 (4 pages).
83. Aslam A., Jamil M., and Myrzakulov R., *Noether gauge symmetry for the Bianchi type I model in  $f(T)$  gravity*, *Phys. Scr.* **88** (2013) 025003.
84. Assad M.J.D. and Letelier P.S., *On a class of inflationary universes of the self-consistent Einstein-Cartan theory*, *Phys. Lett.* **A145** (1990) 74-78.
85. Assad M.J.D. and Romero C., *Eternal and pulsating universe in Einstein-Cartan space-time*, *Astrophys. Space Sci.* **191** (1992) 289-297.
86. Asselmeyer T., *Generation of source terms in general relativity by differential structures*, *Class. Quantum Grav.* **14** (1997) 749-758.
87. Atazadeh K., *Stability of the Einstein static universe in Einstein-Cartan theory*, *JCAP* **06** (2014) 020.
88. Atkins W.R., Baker W.M., and Davis W.R., *On variational theories involving torsion*, *Phys. Lett.* **A61** (1977) 363-366.
89. Atkins W.K., Baker W.M., and Davis W.R., *On the properties of torsion in Riemann-Cartan space-time. II. Reinterpretation of gravitational-radiation solutions in  $U_4$* , *Nuovo Cim.* **B44** (1978) 17-22.
90. Attard J., François J., and Lazzarini S., *Weyl gravity and Cartan geometry*, *Phys. Rev.* **D93** (2016) 085032 (9 pages).
91. Audretsch J., *Dirac electron in space-times with torsion. Spinor propagation, spin precession, and nongeodesic orbit*, *Phys. Rev.* **D24** (1981) 1470-1477.
92. Audretsch J. and Lämmerzahl C., *Neutron interference: general theory of the influence of gravity, inertia and space-time torsion*, *J. Phys.* **A16** (1983) 2457-2477.
93. Audretsch J. and Lämmerzahl C., *Constructive approach to spacetime torsion*, *Class. Quantum Grav.* **5** (1988) 1285-1295.
94. Audretsch J., Hehl F.W., and Lämmerzahl C., *Matter wave interferometry and why quantum objects are fundamental for establishing a gravitational theory*, in: *Relativistic Gravity Research (Bad Honnef School, 1991)*, Eds. J.Ehlers and G.Schäfer, *Lect. Notes Phys.* **410** (1992) 369-407.
95. Aurilia A. and Martellini M., *A model for quantum gravity: an effective approach to Einstein theory*, *Phys.Lett.* **B123** (1983) 303-307.
96. Aurilia A. and Spallucci E., *New look at the Riemann-Cartan theory*, *Phys. Rev.* **D42** (1990) 464-468.
97. Avalos R., Dahia F., and Romero C., *On the embedding of Weyl manifolds*, *J. Math. Phys.* **58** (2017) 012502 (18 pages).
98. Ayón-Beato E., García A., Macías A., and Quevedo H., *Uniqueness theorems for static black holes in metric-affine gravity*, *Phys. Rev.* **D61** (2000) 084017 (6 pages).
99. Ayón-Beato E., García A., Macías A., and Quevedo H., *Static black holes of metric-affine gravity in the presence of matter*, *Phys. Rev.* **D64** (2001) 024026 (7 pages).

## B

100. Baaklini N.S., *Spin  $\frac{3}{2}$  field and Cartan's geometry*, *Lett. Math. Phys.* **2** (1977) 33-47.
101. Baaklini N.S., *Cartan's geometrical structure of supergravity*, *Lett. Math. Phys.* **2** (1977) 115-117.
102. Baaklini N.S., *Quantum spinors and the singularity theorem of general relativity*, *Phys. Lett.* **A66** (1978) 357-358.
103. Babourova O.V., *Variational theory of ideal fluid in quadratic gravitational theory in Riemann-Cartan space-time*, *Sov. J. Phys.* **32** (1989) 849-853 [*Izvestia VUZov, Fizika*, n. 10 (1989) 101-105 (in Russian)].
104. Babourova O.V., *Modified Friedmann-Lemaitre equation for dilaton-spin dark matter in Weyl-Cartan space*, *Grav. Cosmol.* **10** (2004) 121-126.
105. Babourova O.V. and Frolov B.N., *Pontryagin, Euler forms and Chern-Simons terms in Weyl-Cartan space*, *Mod. Phys. Lett.* **A12** (1997) 1267-1274.
106. Babourova O.V. and Frolov B.N., *The variational theory of the perfect dilaton-spin fluid in a Weyl-Cartan space*, *Mod. Phys. Lett.* **A12** (1997) 2943-2950.
107. Babourova O.V. and Frolov B.N., *Gauss-Bonnet type identity in Weyl-Cartan space*, *Int. J. Mod. Phys.* **A12** 3665-3668.
108. Babourova O.V. and Frolov B.N., *Perfect fluid and test particle with spin and dilatonic charge in a Weyl-Cartan space*, *Mod. Phys. Lett.* **A13** (1998) 7-13.
109. Babourova O.V. and Frolov B.N., *Perfect hypermomentum fluid: Variational theory and equations of motion*, *Int. J. Mod. Phys.* **A13** (1998) 5391-5407.
110. Babourova O.V. and Frolov B.N., *Matter with dilaton charge in Weyl-Cartan spacetime and evolution of the universe*, *Class. Quantum Grav.* **20** (2003) 1423.
111. Babourova O.V. and Frolov B.N., *Interaction of the 4-rotational gauge field with orbital momentum, gravidiamagnetic effect, and orbital experiment "Gravity Probe B"*, *Phys. Rev.* **D82** (2010) 027503 (3 pages).
112. Babourova O.V. and Frolov B.N., *The "Gravity Probe B" experiment, interaction between the orbital moment and gauge field, and a gravidiamagnetic effect*, *Russ. Phys. J.* **54** (2011) 124-126.
113. Babourova O.V., Frolov B.N., and Febres E.V., *Spherically symmetric solution of gravitation theory with a Dirac scalar field in the Cartan-Weyl space*, *Russ. Phys. J.* **57** (2015) 1297-1299.
114. Babourova O.V., Frolov B.N., and Febres E.V., *On the variational principle in the Cartan-Weyl space*, *Russ. Phys. J.* **58** (2015) 283-285.
115. Babourova O.V., Frolov B.N., and Klimova E.A., *Plane torsion waves in quadratic gravitational theories in Riemann-Cartan space*, *Class. Quantum Grav.* **16** (1999) 1149-1162.
116. Babourova O.V., Frolov B.N., Kudlaev P.E., and Romanova E.V., *Spherically symmetric solution in Cartan-Weyl space with Dirac scalar field*, in: "Gravitation, Astrophysics, and Cosmology", *Proc. Twelfth Asia-Pacific Internat. Conf., Moscow, 28 Jun - 5 July 2015*, Eds. V. Melnikov and J.-P. Hsu (World Scientific: Singapore, 2016) 191-195.
117. Babourova O.V., Frolov B.N., and Lipkin K.N., *Gravitation theory with a Dirac scalar field in the exterior form formalism and the cosmological constant problem*, *Grav. Cosmol.* **18** (2012) 225-231.
118. Babourova O.V., Frolov B.N., and Portnov Yu.A., *On inflation of dark matter with dilatation charge in Weyl-Cartan spacetime*, *Grav. Cosmol.* **11** (2005) 310-312.
119. Babourova O.V., Frolov B.N., and Scherban V.N., *Plane torsion waves in the Poincaré gauge gravitation theory*, *Russ. Phys. J.* **55** (2012) 726-728.

120. Babourova O.V., Frolov B.N., and Scherban V.N., *A study of plane torsion waves in the Poincaré gauge theory of gravity*, *Grav. Cosmol.* **19** (2013) 144-150.
121. Babourova O.V., Frolov B.N., and Zhukovsky V.Ch., *Gauge field theory for the Poincaré-Weyl group*, *Phys. Rev.* **D74** (2006) 064012 (12 pages).
122. Babourova O.V., Frolov B.N., and Zhukovsky V.Ch., *Theory of gravitation on the basis of the Poincaré-Weyl gauge group*, *Grav. Cosmol.* **15** (2009) 13-15.
123. Babourova O.V., and Korolev M.Yu., *Variational formalism and field equations of tetrad gravitation theory in the Weyl-Cartan space*, *Russ. Phys. J.* **49** (2006) 682-632.
124. Babourova O.V., Korolev M.Yu., and Frolov B.N., *Motion of matter in metric-affine theory of gravitation*, *Russ. Phys. J.* **37** (1994) 69-74 [*Izvestiya VUZov, Fizika*, n. 1 (1994) 76-82 (in Russian)]
125. Babourova O.V., Korolev M.Yu., and Umyarova I.Ya., *Variational formalism for the quadratic Lagrangians in tetrad gravitation theory*, *Russ. Phys. J.* **49** (2006) 531-536.
126. Babourova O.V. and Kostkin R.S., *Matter with dilatation charge in the Riemann-Cartan space*, *Russ. Phys. J.* **52** (2009) 487-493.
127. Babourova O.V., Kostkin R.S., and Frolov B.N., *Extension of the CARTAN package for symbolic calculations to space-time models with Weyl-Cartan structure*, *Grav. Cosmol.* **15** (2009) 302-305.
128. Babourova O.V., Kostkin R.S., and Frolov B.N., *The problem of a cosmological constant within the conformal gravitation theory in the Weyl-Cartan space*, *Russ. Phys. J.* **54** (2011) 121-123.
129. Babourova O.V., Kostkin R.S., and Frolov B.N., *The use of symbolic calculations in post-Riemannian and multidimensional theories of gravity*, *Grav. Cosmol.* **17** (2011) 335-338.
130. Babourova O.V., Lipkin K.N., and Frolov B.N., *Theory of gravity with the Dirac scalar field and the problem of cosmological constant*, *Russ. Phys. J.* **55** (2012) 855-857.
131. Babourova O.V., Lipkin K.N., Frolov B.N., and Scherban V.N., *Variational principle in post-Riemannian theories of gravitation*, *Russ. Phys. J.* **56** (2013) 722-724.
132. Babourova O.V., Vshivtsev A.S., Myasnikov A.P., and Frolov B.N., *The model of perfect fluid with spin and non-abelian color charge* *Doklady Akad. Nauk* **357** (1997) 183-186 (in Russian).
133. Babourova O.V., Vshivtsev A.S., Myasnikov A.P., and Frolov B.N., *Perfect spin fluid with intrinsic color charge*, *Phys. of Atom. Nucl.* **61** (1998) 802-807.
134. Babourova O.V., Vshivtsev A.S., Myasnikov A.P., and Frolov B.N., *Perfect spin fluid with a color charge in a color field in Riemann-Cartan space*, *Phys. of Atom. Nucl.* **61** (1998) 2175-2179.
135. Baekler P., *The unique spherically symmetric solution of the  $U_2$ -theory of gravity in the teleparallelism limit*, *Phys. Lett.* **B94** (1980) 44-50.
136. Baekler P., *A spherically symmetric vacuum solution of the quadratic Poincaré gauge field theory of gravitation with Newtonian and confinement potentials*, *Phys. Lett.* **B99** (1981) 329-332.
137. Baekler P., *A magnetic-monopole-type solutions in the Poincaré gauge field theory of gravity*, *Gen. Relat. Grav.* **18** (1986) 31-43.
138. Baekler P., *Prolongation structure and Backlund transformations of gravitational double duality equations*, *Class. Quantum Grav.* **8** (1991) 1023-1046.
139. Baekler P., Boulanger N., and Hehl F.W., *Linear connections with a propagating spin-3 field in gravity*, *Phys. Rev.* **D74** (2006) 125009 (20 pages).
140. Baekler P., Favaro A., Itin Y., and Hehl F.W., *The Kummer tensor density in electrodynamics and in gravity*, *Ann. Phys. (USA)* **349** (2014) 297-324.

141. Baekler P., and Gürses M., *Exact solutions of the Poincaré gauge theory from its linearized field equations*, *Lett. Math. Phys.* **14** (1987) 185-191.
142. Baekler P., Gürses M., and Hehl F.W., *A new method to solve the field equations of Poincaré gauge theories*, *Class. Quantum Grav.* **5** (1988) L105-L112.
143. Baekler P., Gürses M., Hehl F.W., and McCrea J.D., *The exterior gravitational field of a charged spinning source in the Poincaré gauge theory: a Kerr-Newman metric with dynamical torsion*, *Phys. Lett.* **A128** (1988) 245-250.
144. Baekler P., Hecht R., Hehl F.W., and Shirafuji T., *Mass and spin of exact solutions of the Poincaré gauge theory*, *Prog. Theor. Phys.* **78** (1987) 16-21.
145. Baekler P. and Hehl F.W., *A micro - de Sitter space-time with constant torsion: a new vacuum solution of the Poincaré gauge field theory*, in: "Internat. Symp. on Gauge Theory and Gravitation" (Nara, Japan,1982) *Lect. Notes Phys.* **176** (1983) 1-15.
146. Baekler P. and Hehl F.W., *A charged Taub-NUT metric with torsion. A new axially symmetric solution of the Poincaré gauge field theory*, *Phys. Lett.* **A100** (1984) 392-396.
147. Baekler P. and Hehl F.W., *Beyond Einstein-Cartan gravity: quadratic torsion and curvature invariants with even and odd parity including all boundary terms*, *Class. Quantum Grav.* **28** (2011) 215017.
148. Baekler P. and Hehl F.W., *On the dynamics of the torsion of space-time: exact solutions in a gauge theoretical model of gravity*, in: "From  $SU(3)$  to Gravity. Festschrift in honor of Y.Ne'eman", Eds. E. Gotsman, and G. Tauber (Cambridge Univ. Press: Cambridge, 1985) 341-359.
149. Baekler P., Hehl F.W., and Lenzen H.J., *Vacuum solutions with double duality properties of the Poincaré gauge field theory II*, in: "Proc. 3rd Marcel Grossmann Meeting on General Relativity", Ed. Hu Ning (North Holland: Amsterdam, 1983) 107-128.
150. Baekler P., Hehl F.W., and Mielke E.W., *Vacuum solutions with double duality properties of a quadratic Poincaré gauge field theory*, in: "Proc. of the 2nd Marcel Grossmann Meeting on Recent Progress of the Fundamentals of General Relativity", Ed. R. Ruffini (North Holland: Amsterdam, 1981) 413-450.
151. Baekler P., Hehl F.W., and Mielke E.W., *Nonmetricity and torsion: facts and fancies in gauge approaches to gravity*, in: "Proc. of the 4th Marcel Grossmann Meeting on General relativity", Ed. R. Ruffini (North Holland: Amsterdam, 1986) 277-316.
152. Baekler P., Hehl F.W., and Nester J.M., *Poincaré gauge theory of gravity: Friedman cosmology with even and odd parity modes: Analytic part*, *Phys. Rev.* **D83** (2011) 024001 (23 pages).
153. Baekler P. and Mielke E.W., *Effective Einsteinian gravity from Poincaré gauge field theory*, *Phys. Lett.* **A113** (1986) 471-475.
154. Baekler P. and Mielke E.W., *Hamiltonian structure of the Poincaré gauge theory and separation of non-dynamical variables in exact torsion solutions*, *Fortschr. Phys.* **36** (1988) 549-594.
155. Baekler P., Mielke E.W., and Hehl F.W., *Dynamical symmetries in topological 3D gravity with torsion*, *Nuovo Cim.* **B107** (1992) 91-110.
156. Baekler P., Mielke E.W., Hecht R., and Hehl F.W., *Kinky torsion in a Poincaré gauge model of gravity coupled to a massless scalar field*, *Nucl. Phys.* **B288** (1987) 800-812.
157. Baekler P., Seitz M., and Winkelmann V., *Cylindrically symmetric solutions of self-consistently coupled Dirac fields in gauge theories of gravity*, *Class. Quantum Grav.* **5** (1988) 479-490.
158. Baekler P. and Yasskin P.B., *All torsion-free spherical solutions of the quadratic Poincaré gauge theory of gravity*, *Gen. Relat. Grav.* **16** (1984) 1135-1155.

159. Bagrov V.G., Belov V.V., Trifonov A.Yu., and Yevseyevich A.Y., *The complex WKB-Maslov method for the Dirac equation in a torsion field. 1. Construction of trajectory-coherent states and the equation for spin*, *Class. Quantum Grav.* **8** (1991) 1349-1359.
160. Bagrov V.G., Belov V.V., Trifonov A.Yu., and Yevseyevich A.Y., *Quasi-classical trajectory-coherent approximation for the Dirac equation with an external electromagnetic field in Riemann-Cartan space. 2. Construction of TCS and equation of motion for spin*, *Class. Quantum Grav.* **8** (1991) 1833-1846.
161. Bagrov V.G., Belov V.V., Trifonov A.Yu., and Yevseyevich A.Y., *Quantization of closed orbits in Dirac theory by Maslov's complex germ method*, *J. Phys. A: Math. and Gen.* **A27** (1994) 1021-1043.
162. Bagrov V.G., Bukhbinder I.L., and Shapiro I.L., *Possible experimental manifestations of the torsion field*, *Russ. Phys. J.* **35** (1992) 208-213 [*Izvestiya VUZov, Fizika* n. **3** (1992) 5-12 (in Russian)].
163. Bagrov V.G., Trifonov A.Yu., and Yevseyevich A.Y., *Quantum mechanics of charged massive spin-1 particle in curved spacetime with torsion: quasiclassical analysis of the Proca equation based on the Maslov complex sprout method*, *Class. Quantum Grav.* **9** (1992) 533-544.
164. Bahamonde S., Böhmer C., and Wright M., *Modified teleparallel theories of gravity*, *Phys. Rev.* **D92** (2015) 104042 (7 pages).
165. Bahamonde S. and Wright M., *Teleparallel quintessence with a nonminimal coupling to a boundary term*, *Phys. Rev.* **D92** (2015) 084034 (12 pages).
166. Baikov P.A., Nelipa N.F., and Ostapchenko S.S., *Gauge-invariant affine and projective Lagrangians with physical propagators*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **29**, n. 4 (1988) 87-88 (in Russian).
167. Baikov P., Hayashi M., Nelipa N., and Ostapchenko S., *Ghost- and tachyon-free gauge-invariant, Poincaré, affine and projective Lagrangians*, *Gen. Relat. Grav.* **24** (1992) 867-880.
168. Baker W.M., *Torsion fields near singularities*, *Class. Quantum Grav.* **3** (1986) L19-L22.
169. Baker W.M., *Cosmic strings in Riemann-Cartan space-time*, *Class. Quantum Grav.* **7** (1990) 717-730.
170. Baker W.M., Atkins W.K., and Davis W.R., *On the properties of torsion in Riemann-Cartan space-times. I. Torsion associated with  $U_4$  reinterpretation of null electromagnetic solutions of Einstein's equations*, *Nuovo Cim.* **B44** (1978) 1-16.
171. Bakke K., *Torsion and noninertial effects on a nonrelativistic Dirac particle*, *Ann. Phys. (USA)* **346** (2014) 51-58.
172. Balek V., Melek M., and Petras M., *Description of spinning particle in general relativity using commuting spin variables*, *Czech. J. Phys.* **B37** (1987) 1321-1339.
173. Bamba K., Capozziello S., De Laurentis M., Nojiri S., and Sáez-Gómez D., *No further gravitational wave modes in  $F(T)$  gravity*, *Phys. Lett.* **B727** (2013) 194-198.
174. Bamba K. and Geng C.-G., *Thermodynamics of cosmological horizons in  $f(T)$  gravity*, *JCAP* **11** (2011) 008.
175. Bamba K., Geng C.-G., Lee C.-C., and Luo L.-W., *Equation of state for dark energy in  $f(T)$  gravity*, *JCAP* **01** (2011) 021.
176. Bamba K., Geng C.-G., and Luo L.-W., *Generation of large-scale magnetic fields from inflation in teleparallelism*, *JCAP* **10** (2012) 058.
177. Bamba K., Nashed G.G.L., El Hanafy W., and Ibraheem S.K., *Bounce inflation in  $f(T)$  cosmology: A unified inflaton-quintessence field*, *Phys. Rev.* **D94** (2016) 083513 (23 pages).

178. Bandyopadhyay P., *Quantum space-time, quantum gravity and torsion*, in “Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko”, Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 239-252.
179. Banerji K., *Some aspects of Holst and Nieh-Yan terms in general relativity with torsion*, *Class. Quantum Grav.* **27** (2010) 135012.
180. Banerjee R. and Roy D., *Trivial symmetries in a 3D topological torsion model of gravity*, *J. Phys.: Conf. Ser.* **405** (2012) 012028.
181. Banerji S., *Bounce of spheres in Einstein-Cartan theory*, *Gen. Rel. Grav.* **9** (1978) 783-792.
182. Barbashov B.M. and Leonovich A.A., *Differential identities and topological invariants in the Riemann geometry and in the Riemann-Cartan geometry, and in space-time of an arbitrary affine connection* (JINR: Dubna, 1983) **R5-83-398**, 4 pp.
183. Barbashov B.M. and Pestov A.B., *Weyl connection, non-Abelian gauge field, and torsion*, *Teor. Mat. Fiz.* **104**, N 3 (1995) 429-434 (in Russian); English transl.: *Theor. Math. Phys.* **104** (1995) 1104-1107.
184. Barbashov B.M., and Pestov A.B., *Weyl non-Abelian gauge field*, *Mod. Phys. Lett.* **A10** (1995) 193-197.
185. Barbashov B.M. and Pestov A.B., *On spinor representations in the Weyl gauge theory*, *Mod. Phys. Lett.* **A12** (1997) 1957-1968.
186. Barbashov B.M. and Pestov A.B., *Antisymmetric tensor fields and the Weyl gauge theory*, *Theor. Math. Phys.* **113** (1997) 1299-1308.
187. Barbashov B.M. and Pestov A.B., *The Weyl non-Abelian gauge field and the Thomas precession*, *Mod. Phys. Lett.* **A13** (1998) 2117-2121.
188. Barbosa A.L., Guillen L.T.C., and Pereira J.G., *Teleparallel equivalent of non-Abelian Kaluza-Klein theory*, *Phys. Rev.* **D66** (2002) 064028 (7 pages).
189. Barducci A., Casalbuoni R., and Lusanna L., *Classical spinning particles interacting with external gravitational field*, *Nucl. Phys.* **B24** (1977) 521-538.
190. Barrett J.N., *Holonomy and path structures in general relativity and Yang-Mills theory*, *Int. J. Theor. Phys.* **30** (1991) 1171-1215.
191. Barros M., Ferrández A., Javaloyes M.A., and Lucas P., *Geometry of relativistic particles with torsion*, *Int. J. Mod. Phys.* **A19** (2004) 1737.
192. Barth N.H., *Heat kernel expansion coefficient. I. An extension*, *J. Phys.* **A20** (1987) 857-874.
193. Barth N.H., *Heat kernel expansion coefficient. II. High-order operator*, *J. Phys.* **A20** (1987) 875-888.
194. Barth N.H. and Christensen S.M., *Arbitrary spin field equations on curved manifolds with torsion*, *J. Phys.* **A16** (1983) 543-563.
195. Barvinski A.O. and Gryzov Yu.V., *Generalized condition of causality in quantum field theory with torsion*, *Sov. J. Phys.* **32** (1989) 383-387 [*Izvestia VUZov, Fizika*, n 5 (1989) 74-78 (in Russian)].
196. Basilakos S., Capozziello S., De Laurentis M., Paliathanasis A., and Tsamparlis M., *Noether symmetries and analytical solutions in  $f(T)$  cosmology: A complete study*, *Phys. Rev.* **D88** (2013) 103526 (12 pages).
197. Basombrio F.G., *On spinors and gravity*, *Nuovo Cim.* **B57** (1980) 269-282.
198. Basombrio F.G., *A comparative review of certain gauge theories of the gravitational field*, *Gen. Relat. Grav.* **12** (1980) 109-136.

199. Bassetto A., Griguolo L., Giacconi P., and Soldati R., *Extended integrability conditions for vector - axial vector anomalies on Riemann-Cartan manifold*, in: "General relativity and gravitational physics". *Proc. of the 9th Ital. Conf. (Capri, Italy, September 25-28, 1990)* Eds. R. Cianci, R. de Ritis, G. Marmo, C. Rubano, P. Scudellaro, and M. Francaviglia (World Scientific: Singapore, 1991) 658-664.
200. Batakis N., *A unification scheme for the Einstein and Maxwell theories*, *Phys. Lett.* **A90** (1982) 115-118.
201. Batakis N.A. and Tsoubelis D., *Dual description of a spacetime in the Einstein and Einstein-Cartan theories*, *Phys. Lett.* **A85** (1981) 261-262.
202. Batakis N. and Tsoubelis D., *Study of Bianchi type V cosmological model with torsion*, *Phys. Rev.* **D26** (1982) 2611-2614.
203. Battiti R. and Toller M., *Zero-mass normal modes in linearized Poincaré gauge theories*, *Lett. Nuovo Cim.* **44** (1985) 35-39.
204. Bauer M., Freedman D.Z., and Haagensen P.E., *Spatial geometry of the electric field representation of non-abelian gauge theories*, *Nucl. Phys.* **B428** (1994) 147-168.
205. Bauerle G.G.A., *Elements of gauge theory and theory of gravity*, In: "Mathematical structures in field theories". *Proc. Seminar, Amsterdam, Netherlands, 1981-1982*, Eds. E.M. De Jager, and H.G.J. Pijls. (Amsterdam: CWI, 1984) 59-117.
206. Bauerle G.G.A. and Haneveld C.J., *Spin and torsion in the early Universe*, *Physica* **A121** (1983) 541-551.
207. Bauerle G.G.A. and Twelker H.F., *On the non-relativistic limit of a spin-1/2 particle in a classical gravitational field*, *Physica* **A130** (1985) 553-564.
208. Baurov A.Yu. and Pronin P.I., *Antisymmetric tensor field in the space with torsion*, *Moscow University Physics Bulletin*, **52**, n. 6 (1997) 19-23 (in Russian).
209. Baykal A. and Dereli T., *Nonminimally coupled Einstein-Maxwell model in a non-Riemann spacetime with torsion*, *Phys. Rev.* **D92** (2015) 065018 (10 pages).
210. Bécarr R., González P.A., and Vásquez Y., *Dirac quasinormal modes of Chern-Simons and BTZ black holes with torsion*, *Phys. Rev.* **D89** (2014) 023001 (9 pages).
211. Becattini F. and Tinti L., *Thermodynamical inequivalence of quantum stress-energy and spin tensors*, *Phys. Rev.* **D84** (2011) 025013 (21 pages).
212. Becker R.E., *The value of Einstein's mistakes*, *Physics Today* **59**, n. 4 (2006) 14-15.
213. Bedran M.L. and Garcia de Andrade L.C., *Cylindrically symmetric cluster in Einstein-Cartan theory*, *Progr. Theor. Phys.* **70** (1983) 1583-1587.
214. Bedran M.L. and Som M.M., *Conformally flat solution of a static dust sphere in Einstein-Cartan theory*, *J. Phys.* **A15** (1982) 3539-3542.
215. Bedran M.L. and Vaidya V.E.P., *The role of spin in cosmological models*, *Lett. Nuovo Cim.* **41** (1984) 73-74.
216. Bedran M.L., Vaidya V.E.P., and Som M.M., *Stationary cosmological solution with torsion*, *Nuovo Cim.* **B87** (1985) 101-108.
217. Bekaert X. and Boulanger N., *Tensor gauge fields in arbitrary representations of  $GL(D, R)$ . Duality and Poincaré lemma*, *Commun. Math. Phys.* **245** (2004) 27-67.
218. Bekaert X. and Boulanger N., *Tensor gauge fields in arbitrary representations of  $GL(D, R)$ : II. Quadratic actions*, *Commun. Math. Phys.* **271** (2007) 723-773.
219. Belyaev A.S. and Shapiro I.L., *The action for the (propagating) torsion and the limits on the torsion parameters from present experimental data*, *Phys. Lett.* **B425** (1998) 246-254.
220. Belyaev A.S. and Shapiro I.L., *Torsion action and its possible observables*, *Nucl. Phys.* **B543** (1999) 20-46.

221. Belyaev A.S., Shapiro I.L., and do Vale M.A.B., *Torsion phenomenology at the CERN LHC*, *Phys. Rev.* **D75** (2007) 034014 (8 pages).
222. Belyaev A.S., Thomas M.C., and Shapiro I.L., *Torsion as a dark matter candidate from the Higgs portal*, *Phys. Rev.* **D95** (2017) 095033 (10 pages).
223. Bengochea G.R., *Observational information for  $f(T)$  theories and dark torsion*, *Phys. Lett.* **B695** (2011) 405-411.
224. Bengochea G.R. and Ferraro R., *Dark torsion as the cosmic speed-up*, *Phys. Rev.* **D79** (2009) 124019 (5 pages).
225. Benn I.M., Dereli T., and Tucker R.W., *Gauge field interactions in spaces with arbitrary torsion*, *Phys. Lett.* **B96** (1980) 100-104.
226. Benn I.M., Dereli T., and Tucker R.W., *Gravitational monopoles with classical torsion*, *J. Phys.* **A13** (1980) L359-L363.
227. Benn I.M., Dereli T., and Tucker R.W., *Double-dual solutions of generalized theories of gravitation*, *Gen. Relat. Grav.* **13** (1981) 581-589.
228. Benn I.M., Dereli T., and Tucker R.W., *A critical analysis of some fundamental differences in gauge approach to gravitation*, *J. Phys.* **A15** (1982) 849-866.
229. Benn I.M. and Tucker R.W., *An introduction to spinors and geometry with applications in physics* (Adam Hilger: Bristol and Philadelphia, 1987) 358 p.
230. Berdichevsky V.L., *Continuum theory of dislocations revisited*, *Continuum Mechanics and Thermodynamics* **18** (2006) 195-222.
231. Bergmann P.G., de Sabbata V., Gillies G.T., and Pronin P.I. (Eds), "Spin in gravity. Is it possible to give an experimental basis to torsion?", Int. School of Cosmology and Gravitation XV Course, Erice, Italy, 13-20 May, 1997 (World Scientific: Singapore, 1998) 272 pp.
232. Bergmann P.G., and Komar A.B., *Generalized Weyl-type gauge geometry*, *J. Math. Phys.* **26** (1985) 2030-2031.
233. Bergshoeff E.A., Hartong J., and Rosseel J., *Torsional Newton-Cartan geometry and the Schrödinger algebra*, *Class. Quantum Grav.* **32** (2015) 135017 (35 pages).
234. Bergshoeff E.A. and Rosseel J., *A new look at Newton-Cartan gravity*, *Int. J. Mod. Phys.* **A31** (2016) 1630040 (19 pages).
235. Berman M.S., *A static universe with magnetic field in Einstein-Cartan theory*, *Nuovo Cim.* **B105** (1990) 1373-1375.
236. Berman M.S., *Inhomogeneous inflation with spinning fluid*, *Nuovo Cim.* **B105** (1990) 235-237.
237. Berman M.S., *Inflation in the Einstein-Cartan cosmological model*, *Gen. Relat. Grav.* **23** (1991) 1083-1088.
238. Berman M.S., *Shear and vorticity in a combined Einstein-Cartan-Brans-Dicke inflationary lambda-*Universe**, *Astrophys. Space Sci.* **317** (2008) 79-82.
239. Berman M.S., *Shear and vorticity in an accelerating Brans-Dicke lambda-*Universe* with torsion*, *Astrophys. Space Sci.* **317** (2008) 279-281.
240. Berman M.S., *Pryce-Hoyle tensor in a combined Einstein-Cartan-Brans-Dicke model*, *Int. J. Theor. Phys.* **48** (2009) 836-846.
241. Berredo-Peixoto G. de, *On the 1-loop calculations of softly broken fermion-torsion theory in curved space using the Stückelberg procedure*, *Int. J. Mod. Phys.* **A24** (2009) 1570.
242. Berredo-Peixoto G. de, Helayël-Neto J.A., and Shapiro I.L., *On the consistency of a fermion-torsion effective theory*, *JHEP* **02** (2000) 003.

243. Berredo-Peixoto G. de, Freidel L., Shapiro I.L., and de Souza C.A., *Dirac fields, torsion and Barbero-Immirzi parameter in cosmology*, *JCAP* **06** (2012) 017.
244. Berredo-Peixoto G. de, and Freitas E.A. de, *On the torsion effects of a relativistic spin fluid in early cosmology*, *Class. Quantum Grav.* **26** (2009) 175015.
245. Berredo-Peixoto G. de, and Maicá A.E., *Form factors and non-local multiplicative anomaly for fermions with background torsion*, *Class. Quantum Grav.* **31** (2014) 125002.
246. Berthias J.-P. and Shahid-Saless B., *Torsion and non-metricity in scalar-tensor theories of gravity*, *Class. Quantum Grav.* **10** (1993) 1039-1044.
247. Bhattacharya B., and Chatterjee K.R., *On the possible existence of a lepton dominated era in the very early universe*, *Phys. Lett.* **A133** (1988) 105-108.
248. Bilby B.A., *Continuous distribution of dislocations*, *Progr. Solid Mech.* **1** (1960) 331.
249. Bilby B.A., Bullogh R., Gardner L.R.T., and Smith E., *Continuous distribution of dislocations IV. Single glide and plain strain*, *Proc. Roy. Soc. Lond.* **A244** (1958) 538-557.
250. Bilby B.A., Bullogh R., and Smith E., *Continuous distributions of dislocations: a new application of the methods of non-Riemannian geometry*, *Proc. Roy. Soc. Lond.* **A231** (1955) 263-273.
251. Bilby B.A. and Gardner L.R.T., *Continuous distribution of dislocations V. Twisting under conditions of single glide*, *Proc. Roy. Soc. Lond.* **A247** (1958) 92-108.
252. Bilby B.A. and Smith E., *Continuous distribution of dislocations III*, *Proc. Roy. Soc. Lond.* **A236** (1956) 481-505.
253. Bini D. and Mashhoon B., *Weitzenböck's torsion, Fermi coordinates, and adapted frames*, *Phys. Rev.* **D91** (2015) 084026 (9 pages).
254. Biswas S.K. and Chakraborty S., *Interacting dark energy in  $f(T)$  cosmology: A dynamical system analysis*, *Int. J. Mod. Phys.* **D24** (2015) 1550046 (20 pages).
255. Blagojević M., *Gravitacija i lokalne simetrije* (Institute of Physics: Belgrade, 1997) 484 p. (in Serbo-Croatian).
256. Blagojević M., *Conservation laws in Poincaré gauge theory*, *Acta Phys. Pol.* **29** (1998) 881-890.
257. Blagojević M., *Hamiltonian structure and gauge symmetries of Poincaré gauge theory*, *Ann. d. Phys. (Leipzig)* **10** (2001) 367-391.
258. Blagojević M., *Gravitation and gauge symmetries* (Institute of Physics Publishing: Bristol, 2002) 534 p.
259. Blagojević M. and Cvetković B., *Black hole entropy from the boundary conformal structure in 3D gravity with torsion*, *JHEP* **10** (2006) 005.
260. Blagojević M. and Cvetković B., *Canonical structure of 3D gravity with torsion*, in: "Trends in General Relativity and Quantum Cosmology", vol. 2, C. Benton ed. (Nova Science Publishers: New York, 2006) pp. 103-123.
261. Blagojević M. and Cvetković B., *Asymptotic charges in 3d gravity with torsion*, *J. Phys.: Conf. Ser.* **33** (2006) 248.
262. Blagojević M. and Cvetković B., *Black hole entropy in 3D gravity with torsion*, *Class. Quantum Grav.* **23** (2006) 4781.
263. Blagojević M. and Cvetković B., *Covariant description of the black hole entropy in 3D gravity*, *Class. Quantum Grav.* **24** (2007) 129.
264. Blagojević M. and Cvetković B., *Electric field in 3D gravity with torsion*, *Phys. Rev.* **D78** (2008) 044036 (8 pages).

265. Blagojević M. and Cvetković B., *Self-dual Maxwell field in 3D gravity with torsion*, *Phys. Rev.* **D78** (2008) 044037 (8 pages).
266. Blagojević M. and Cvetković B., *Canonical structure of topologically massive gravity with a cosmological constant*, *JHEP* **05** (2009) 073.
267. Blagojević M. and Cvetković B., *Self-dual Maxwell field in 3D gravity with torsion and dynamical role of central charges*, *J. Phys.: Conf. Ser.* **189** (2009) 012010.
268. Blagojević M. and Cvetković B., *Conserved charges in 3D gravity*, *Phys. Rev.* **D81** (2010) 124024 (7 pages).
269. Blagojević M. and Cvetković B., *3D gravity with propagating torsion: The AdS sector*, *Phys. Rev.* **D85** (2012) 104003 (10 pages).
270. Blagojević M. and Cvetković B., *Three-dimensional gravity with propagating torsion: Hamiltonian structure of the scalar sector*, *Phys. Rev.* **D88** (2013) 104032 (15 pages).
271. Blagojević M. and Cvetković B., *Gravitational waves with torsion in 3D*, *Phys. Rev.* **D90** (2014) 044006 (9 pages).
272. Blagojević M. and Cvetković B., *Siklos waves with torsion in 3D*, *JHEP* **11** (2014) 141 (17 pages).
273. Blagojević M. and Cvetković B., *Vaidya-like exact solutions with torsion*, *JHEP* **05** (2015) 101 (16 pages).
274. Blagojević M. and Cvetković B., *Siklos waves in Poincaré gauge theory*, *Phys. Rev.* **D92** (2015) 024047 (9 pages).
275. Blagojević M. and Cvetković B., *Conformally flat black holes in Poincaré gauge theory*, *Phys. Rev.* **D93** (2016) 044018 (9 pages).
276. Blagojević M. and Cvetković B., *Generalized pp waves in Poincaré gauge theory*, *Phys. Rev.* **D95** (2017) 104018 (10 pages).
277. Blagojević M., Cvetković B., and Mišcović O., *Nonlinear electrodynamics in 3D gravity with torsion*, *Phys. Rev.* **D80** (2009) 024043 (9 pages).
278. Blagojević M., Cvetković B., Mišcović O., and Olea R., *Holography in 3D AdS gravity with torsion*, *JHEP* **5** (2013) 103.
279. Blagojević M., Cvetković B., and Vasilčić M., *“Exotic” black holes with torsion*, *Phys. Rev.* **D88** (2013) 101501(R) (3 pages).
280. Blagojević M. and Hehl F.W., *Gauge theories of gravitation: A reader with commentaries* (Imperial College Press: London, 2013) 635 p.
281. Blagojević M. and Nicolčić I.A., *Various aspects of the time gauge in Poincaré gauge theory*, *Lett. Nuovo Cim.* **38** (1983) 77-82.
282. Blagojević M. and Nicolčić I.A., *Hamiltonian structure of the theory of gravity with  $R + T^2$  type of Lagrangian*, *Nuovo Cim.* **B73** (1983) 238-273.
283. Blagojević M. and Nicolčić I.A., *Hamiltonian dynamics of Poincaré gauge theory: general structure in the time gauge*, *Phys. Rev.* **D28** (1983) 2455-2463.
284. Blagojević M. and Nikolić I.A., *Local Poincaré generators in the  $R + T^2 + R^2$  theory of gravity*, *Nuovo Cim.* **B101** (1988) 439-451.
285. Blagojević M. and Nikolić I.A., *Hamiltonian structure of the teleparallel formulation of general relativity*, *Phys. Rev.* **D62** (2000) 024021 (10 pages).
286. Blagojević M., Nikolić I.A., Popović D.S. and Zivanović D., *Poincaré gauge theory of gravitation and its Hamiltonian formulation*, *Nuovo Cim.* **B62** (1981) 257-272.
287. Blagojević M., Popović D.S., and Zivanović Dj., *Gravitational singularity in Poincaré gauge theory*, *Phys. Lett.* **B109** (1982) 431-433.

288. Blagojević M. and Vasilic M., *Hamiltonian analysis of extra gauge symmetries in an  $R+T^2$  theory of gravity*, *Phys. Rev.* **D34** (1986) 357-366.
289. Blagojević M. and Vasilic M., *Extra-gauge symmetries in a weak-field approximation of a  $R + R^2 + T^2$  theory of gravity*, *Phys. Rev.* **D35** (1987) 3748-3759.
290. Blagojević M. and Vasilic M., *Asymptotic symmetry and conserved quantities in the Poincaré gauge theory of gravity*, *Class. Quantum Grav.* **5** (1988) 1241-1257.
291. Blagojević M. and Vasilic M., *Gauge symmetries of the teleparallel theory of gravity*, *Class. Quantum Grav.* **17** (2000) 3785-3797.
292. Blagojević M. and Vasilic M., *Conservation laws in the teleparallel theory of gravity*, *Phys. Rev.* **D64** (2001) 044010 (12 pages).
293. Blagojević M. and Vasilic M., *Conservation laws in the teleparallel theory with a positive cosmological constant*, *Class. Quantum Grav.* **19** (2002) 3723.
294. Blagojević M. and Vasilic M., *Asymptotic symmetries in 3D gravity with torsion*, *Phys. Rev.* **D67** (2003) 084032 (14 pages).
295. Blagojević M. and Vasilic M., *Three-dimensional gravity with torsion as a Chern-Simons gauge theory*, *Phys. Rev.* **D68** (2003) 104023 (6 pages).
296. Blagojević M. and Vasilic M., *Asymptotic dynamics in 3D gravity with torsion*, *Phys. Rev.* **D68** (2003) 124007 (7 pages).
297. Blagojević M. and Vasilic M., *Anti-de Sitter 3-dimensional gravity with torsion*, *Mod. Phys. Lett.* **A20** (2005) 1285.
298. Blagojević M., Vasilic M., and Vukašinac T., *Asymptotic symmetry and conservation laws in the two-dimensional Poincaré gauge theory of gravity*, *Class. Quantum Grav.* **13** (1996) 3003-3019.
299. Bleecker D.D., *Gauge theory and variational theory* (Addison-Wesley Publ.: Reading, 1981) 179 p.
300. Bleecker D.D., *A new view of the ECSK theory with a Dirac spinor*, *Gen. Relat. Grav.* **22** (1990) 299-347.
301. Blome H.-J., Chicone C., Hehl F.W., and Mashhoon B., *Nonlocal modification of Newtonian gravity*, *Phys. Rev.* **D81** (2010) 065020 (11 pages).
302. Bloomer I., *A maximally symmetric space with torsion*, *Gen. Relat. Grav.* **9** (1978) 763-771.
303. Bloomer I. and Charap J.M., *The continuity of dislocations*, *J. Math. Phys.* **26** (1985) 2383-2385.
304. Bluhm R. and Kostelecky V.A., *Lorentz and CPT tests with spin-polarized solids*, *Phys. Rev. Lett.* **84** (2000) 1381-1384.
305. Boccaletti D., Agostini W., and Festa P., *Motion of spinning particles: Post-Newtonian approximation in the Einstein-Cartan theory*, *Nuovo Cim.* **B49** (1979) 45-54.
306. Böhmer C.G., *The Einstein static universe with torsion and the sign problem of the cosmological constant*, *Class. Quantum Grav.* **21** (2004) 1119.
307. Böhmer C.G., *On inflation and torsion in cosmology*, *Acta Phys. Pol.* **B36** (2005) 2841-2845.
308. Böhmer C.G., *The Einstein-Cartan-Elko system*, *Annalen der Physik (Berlin)* **16** (2007) 38-44.
309. Böhmer C.G., *Dark spinor inflation: Theory primer and dynamics*, *Phys. Rev.* **D77** (2008) 123535 (6 pages).
310. Böhmer C.G. and Burnett J., *Dark spinors with torsion in cosmology*, *Phys. Rev.* **D78** (2008) 104001 (6 pages).

311. Böhmer C.G., Burnett J., Mota D.F., and Shaw D.J., *Dark spinor models in gravitation and cosmology*, *JHEP* **7** (2010) 53.
312. Böhmer C.G., Downes R.J., and Vassiliev D., *Rotational elasticity*, *Quarterly Journal of Mechanics and Applied Mathematics* **64** (2011) 415-439.
313. Böhmer C.G., Harko T., and Lobo F.S.N., *Wormhole geometries in modified teleparallel gravity and the energy conditions*, *Phys. Rev.* **D85** (2012) 044033 (8 pages).
314. Böhmer C.G., Mussa A., and Tamanini N., *Existence of relativistic stars in  $f(T)$  gravity*, *Class. Quantum Grav.* **28** (2011) 245020.
315. Böhmer C.G., Neff P., and Seymenoğlu B., *Soliton-like solutions based on geometrically nonlinear Cosserat micropolar elasticity*, *Wave Motion* **60** (2016) 158-165.
316. Bogoyavlensky O.I., *On incompleteness of some solutions of the Einstein-Cartan equations with motion of matter*, *Phys. Lett.* **A60** (1977) 163-164.
317. Boldo J.L., Helayël-Neto J.A., and Panza N., *Propagating torsion in 3D gravity and dynamical mass generation*, *Class. Quantum Grav.* **19** (2002) 2201.
318. Boldo J.L., de Moraes L.M., and Helayël-Neto J.A., *Comments on topologically massive gravity with propagating torsion*, *Class. Quantum Grav.* **17** (2000) 813-823.
319. Boldo J.L. and Sasaki C.A.G., *Symmetry aspects of fermions coupled to torsion and electromagnetic fields*, *Europhys. Lett.* **59** (2002) 180.
320. Bonder Y., *Torsion or not torsion, that is the question*, *Int. J. Mod. Phys.* **D25** (2016) 1644013 (5 pages).
321. Borchenius K., *Symmetries in unified gauge theory with torsion*, *Nuovo Cim.* **A46** (1978) 403-418.
322. Borisov A.B., *The unitary representation of the general covariant group algebra*, *J. Phys.: Math. and Gen.* **A11** (1978) 1057-1067.
323. Borowiec A., Ferraris M., Francaviglia M., and Volovich I., *Energy-momentum complex for nonlinear gravitational Lagrangians in the 1st-order formalism*, *Gen. Relat. Grav.* **26** (1994) 637-645.
324. Borovko E.V., *Space-time with flat symmetry in Poincaré gauge gravitational theory*, *Sov. Phys. J.* **33** (1990) 361-366 [*Izvestia VUZov, Fizika*, n 4 (1990) 92-98 (in Russian)].
325. Borzeszkowski H.H. von, and Treder H.-J., *Classical gravity and quantum matter fields in unified theory*, *Gen. Rel. Grav.* **28** (1996) 1-14.
326. Borzeszkowski H.H. von, and Treder H.-J., *Mixed field theories and Weyl-Cartan geometry*, in: "Proc. of the 14th Course of Internat. School of Cosmology and Gravitation: Quantum Gravity" (Erice, 1995), Eds. P.G. Bergmann, V. de Sabbata, and H.-J. Treder (World Scientific: Singapore, 1996) 43-53.
327. Borzeszkowski H.H. von, and Treder H.-J., *The Weyl-Cartan space problem in purely affine theory*, *Gen. Rel. Grav.* **29** (1997) 455-466.
328. Borzeszkowski H.H. von, and Treder H.-J., *Torsion and the Weyl-Cartan problem in purely affine theory*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?", *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. de Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 9-32.
329. Borzeszkowski H.H. von, and Treder H.-J., *Spinorial matter in affine theory of gravity and the space problem*, *Gen. Rel. Grav.* **33** (2001) 1351-1369.
330. Borzou A., *A Lorentz gauge theory of gravity*, *Class. Quantum Grav.* **33** (2016) 025008 (16 pages).
331. Borzou A., *A macroscopically effective Lorentz gauge theory of gravity*, *Class. Quantum Grav.* **33** (2016) 235006 (19 pages).

332. Boulanger N. and Kirsch I., *Higgs mechanism for gravity. II. Higher spin connections*, *Phys. Rev.* **D73** (2006) 124023 (17 pages).
333. Boyadjiev T., Fiziev P., and Yazadjiev S., *Neutron star in the presence of a torsion-dilaton field*, *Class. Quantum Grav.* **16** (1999) 2359-2380.
334. Bradas J., Fennelly A., and Smalley L.L., *Torsion as a source of expansion in a Bianchi type-I universe in the self-consistent Einstein-Cartan theory of a perfect fluid with spin density*, *Phys. Rev.* **D35** (1987) 2302-2308.
335. Bradfield T., *Spontaneously broken Lorentz symmetry: an alternative model*, *Can. J. Phys.* **67** (1989) 881-883.
336. Brandt H.E., *Spacetime tangent bundle with torsion*, *Found. Phys. Lett.* **6** (1993) 339-369.
337. Brandt H.E., *Solution to torsion relations in Finsler-spacetime tangent bundle*, *Found. Phys. Lett.* **7** (1994) 297-301.
338. Brauns G., *Nichtlineare Spinorgleichungen und affiner Zusammenhang*, *Zeits. f. Naturforsch.* **19a** (1964) 825-827.
339. Brechet S.D., Hobson M.P., and Lasenby A.N., *Weyssenhoff fluid dynamics in general relativity using a 1 + 3 covariant approach*, *Class. Quantum Grav.* **24** (2007) 6329.
340. Brechet S.D., Hobson M.P., and Lasenby A.N., *Classical big-bounce cosmology: dynamical analysis of a homogeneous and irrotational Weyssenhoff fluid*, *Class. Quantum Grav.* **25** (2008) 245016.
341. Bregman A., *Weyl transformations and Poincaré gauge invariance*, *Progr. Theor. Phys.* **49** (1973) 667-692.
342. Bressan O., *Unified classical physics on a metric space with torsion*, In: "Relativity, supersymmetry and cosmology". *Proc. 5th Latin American Symposium on relativity and gravitation (SILARG V), Bariloche, Argentina, January 4-11, 1985*, Eds. O.J. Bressan, M.A. Castagnino, and V.H. Hamity. (Singapore: World Scientific, 1985) 232-242.
343. Bressan O., *Classical unified fields: physical space*, *Gen. Relat. Grav.* **18** (1986) 375-388.
344. Bressange G.F., *On the extension of the concept of thin shells to the Einstein-Cartan theory*, *Class. Quantum Grav.* **17** (2000) 2509-2523.
345. Brodskii A.M., Ivanenko D.D., and Sokolik H.A., *New treatment of the gravitational field*, *ZhETF* **41** (1962) 1307-1309 (in Russian).
346. Brodskii A.M., Ivanenko D.D., and Sokolik H.A., *A new conception of the gravitational field*, *Acta Phys. Hungarica* **14** (1962) 21-25.
347. Bronnikov K.A. and Galiakhmetov A.M., *Wormholes without exotic matter in Einstein-Cartan theory*, *Gravitation and Cosmology* **21** (2015) 283-288.
348. Bronnikov K.A. and Galiakhmetov A.M., *Wormholes and black universes without phantom fields in Einstein-Cartan theory*, *Phys. Rev.* **D94** (2016) 124006 (9 pages).
349. Bruggen M., *Effects of a torsion field on big bang nucleosynthesis*, *Gen. Rel. Grav.* **31** (1999) 1935-1939.
350. Bruhn G.W., Hehl F.W., and Jadczyk A., *Comments on "Spin connection resonance in gravitational general relativity"*, *Acta Phys. Pol.* **39** (2008) 51-58.
351. Brunelli R., Cognola G., and Zerbini S., *Heat-kernel expansion in a Riemann-Cartan spacetime*, In: "General relativity and gravitational physics". *Proc. 7th Italian Conf. Rapallo, Italy, September 3-7, 1986*, Eds. U. Bruzzo, R. Cianci, E. Massa. (Singapore: World Scientific, 1987) 269-273.
352. Bruzzo U., *The global Utiyama theorem in Einstein-Cartan theory*, *J. Math. Phys.* **28** (1987) 2074-2077.

353. Bruzzo U. and Cianci R., *Reduced fiber bundles and symmetries of gauge field theories. Rigid reductions and the Einstein-Cartan theory*, *Lett. Math. Phys.* **6** (1982) 471-477.
354. Buchdahl H.A., *Quadratic Lagrangians and Palatini's device*, *J. Phys. A: Math. and Gen.* **A12** (1979) 1229-1234.
355. Buchdahl H.A., *Representation of the Einstein-Proca field by an  $A_4^*$* , *J. Phys. A: Math. and Gen.* **A12** (1979) 1235-1238.
356. Buchdahl H.A., *On the compatibility of relativistic wave equations in Riemann-Cartan spaces*, *J. Math. Phys.* **30** (1989) 700-705.
357. Buchdahl H.A., *On the compatibility of relativistic wave equations in Riemann-Cartan spaces II*, *J. Math. Phys.* **31** (1990) 712-714.
358. Buchdahl H.A., *P-variations and asymmetric connexions*, *Nuovo Cim.* **B107** (1992) 1427-1435.
359. Bucataru I. and Epstein M., *Geometrical theory of dislocations in bodies with microstructure*, *J. Geom. Phys.* **52** (2004) 57-73.
360. Bugajska K., *From spinor structure to magnetic monopoles*, *J. Math. Phys.* **26** (1985) 2909-2914.
361. Bukhbinder I.L. and Odintsov S.D., *Nonsingular cosmological models with torsion determined by vacuum polarization of quantum fields*, *Pramana* **28** (1987) 241-245.
362. Bukhbinder I.L. and Odintsov S.D., *Effective action in the de Sitter space with torsion*, *Acta Phys. Pol.* **B18** (1987) 237-242.
363. Bukhbinder I.L. and Odintsov S.D., *The behavior of effective coupling constants in "finite" grand unification theories in curved space-time with torsion*, *Europhys. Lett.* **8** (1989) 595-598.
364. Bukhbinder I.L., Odintsov S.D., and Shapiro I.L., *Nonsingular cosmological model with torsion induced by vacuum quantum effects*, *Phys. Lett.* **B162** (1985) 92-96.
365. Bukhbinder I.L., Odintsov S.D., and Shapiro I.L., *Calculation of an effective potential in an external gravitational field with torsion and phase transitions induced by external fields*, *Sov. J. Phys.* **30** (1987) 183-189 [*Izvestiya VUZov, Fizika* n **3** (1987) 3-9 (in Russian)].
366. Bukhbinder I.L., Odintsov S.D., and Shapiro I.L., *Renormalization group approach to quantum field theory in curved space-time*, *Riv. Nuovo Cim.* **12** (1989) 1-112.
367. Bukhbinder I.L., Odintsov S.D., and Shapiro I.L., *Effective action in quantum gravity* (Institute of Physics Publishing: Bristol, 1992) 413 pp.
368. Bukhbinder I.L. and Shapiro I.L., *Renormalization of a quantum field theory model in a curved spacetime with torsion*, *Sov. J. Phys.* **28** (1985) 685-689 [*Izvestiya VUZov, Fizika*, n. 8 (1985) 94-100 (in Russian)].
369. Bukhbinder I.L. and Shapiro I.L., *Behavior of effective charges in a quantum field theory model in an external gravitational field with torsion*, *Sov. J. Phys.* **28** (1985) 1000-1003 [*Izvestiya VUZov, Fizika*, n. 12 (1985) 58-62 (in Russian)].
370. Bukhbinder I.L. and Shapiro I.L., *On the renormalization of models of quantum field theory in an external gravitational field with torsion*, *Phys. Lett.* **B151** (1985) 263-266.
371. Bukhbinder I.L. and Shapiro I.L., *On the renormalisation group equations in curved space-time with torsion*, *Class. Quantum Grav.* **7** (1990) 1197-1206.
372. Burdet G. and Perrin M., *Global formulation of the Fokker-Planck equation over Newton-Cartan and Einstein space-times*, *Europhys. Lett.* **12** (1990) 199-203.
373. Burnett J., Chervova O., and Vassiliev D., *Dirac equation as a special case of Cosserat elasticity*, in: "Analysis, Partial Differential Equations and Applications", The Vladimir Maz'ya Anniversary Volume, eds. A.Cialdea, F.Lanzara and P.E.Ricci, series Operator Theory: Advances and Applications (Birkhauser) **193** (2009) 15-29.

374. Burnett J. and Vassiliev D., *Weyl's Lagrangian in teleparallel form*, *J. Math. Phys.* **50** (2009) 102501.
375. Burnett J. and Vassiliev D., *Modeling the electron with Cosserat elasticity*, *Mathematika* **58** (2012) 349-370.
376. Burton D.A., Tucker R.W., and Wang C.H., *Spinning particles in scalar-tensor gravity*, *Phys. Lett.* **A372** (2008) 3141-3144.
377. Butcher L.M., Hobson M., and Lasenby A., *Localized energetics of linear gravity: Theoretical development*, *Phys. Rev.* **D86** (2012) 084013 (17 pages).
378. Bytsenko A.A., Elizalde E., and Odintsov S.D., *The renormalization group and the effective potential in a curved space-time with torsion*, *Progr. Theor. Phys.* **90** (1993) 677-688.

## C

379. Cacciatori S.L., Caldarelli M.M., Giacomini A., Klemm D., and Mansi D.S., *Chern-Simons formulation of three-dimensional gravity with torsion and nonmetricity*, *J. Geom. Phys.* **56** (2006) 2523-2543.
380. Cagnemi D., *One formulation for both lineal gravities through a dimensional reduction*, *Phys. Lett.* **B297** (1992) 261-265.
381. Cagnemi D. and Jackiw R., *Geometric gravitational force on particles moving in a line*, *Phys. Lett.* **B299** (1993) 24-29.
382. Cagnemi D. and Jackiw R., *Poincaré gauge theory for gravitational forces in (1+1) dimensions*, *Ann. Phys.* **225** (1993) 229-263.
383. Cai Y.-F., Capozziello S., De Laurentis M., and Saridakis E.N.,  *$f(T)$  teleparallel gravity and cosmology*, *Rep. Prog. Phys.* **79** (2016) 106901 (121 pages).
384. Cai Y.-F., Chen S.-H., Dent J.B., Dutta S., and Saridakis E.N., *Matter bounce cosmology with the  $f(T)$  gravity*, *Class. Quantum Grav.* **28** (2011) 215011.
385. Camacho A. and Camacho-Galván A., *Test of some fundamental principles in physics via quantum interference with neutrons and photons*, *Rep. Prog. Phys.* **70** (2007) 1937-1993.
386. Camacho A. and Macías A., *Space-time torsion contribution to quantum interference phases*, in: "Topics in Mathematical Physics, General Relativity and Cosmology in Honor of Jerzy Plebanski" *Proc. 2002 Int. Conf., Cinvestav, Mexico City, 17-20 Sept. 2002*, Eds. H. García-Compeán, B. Mielnik, M. Montesinos, and M. Przanowski (World Scientific: Singapore, 2002) 101-107.
387. Camacho A. and Macías A., *Space-time torsion contribution to quantum interference phases*, *Phys. Lett.* **B617** (2005) 118-123.
388. Camenzind M., *On the Yang-Mills structure of gravitation: a new issue of the final state*, *Gen. Relat. Grav.* **8** (1977) 103-108.
389. Camenzind M., *The gravitational field of spherically symmetric matter distributions in the Yang-Mills gauge theory of gravity*, *Phys. Lett.* **A63** (1977) 69-72.
390. Camenzind M., *Homogeneous and isotropic world models in the Yang-Mills dynamics of gravity*, *J. Math. Phys.* **19** (1978) 624-634.
391. Camenzind M., *On the way towards a nonsingular gravitational collapse – The properties of the Yang-Mills cosmos*, *Gen. Relat. Grav.* **8** (1978) 661-671.
392. Camenzind M., *Lepton pair creation and the uniform gravitational collapse*, *Nuovo Cim.* **B46** (1978) 153-166.
393. Camenzind M., *Weak and strong sources of gravity –  $SO(1,3)$ -gauge theory of gravity*, *Phys. Rev.* **D18** (1978) 1068-1081.

394. Camera S., Cardone V.F., and Radicella N., *Detectability of torsion gravity via galaxy clustering and cosmic shear measurements*, *Phys. Rev.* **D89** (2014) 083520 (15 pages).
395. Campigotto M. and Fatibene L., *Conformal gravity as a Gauge natural theory*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650052 (8 pages).
396. Camporesi R., Destri C., Melegari G., and Orzalesi C.A., *Kaluza-Klein spectrum on a contorted vacuum*, *Class. Quantum Grav.* **2** (1985) 461-686.
397. Canale A., De Ritis R., and Tarantino C., *Cosmological solutions of a quadratic theory of gravity with torsion*, *Phys. Lett.* **A100** (1984) 178-181.
398. Canarutto D. and Jadczyk A., *Two-spinors and Einstein-Cartan-Maxwell-Dirac fields*, *Nuovo Cim.* **B113** (1998) 49-67.
399. Cant A. and Ne'eman Y., *Spinorial infinite equations fitting metric-affine gravity*, *J. Math. Phys.* **26** (1985) 3180-3189.
400. Cantcheff M.B., *Einstein-Cartan formulation of Chern-Simons Lorentz-violating gravity*, *Phys. Rev.* **D78** (2008) 025002 (8 pages).
401. Cao Z.-B., and Duan Y.-S., *Extra dimensions with nontrivial torsion and spins of cosmic strings*, *Int. J. Mod. Phys.* **A24** (2009) 1147.
402. Cappelli A., Ciafaloni M., and Valtancoli P., *Classical scattering in 2 + 1 gravity with N-spinning sources*, *Phys. Lett.* **B273** (1991) 431-437.
403. Capozziello S., *Torsion quintessence*, *Mod. Phys. Lett.* **A17** (2002) 1621.
404. Capozziello S., Basini G., and De Laurentis M., *Deriving the mass of particles from extended theories of gravity in LHC era*, *Eur. Phys. J.* **C71** (2011) 1679.
405. Capozziello S., Cardone V.F., Carloni S., and Troisi A., *Torsion  $\Lambda$  term*, in: "Thinking, Observing and Mining the Universe" *Proc. Int. Conf., Sorrento, Italy, 22-27 Sept. 2003*, Eds. G. Miele and G. Longo (World Scientific: Singapore, 2004) 309-310.
406. Capozziello S., Cardone V.F., Farajollahi H., and Ravanpak A., *Cosmography in  $f(T)$  gravity*, *Phys. Rev.* **D84** (2011) 043527 (11 pages).
407. Capozziello S., Cardone V.F., Piedipalumbo E., Sereno M., and Troisi A., *Matching torsion  $\Lambda$ -term with observations*, *Int. J. Mod. Phys.* **D12** (2003) 381.
408. Capozziello S., Cianci R., Stornaiolo C., and Vignolo S.,  *$f(R)$  gravity with torsion: the metric-affine approach*, *Class. Quantum Grav.* **24** (2007) 6417.
409. Capozziello S., Cianci R., Stornaiolo C., and Vignolo S.,  *$f(R)$  cosmology with torsion*, *Phys. Scr.* **78** (2008) 065010.
410. Capozziello S., Cirilo-Lombardo D.J., and Dorokhov A.E., *Fermion interactions, cosmological constant and space-time dimensionality in a unified approach based on affine geometry*, *Int. J. Theor. Phys.* **53** (2014) 3882-3892.
411. Capozziello S. and De Laurentis M., *Gravity from local Poincaré gauge invariance*, *Int. J. Geom. Methods Mod. Phys.* **06** (2009) 1-24.
412. Capozziello S. and De Laurentis M., *A review about invariance induced gravity: gravity and spin from local conformal-affine symmetry*, *Found. Phys.* **40** (2010) 867-899.
413. Capozziello S. and De Laurentis M., *Metric and connections in theories of gravity. The role of equivalence principle*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1640007 (7 pages).
414. Capozziello S., De Laurentis M., Fabbri L., and Vignolo S., *Running coupling in electroweak interactions of leptons from  $f(R)$ -gravity with torsion*, *Eur. Phys. J.* **C72** (2012) 1908.
415. Capozziello S., Fabbri L., and Vignolo S., *Weak forces and neutrino oscillations under the standards of hybrid gravity with torsion*, *Mod. Phys. Lett.* **A28** (2013) 1350155 (20 pages).

416. Capozziello S. and Francaviglia M., *Extended theories of gravity and their cosmological and astrophysical applications*, *Gen. Rel. Grav.* **40** (2008) 357-420.
417. Capozziello S., González P.A., Saridakis E.N., and Vásquez Y., *Exact charged black-hole solutions in  $D$ -dimensional  $f(T)$  gravity: torsion vs curvature analysis*, *JHEP* **2** (2013) 39.
418. Capozziello S., Iovane G., Lambiase G., and Stornaiolo C., *Fermion helicity flip induced by torsion field*, *Europhys. Lett.* **46** (1999) 710-715.
419. Capozziello S. and Lambiase G., *Berry's phase of neutrino oscillations in the presence of torsion*, *Europhys. Lett.* **52** (2000) 15-21.
420. Capozziello S., Lambiase G., and Stornaiolo C., *Berry and geometrical phase induced by torsion field*, *Europhys. Lett.* **48** (1999) 482-485.
421. Capozziello S., Lambiase G., and Stornaiolo C., *Geometric classification of the torsion tensor of space-time*, *Ann. d. Phys. (Leipzig)* **10** (2001) 713-727.
422. Capozziello S., Luongo O., and Saridakis E.N., *Transition redshift in  $f(T)$  cosmology and observational constraints*, *Phys. Rev.* **D91** (2015) 124037 (11 pages).
423. Capozziello S. and Stornaiolo C., *Scalar perturbations in a cosmological model with torsion*, *Nuovo Cim.* **B113** (1998) 879-886.
424. Capozziello S. and Stornaiolo C., *Torsion tensor and its geometric interpretation*, *Annales de la Fondation L. de Broglie* **32** (2007) 195-214.
425. Capozziello S. and Vignolo S., *Metric-affine  $f(R)$ -gravity with torsion: an overview*, *Ann. d. Phys. (Berlin)* **19** (2010) 238-248.
426. Capriotti S., *Differential geometry, Palatini gravity and reduction*, *J. Math. Phys.* **55** (2014) 012902 (29 pages).
427. Cardone V.F., Radicella N., and Camera S., *Accelerating  $f(T)$  gravity models constrained by recent cosmological data*, *Phys. Rev.* **D85** (2012) 124007 (9 pages).
428. Carlevaro N., Lecian O.M., and Montani G., *Macroscopic and microscopic paradigms for the torsion field: from the test-particle motion to a Lorentz gauge theory*, *Annales de la Fondation L. de Broglie* **32** (2007) 281-295.
429. Carlevaro N., Lecian O.M., and Montani G., *Lorentz gauge theory and spinor interaction*, *Int. J. Mod. Phys.* **A23** (2008) 1282-1285.
430. Carlevaro N., Lecian O.M., and Montani G., *Fermion dynamics by internal and spacetime symmetries*, *Mod. Phys. Lett.* **A24** (2009) 415.
431. Carlevaro N. and Montani G., *A novel approach to Lorentz gauge theory*, in: *Proc. Third Stueckelberg Workshop on "Relativistic Field Theories"*, Eds. N. Carlevaro, G.V. Vereshchagin, and R. Ruffini (Cambridge Scientific Publishers: Cambridge, 2010) p. 149.
432. Carloni S., Lobo F.S.N., Otalora G., and Saridakis E.N., *Dynamical system analysis for a nonminimal torsion-matter coupled gravity*, *Phys. Rev.* **D93** (2016) 024034 (12 pages).
433. Carloni S., Vignolo S., and Fabbri L., *The dynamics of Bianchi I universes in  $\mathcal{R}^n$  cosmologies with torsion*, *Class. Quantum Grav.* **30** (2013) 205010.
434. Carmeli M., *Integral formalism of gauge fields and general relativity*, *Phys. Rev. Lett.* **36** (1974) 59-61.
435. Carmeli M. and Kaye M., *Gauge group theory of combined gravitational and electromagnetic interactions*, *Lett. Nuovo Cim.* **17** (1976) 275-278.
436. Carmeli M. and Malin S., *Reformulation of general relativity as a gauge theory*, *Ann. Phys.* **103** (1977) 208-232.
437. Carroll S.M. and Field G.B., *Consequences of propagating torsion in connection dynamic theories of gravity*, *Phys. Rev.* **D50** (1994) 3867-3873.

438. Cartan É., *Sur une généralisation de la notion de courbure de Riemann et les espaces à torsion*, *C.R. Acad. Sci. (Paris)* **174** (1922) 593-595; English translation: *On a generalization of the notion of Riemann curvature and spaces with torsion*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 489-491.
439. Cartan É., *Sur les variétés à connexion affine et la théorie de la relativité généralisée, 1 partie*, *Ann. Éc. Norm.* **40** (1923) 325-412.
440. Cartan É., *Sur les variétés à connexion affine et la théorie de la relativité généralisée, 1 partie, suite*, *Ann. Éc. Norm.* **41** (1924) 1-25.
441. Cartan É., *Sur les variétés à connexion affine et la théorie de la relativité généralisée, 2 partie*, *Ann. Éc. Norm.* **42** (1925) 17-88.
442. Cartan É., *On manifolds with an affine connection and the theory of general relativity* (Bibliopolis: Napoli, 1986) 200 p.
443. Cartas-Fuentevilla R., *BF-like symmetries for general relativity and Ricci flows on the space of connections*, *Phys. Rev.* **D80** (2009) 085003 (7 pages).
444. Cartas-Fuentevilla R., *Fully torsion-based formulation for curved manifolds*, *Int. J. Geom. Methods Mod. Phys.* **10** (2013) 1350021 (32 pages).
445. Cartas-Fuentevilla R., *Twisted entropy functional for three-manifolds and flows induced on the boundary*, *Int. J. Geom. Methods Mod. Phys.* **11** (2014) 1450033 (30 pages).
446. Cartas-Fuentevilla R., Solano-Altamirano J.M., and Enriquez-Silverio P., *Post-Riemannian approach for the symplectic and elliptic geometries of gravity*, *J. Phys. A: Math. and Theor.* **44** (2011) 195206.
447. Carvalho A.M.D., Furtado C., and Moraes F., *Self-forces on electric and magnetic linear sources in the presence of a torsional defect*, *Phys. Rev.* **D62** (2000) 067504 (4 pages).
448. Casana R., Fainberg V.Ya., Lunardi J.T., Pimentel B.M., and Teixeira R.G., *Massless DKP fields in Riemann-Cartan spacetimes*, *Class. Quantum Grav.* **20** (2003) 2457.
449. Casanova S., Montani G., Ruffini R., and Zalaletdinov R., *On the non-Riemannian manifolds as framework for geometric unification theories: Affine connection geometry with asymmetric metric*, in: Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories" (Jerusalem, 22-27 June 1997). Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 436-438.
450. Casanova S., Lecian O.M., Montani G., Ruffini R., and Zalaletdinov R., *Extended Schouten classification for non-Riemannian geometries*, *Mod. Phys. Lett.* **A23** (2008) 17-23.
451. Castagnino M. and Levinas M.L., *On the post-Newtonian approximation of the ECSK theory. II*, *Gen. Relat. Grav.* **19** (1987) 545-562.
452. Castagnino M. and Levinas M., *Photon motion in the ECSK theory*, *Gen. Relat. Grav.* **20** (1988) 715-721.
453. Castagnino M., Levinas M.L., and Umerez N., *On the post-Newtonian approximation of the ECSK theory*, *Gen. Relat. Grav.* **17** (1985) 683-693.
454. Castagnino M., Levinas M.L., and Umerez N., *Energy-momentum and equivalence principle in non-Riemannian geometries*, *Gen. Rel. Grav.* **29** (1997) 691-703.
455. Castagnino M., Levinas M.L., N Umerez N., *Gravitational and matter energy-momentum densities and equivalence principle in non-Riemannian geometries*, *Int. J. Mod. Phys.* **A14** (1999) 4721-4734.
456. Castillo-Felisola O., Corral C., Kovalenko S., Schmidt I., and Lyubovitskij V.E., *Axions in gravity with torsion*, *Phys. Rev.* **D91** (2015) 085017 (7 pages).

457. Castillo-Felisola O., Corral C., Schmidt I., and Zerwekh A.R., *Updated limits on extra dimensions through torsion and LHC data*, *Mod. Phys. Lett.* **A29** (2014) 1450081 (9 pages).
458. Castillo-Felisola O., Corral C., Villavicencio C., and Zerwekh A.R., *Fermion masses through condensation in spacetimes with torsion*, *Phys. Rev.* **D88** (2013) 124022 (7 pages).
459. Castro C., *A supersymmetric Lagrangian for Poincaré gauge theories of gravity*, *Progr. Theor. Phys.* **82** (1989) 616-630.
460. Castro C. and Pavšič M., *Higher derivative gravity and torsion from the geometry of C-spaces*, *Phys. Lett.* **B539** (2002) 133-142.
461. Catren G., *Geometric foundations of Cartan gauge gravity*, *Int. J. Geom. Methods Mod. Phys.* **12** (2015) 1530002 (33 pages).
462. Cebeci H. and Dereli T., *Conformal black hole solutions of axidilaton gravity in D dimensions*, *Phys. Rev.* **D65** (2002) 047501 (4 pages).
463. Cebeci H. and Dereli T., *Axi-dilaton gravity in  $D \geq 4$  dimensional space-times with torsion*, *Phys. Rev.* **D71** (2005) 024016 (7 pages).
464. Cembranos J.A.R., and Valcarcel J.G., *New torsion black hole solutions in Poincaré gauge theory*, *JCAP* **01** (2017) 014 (17 pages).
465. Challinor A., Lasenby A., Doran C., and Gull S., *Massive, non-ghost solutions for the Dirac field coupled self-consistently to gravity*, *Gen. Rel. Grav.* **29** (1997) 1527-1544.
466. Chandia O. and Zanelli J., *Topological invariants, instantons and chiral anomaly on spaces with torsion*, *Phys. Rev.* **D55** (1997) 7580-7585.
467. Chandia O. and Zanelli J., *Reply to "Comment on 'Topological invariants, instantons, and the chiral anomaly on spaces with torsion'"*, *Phys. Rev.* **D63** (2001) 048502 (4 pages).
468. Chang C.-C., Nester J.M., and Chen C.-M., *Pseudotensors and quasilocal energy-momentum*, *Phys. Rev. Lett.* **83** (1999) 1897-1901.
469. Chang D.B. and Johnson H.H., *No black holes: a gravitational gauge theory possibility*, *Phys. Lett.* **A77** (1980) 411-415.
470. Chang D.B. and Johnson H.H., *Theory of gravitation with an alternative to black holes*, *Phys. Rev.* **D21** (1980) 874-883.
471. Chang L.N., Lebedev O., Loinaz W., and Takeuchi T., *Universal torsion-induced interaction from large extra dimensions*, *Phys. Rev. Lett.* **85** (2000) 3765-3768.
472. Chang L.N. and Chopin Soo C., *Massive torsion modes, chiral gravity and the Adler-Bell-Jackiw anomaly*, *Class. Quantum Grav.* **20** (2003) 1379.
473. Changgui S., Cunhon W., and Dehnen H., *De Sitter gauge theory of gravity*, *Int. J. Theor. Phys.* **29** (1990) 885-900.
474. Changgui S. and Dehnen H., *Nonlinear gauge theory of Poincaré gravity*, *Int. J. Theor. Phys.* **30** (1991) 1489-1496.
475. Changgui S. and Xu B.-Q., *ISO(3,1) gauge theory of gravity*, *Int. J. Theor. Phys.* **25** (1986) 347-353.
476. Charap J.M. and Tait W., *A gauge theory of the Weyl group*, *Proc. Roy. Soc. Lond.* **A340** (1974) 249-262.
477. Charap J.M. and Bloomer I., *Spin, torsion and the singularity theorems*, *JINR report* **E2-10178** (1976) 1-8.
478. Chatterjee P., Bandyopadhyay A.K., and Bandyopadhyay P., *SL(2,C)-gauge theory and N=1 supergravity*, *Gen. Relat. Grav.* **18** (1986) 1127-1140.
479. Chatterjee P., and Bhattacharya B., *Space-time torsion, broken Lorentz symmetry and inflation in the early universe*, *Mod. Phys. Lett.* **A8** (1993) 2249-2257.

480. Chauveheid J., *A proposal for a quadratic electromagnetic coupling based on the underlying philosophy of Einstein-Maxwell theory*, *Phys. Essays* **10** (1997) 474-482.
481. Chee Guoying, *A connection dynamic theory of gravity*, *Gen. Relat. Grav.* **28** (1996) 549-563.
482. Chee Guoying, *A Self-dual and anti-self-dual structure of the de Sitter gauge theory and Ashtekar's variables*, *Phys. Rev.* **D54** (1996) 6552-6554.
483. Chee Guoying and Guo Y., *Exact de Sitter solutions in quadratic gravitation with torsion*, *Class. Quantum Grav.* **29** (2012) 235022 (22 pages).
484. Chen C.-M., Chern D.-C., Nester J.M., Yang P.-K., and Zhytnikov V.V., *Poincaré gauge theory Schwarzschild-de Sitter solutions with long range spherically symmetric torsion*, *Chinese J. Phys.* **32** (1994) 29-40.
485. Chen C.-M., Nester J.M., and Tung R.-S., *Quasilocal energy-momentum for geometrical gravity theories*, *Phys. Lett.* **A203** (1995) 5-11.
486. Chen C.-M., Nester J.M., and Tung R.-S., *Gravitational energy for GR and Poincaré gauge theories: A covariant Hamiltonian approach*, *Int. J. Mod. Phys.* **D24** (2015) 11 (73 pages).
487. Chen Fang Pei, *General equations of motion for test particles in space-times with torsion*, *Int. J. Theor. Phys.* **29** (1990) 161-171.
488. Chen Fang Pei, *Momentum, angular momentum and equations of motion for test bodies in space-time with torsion*, *Int. J. Theor. Phys.* **32** (1993) 373-387.
489. Chen Fang Pei, *Field equations and conservation laws derived from the generalized Einstein's Lagrangian density for a gravitational system and their influences upon cosmology*, *Int. J. Theor. Phys.* **47** (2008) 421-437.
490. Chen H., Ho F.-H., Nester J.M., Wang C.-H., and Yo H.-J., *Cosmological dynamics with propagating Lorentz connection modes of spin zero*, *JCAP* **10** (2009) 027.
491. Chen H., Nester J.M., and Yo H.-J., *Acausal PGT modes and the nonlinear constraint effect*, *Acta Phys. Pol.* **29** (1998) 961-970.
492. Chen H.H., Chen M.-Q., and Chern D.-C., *Nonanalyticity of the coupling constant in the vacuum solutions of the general  $R + R^2$  theories of gravity with torsion*, *Chinese J. Phys.* **20** (1982) 102-105.
493. Chen H.H., Chern D.-C., Hsu R.R., and Yeung W.B., *Rotonless solutions in the Poincaré gauge theory of gravitation*, *Phys. Lett.* **A111** (1985) 693-642.
494. Chen H.H., Chern D.-C., Hsu R.R., Nester J.M., and Yeung W.B., *Asymptotically Newtonian conditions for Poincaré gauge theory*, *Progr. Theor. Phys.* **79** (1988) 77-85.
495. Chen H.H., Hsu R.R., and Yeung W.B., *Expanding universe solutions for the Yang-Mills-Higgs system coupled to the Poincaré gauge theory of gravitation*, *Class. Quant. Grav.* **2** (1985) 927-934.
496. Chen M.-Q., Chern D.-C., Hsu R.R., and Yeung W.B., *Plane-fronted torsion waves in a gravitational gauge theory with a quadratic Lagrangian*, *Phys. Rev.* **D28** (1983) 2094-2095.
497. Chen M.-Q., Chern D.-C., Hsu R.R., and Yeung W.B., *Some vacuum solutions for the torsion and metric tensors in the Poincaré gauge field theory of gravitation*, *Proc. Nat. Sci. Council, Part A: Phys. Sci. and Engineering* **9** (1985) 10-13.
498. Chen M.-Q., Hsu R.-R., and Yeung W.B., *Spontaneous compactification in six-dimensional Poincaré gauge field theory of gravitation*, *Phys. Lett.* **B140** (1984) 201-204.
499. Chen S. and Jing J., *The torsion cosmology in Kaluza-Klein theory*, *JCAP* **09** (2009) 001.
500. Chen S.-H., Dent J.B., Dutta S., and Saridakis E.N., *Cosmological perturbations in  $f(T)$  gravity*, *Phys. Rev.* **D83** (2011) 023508 (11 pages).
501. Chen Z.-C., Wu Y., and Wei H., *Post-Newtonian approximation of teleparallel gravity coupled with a scalar field* *Nucl. Phys.* **B894** (2015) 422-438.

502. Cheng K.S. and Ni W.T., *Conditions for the local existence of metric in a generic affine manifold*, *Math. Proc. Camb. Phil. Soc.* **87** (1980) 527-534.
503. Cheng K.S., Harko T., and Wang X.Y., *Radiation transport equations in non-Riemannian space-times* *Phys. Rev.* **D71** (2005) 103001 (11 pages).
504. Cheng W.-H., Chern D.-C., and Nester J.M., *Canonical analysis of the one parameter teleparallel theory*, *Phys. Rev.* **D38** (1988) 2656-2658.
505. Chern D.-C., *On the parallel transport of tetrad in a Riemann-Cartan spacetime*, *Chinese J. Phys.* **19** (1981) 45-47.
506. Chern D.-C. and Nester J.M., *Positive energy and Poincaré gauge theory*, in: "Gravitational Collapse and Relativity. Proc. of XIV Yamada Conf." (Kyoto, 1986), Eds. H. Sato and T. Nakamura (World Scientific: Singapore, 1986) 106-113.
507. Chern D.-C., Nester J.M., and Yo H.-J., *Positive energy test of Poincaré gauge theory*, *Int. J. Mod. Phys.* **A7** (1992) 1993-2003.
508. Chernikov N.A., *Tensor and covector of torsion in the unified field theory*, in: "Nonlocal, nonlinear and nonperturbative field theories" (JINR: Dubna, 1976) **D2-9788**, 120-140 (in Russian).
509. Chernikov N.A., *Space of affine connection with non-symmetrical tensor field*, Preprint (JINR: Dubna, 1977) **P2-11021**, 17 p.
510. Chervova O., and Vassiliev D., *The stationary Weyl equation and Cosserat elasticity*, *J. Phys. A: Math. Theor.* **43** (2010) 335203.
511. Chkareuli J.L., *On the origin of Poincaré gauge gravity*, *Phys. Lett.* **B769** (2017) 377-384.
512. Chkareuli J.L., *Poincaré gauge gravity: An emergent scenario*, *Phys. Rev.* **D95** (2017) 084051 (19 pages).
513. Chmielowski P., *Matching and junction conditions in the Poincaré gauge theory of gravity with quadratic Lagrangian*, *Progr. Theor. Phys.* **88** (1992) 691-702.
514. Cho J.H. and Lee H.J., *Lorentzian approach to the Chern-Simons coefficient quantization in the ISO(2,1) gauge gravity*, *J. Korean Phys. Soc. Suppl.* **33** (1998) S471-S476.
515. Cho Y.M., *Higher-dimensional unifications of gravitation and gauge theories*, *J. Math. Phys.* **16** (1975) 2029-2035.
516. Cho Y.M., *Gauge theory of Poincaré symmetry*, *Phys. Rev.* **D14** (1976) 3335-3340.
517. Cho Y.M., *Gauge theory, gravitation, and symmetry*, *Phys. Rev.* **D14** (1976) 3341-3344.
518. Cho Y.M., *Einstein Lagrangian as the translational Yang-Mills Lagrangian*, *Phys. Rev.* **D14** (1976) 2521-2525.
519. Cho Y.M., *A generalization of Cartan's theory of gravitation*, *J. Phys.* **A11** (1978) 2385-2387.
520. Cho Y.M., and Jang P.S., *Unified geometry of internal space with space-time*, *Phys. Rev.* **D12** (1975) 3789-3792.
521. Cho Y.M. and Freund P.G.O., *Non-abelian gauge fields as Nambu-Goldstone fields*, *Phys. Rev.* **D12** (1975) 1711-1720.
522. Cho Y.M. and Pak D.G., *Vacuum tunneling in gravity*, *Class. Quantum Grav.* **28** (2011) 155008.
523. Cho Y.M., Pak D.G., and Park D.H., *A minimal model of Lorentz gauge gravity with dynamical torsion*, *Int. J. Mod. Phys.* **A25** (2010) 2867.
524. Cho Y.M. and Park D.H., *Closed time-like curves and weak energy condition*, *Phys. Lett.* **B402** (1997) 18-24.

525. Cho Y.M., Soh K.S., Yoon J.H., and Park Q.-H., *Gravitation as a gauge theory of the diffeomorphism group*, *Phys. Lett.* **B286** (1992) 251-255.
526. Choquet-Bruhat Y., *Einstein-Cartan theory with spin 3/2 source and classical supergravity*, in: "On Relativity Theory. Proc. of the Sir A.Eddington Centenary Symp.", Eds. Y. Choquet-Bruhat and T.M. Karade (World Scientific: Singapore, 1985) vol. **2**, 27-56.
527. Choquet-Bruhat Y., *Spin 1/2 fields in arbitrary dimensions and the Einstein-Cartan theory*, In: "Gravitation and geometry: a volume in honor of Ivor Robinson", Eds. W. Rindler, and A. Trautman (Naples: Bibliopolis, 1987) 83-106.
528. Christensen S.M., *Second and fourth order invariants on curved manifold with torsion*, *J. Phys.* **A13** (1980) 3001-3009.
529. Chrusciel P.T., *On the unified affine electromagnetism and gravitation theories*, *Acta Phys. Pol.* **B15** (1984) 35-51.
530. Cianci R., *On generalized torsion two-forms in general relativity. A result of null tetrad formalism*, *Lett. Math. Phys.* **5** (1981) 481-487.
531. Cianci R., Fabbri L., and Vignolo S., *Exact solutions for Weyl fermions with gravity*, *Eur. Phys. J.* **C75** (2015) 478 (10 pages).
532. Cirilo-Lombardo D.J., *On unified field theories, dynamical torsion and geometrical models*, *Phys. Part. Nucl. Lett.* **7** (2010) 299-307.
533. Cirilo-Lombardo D.J., *Unified field theoretical models from generalized affine geometries*, *Int. J. Theor. Phys.* **49** (2010) 1288-1301.
534. Cirilo-Lombardo D.J., *Unified field theoretical models from generalized affine geometries. II*, *Int. J. Theor. Phys.* **50** (2011) 1699-1708.
535. Cirilo-Lombardo D.J., *Unified field theoretical models from generalized affine geometries. III*, *Int. J. Theor. Phys.* **50** (2011) 3621-3634.
536. Cirilo-Lombardo D.J., *On unified field theories, dynamical torsion and geometrical models: II*, *Phys. Part. Nucl. Lett.* **8** (2011) 507-511.
537. Ciubotariu C., Ciubotariu C.-I., and Ciubotariu C., *Plasma, gravitation, and force-free geometries*, *Gen. Rel. Grav.* **27** (1995) 129-141.
538. Clerc R.-L., *Équations d'Einstein sur un espace à connexion non riemannienne*, *C. R. Acad. Sci. Paris, Ser. A* **272** (1971) 1145-1148.
539. Clerc R.-L., *Équations des trajectoires sur un espace à connexion non riemannienne*, *C. R. Acad. Sci. Paris, Ser. A* **272** (1971) 1760-1763.
540. Clerc R.-L., *Une structure de l'espace-temps associée à un tenseur impulsion-énergie asymétrique*, *C. R. Acad. Sci. Paris, Ser. A* **274** (1972) 525-528.
541. Cognola G. and Giacconi P., *The nonabelian covariant anomaly in the presence of torsion*, In: "General relativity and gravitational physics" Proc. 8th Italian Conf., Cavalese, Italy, August 30 - September 3, 1988, Eds. M. Cerdonio, M. Toller, R. Cianci, and M. Francaviglia. (Singapore: World Scientific, 1989) 599-603.
542. Cognola G. and Giacconi P., *Consistent anomalies in the presence of torsion*, In: "Recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories". Proc. of 5th Marcel Grossmann meeting, Perth, Australia, August 8-13, 1988, Eds. D.G. Blair, and M.J. Buckingham (Singapore: World Scientific, 1989) Pt. A, 809-812.
543. Cognola G. and Giacconi P., *Non Abelian anomalies on a curved space with torsion*, *Phys. Rev.* **D39** (1989) 2987-2992.
544. Cognola G., Giacconi P., and Soldati R., *Effective actions with vector-axial vector couplings on Riemann-Cartan manifolds*, *J. Math. Phys.* **31** (1990) 2699-2701.

545. Cognola G., Soldati R., Vanzo L., and Zerbini S., *Lagrangian formulation of a spinning test particle in a curved space-time with torsion*, Phys. Rev. **D25** (1981) 3109-3116.
546. Cognola G., Soldati R., Vanzo L., and Zerbini S., *Lagrangian dynamics of a classical spinning particle with dipole moment*, Nuovo Cim. **B76** (1983) 109-129.
547. Cognola G., Vanzo L., and Zerbini S., *Sum rules for spectral densities of differential operators and generalized heat kernel expansion*, Phys. Lett. **B223** (1989) 416-420.
548. Cognola G. and Zerbini S., *Heat kernel expansion in geometric fields*, Phys. Lett. **B195** (1987) 435-438.
549. Cognola G. and Zerbini S., *Seeley-DeWitt coefficients in a Riemann-Cartan manifold*, Phys. Lett. **B214** (1988) 70-74.
550. Cognola G. and Zerbini S., *Some physical applications of the heat kernel expansion*, Mod. Phys. Lett. **A3** (1988) 599.
551. Coley A., *Analysis of nonmetric theories of gravity. I. Electromagnetism*, Phys. Rev. **D27** (1983) 728-739.
552. Coley A., *Analysis of nonmetric theories of gravity. II. The weak equivalence principle*, Phys. Rev. **D28** (1983) 1829-1843.
553. Coley A., *Analysis of nonmetric theories of gravity. III. Summary of the analysis and its application to theories in the literature*, Phys. Rev. **D28** (1983) 1844-1852.
554. Coley A., *A note on the geometric unification of gravity and electromagnetism*, Gen. Relat. Grav. **16** (1984) 459-464.
555. Coley A. and Tremaine S., *Constraint on nonmetric theories of gravity from supernova 1987A*, Phys. Rev. **D38** (1988) 2927-2929.
556. Concha P.K., Peñafiel D.M., Rodríguez E.K., and Salgado P., *Generalized Poincaré algebras and Lovelock-Cartan gravity theory*, Phys. Lett. **B742** (2015) 310-316.
557. Corichi A. and Reyes J.D., *The gravitational Hamiltonian, first order action, Poincaré charges and surface terms*, Class. Quantum Grav. **32** (2015) 195024 (55 pages).
558. Corichi A., Rubalcava-García I., and Vukašinac T., *Actions, topological terms and boundaries in first-order gravity: A review*, Int. J. Mod. Phys. **D25** (2016) 1630011 (54 pages).
559. Cosserat E., and Cosserat F., *Theorie des corps deformables* (Hermann: Paris, 1909) 226 pp.
560. Costa de Beauregard O., *Sur la dynamique des milieux doués d'une densité de moment cinétique propre*, C.R. Acad. Sci. (Paris) **214** (1942) 904.
561. Costa de Beauregard O., *Translational inertial spin effect*, Phys. Rev. **129** (1963) 466-471.
562. Costa de Beauregard O., *Translational inertial spin effect*, in: "Perspectives in Geometry and Relativity", Ed. B. Hoffmann (Indiana Univ. Press: Bloomington, 1966) 44-57.
563. Crawford J.P., *Spinor metrics, spin connection compatibility and spacetime geometry from spin geometry*, Class. Quantum Grav. **20** (2003) 2945.
564. Cvetković B. and Blagojević M., *Stability of 3D black hole with torsion*, Mod. Phys. Lett. **A22** (2007) 3047.
565. Cvetković B. and Simić M., *5D Lovelock gravity: New exact solutions with torsion*, Phys. Rev. **D94** (2016) 084037 (9 pages).

## D

566. D'Adda A., *On the construction of gravity and supergravity from fundamental spinorial constituents*, *Phys. Lett.* **B119** (1982) 334-338.
567. Dadhich N. and Pons J.M., *On the equivalence of the Einstein-Hilbert and the Einstein-Palatini formulations of general relativity for an arbitrary connection*, *Gen. Rel. Grav.* **44** (2012) 2337-2352.
568. Dai M.-C., Nester J.M., and Chern D.-C., *Anisotropic cosmological models in Poincaré gauge gravity*, *Chinese J. Phys.* **28** (1990) 473-480.
569. Dalakov P. and Ivanov S., *Harmonic spinors of the Dirac operator of connection with torsion in dimension four*, *Class. Quantum Grav.* **18** (2001) 253-263.
570. Damour T., Deser S., and McCarthy J., *Nonsymmetric gauge theories: Inconsistencies and a cure*, *Phys. Rev.* **D47** (1993) 1541-1556.
571. Darabi F., Mousavi M., and Atazadeh K., *Geodesic deviation equation in  $f(T)$  gravity*, *Phys. Rev.* **D91** (2015) 084023 (11 pages).
572. Dariescu C. and Gottlieb I., *Up to the second order spin contributions of the electrostatic field: Reissner-Nordstrom solution representing  $U_4$ -spacetime*, *Tensor N.S.* **54** (1993) 35-45.
573. Da Rocha R. and Pereira J.G., *The quadratic spinor Lagrangian, axial torsion current and generalizations*, *Int. J. Mod. Phys.* **D16** (2007) 1653 (15 pages).
574. Da Rocha Neto J.F., Maluf J.W., and Ulhoa S.C., *Hamiltonian formulation of unimodular gravity in the teleparallel geometry*, *Phys. Rev.* **D82** (2011) 124035 (9 pages).
575. Das A., Mukhopadhyaya B., and SenGupta S., *Why has spacetime torsion such negligible effect on the Universe?* *Phys. Rev.* **D90** (2014) 107901 (4 pages).
576. Das M., Mohanty S., and Prasanna A.R., *Constraints on background torsion from birefringence of CMB polarization*, *Int. J. Mod. Phys.* **D22** (2013) 1350011 (10 pages).
577. Dass T., *Gauge fields, space-time geometry and gravity*, *Pramana J. Phys.* **23** (1984) 433-443.
578. Datta B.K., *Spinor fields in general relativity I: Noether's theorem and the conservation laws in Riemann-Cartan space*, *Nuovo Cim.* **B6** (1971) 1-15.
579. Datta B.K., *Spinor fields in general relativity II: Generalized field equations and application to the Dirac field*, *Nuovo Cim.* **B6** (1971) 16-25.
580. Datta B.K., DeSabbata V., and Ronchetti L., *Quantization of gravity in real spacetime*, *Nuovo Cim.* **B113** (1998) 711-732.
581. Daum J.-E. and Reuter M., *Renormalization group flow of the Holst action*, *Phys. Lett.* **B710** (2012) 215-218.
582. Daum J.-E. and Reuter M., *Einstein-Cartan gravity, Asymptotic Safety, and the running Immirzi parameter*, *Ann. Phys. (USA)* **334** (2013) 351-419.
583. D'Auria R. and Regge T., *Gravity theories with asymptotically flat instantons*, *Nucl. Phys.* **B195** (1982) 308-324.
584. Daviau C., *On the torsion of the intrinsic spacetime*, *Annales de la Fondation L. de Broglie* **32** (2007) 137-156.
585. Davies J.B., *New curvature-torsion relations through decomposition of Bianchi identities*, *Found. Phys.* **18** (1988) 563-569.
586. Davis W.R., *On the role of projective invariance in general metrical or non-metrical space-time with torsion, spinor and Yang-Mills fields*, *Lett. Nuovo Cim.* **22** (1978) 101-107.

587. Davis W.R., Atkins W.K., and Baker W.M., *On the properties of torsion in Riemann-Cartan space-time. III. Classification of torsion and general discussion*, *Nuovo Cim.* **B44** (1978) 23-38.
588. Davis W.R., Baker W.M., and Green L.H., *Simple conformal-projective invariant gravitational theories with torsion*, in: "On Relativity Theory. Proc. of Sir A.Eddington Centenary Symp., Eds. Y. Choquet-Bruchat and T.M. Karade (World Scientific: Singapore, 1985) vol. **2**, 1-26.
589. De Alfaro V., Fubini S., Furlan G., and Roncadelli M., *Quantum spinning particles in a curved metric*, *Phys. Lett.* **B200** (1988) 323-326.
590. De Almeida F.M.L., Jr., Nepomuceno A.A., and do Vale M.A.B., *Torsion discovery potential and its discrimination at CERN LHC*, *Phys. Rev.* **D79** (2009) 014029 (7 pages).
591. De Andrade V.C. and Pereira J.G., *Riemannian and teleparallel descriptions of the scalar field gravitational interaction*, *Gen. Relat. Grav.* **30** (1998) 263-273.
592. De Andrade V.C. and Pereira J.G., *Gravitational Lorentz force and the description of the gravitational interaction*, *Phys. Rev.* **D56** (1997) 4689-4695.
593. De Andrade V.C. and Pereira J.G., *Torsion and the electromagnetic field*, *Int. J. Mod. Phys.* **D8** (1999) 141-151.
594. De Andrade V.C., Guillen L.C.T., and Pereira J.G., *Teleparallel equivalent of the Kaluza-Klein theory*, *Phys. Rev.* **D61** (2000) 084031 (4 pages).
595. De Andrade V.C., Guillen L.C.T., and Pereira J.G., *Gravitational energy-momentum density in teleparallel gravity*, *Phys. Rev. Lett.* **84** (2000) 4533-4536.
596. De Andrade V.C., Guillen L.C.T., and Pereira J.G., *Teleparallel spin connection*, *Phys. Rev.* **D64** (2001) 027502 (2 pages).
597. De Azeredo C.R. and Oliveira C.G., *Hamiltonian formulation for a translation gauge theory of gravitation*, *Nuovo Cim.* **B74** (1983) 83-92.
598. De Azeredo C.R. and Letelier P., *A quadratic theory of gravitation with Einstein limit and dynamical torsion*, *Progr. Theor. Phys.* **75** (1986) 1359-1364.
599. DeBenedictis A. and Ilijic S., *Spherically symmetric vacuum in covariant  $F(T) = T + \frac{\alpha}{2}T^2 + \mathcal{O}(T^\gamma)$  gravity theory*, *Phys. Rev. D* **94** (2016) 124025 (9 pages).
600. De Berredo-Peixoto G., Helayel-Neto J.A. and Shapiro I.L., *On the consistency of a fermion-torsion effective theory*, *JHEP* **02** (2000) 003 (24 pages).
601. Debney G., Fairchild E.E., Jr., and Siklos S.T.C., *Equivalence of vacuum Yang-Mills gravitation and vacuum Einstein gravitation*, *Gen. Relat. Grav.* **9** (1978) 879-887.
602. Debono I. and Smoot G.F., *General relativity and cosmology: Unsolved questions and future directions*, *Universe* **2**(4) (2016) 23 (82 pages).
603. Deffayet C. and Randjbar-Daemi S., *Nonlinear Fierz-Pauli theory from torsion and bi-gravity*, *Phys. Rev.* **D84** (2011) 044053 (8 pages).
604. Dehnen H. and Hitzler E., *Spin-gauge theory of gravity with Higgs-field mechanism*, *Int. J. Theor. Phys.* **33** (1994) 575-592.
605. Delacrétaz L.V., Endlich S., Monin A., Penco P., and Riva F., *(Re-)inventing the relativistic wheel: gravity, cosets, and spinning objects*, *JHEP* **11** (2014) 008 (31 pages).
606. De la Cruz-Dombriz A., Dunsby P.K.S., and Sáez-Gómez D., *Junction conditions in extended teleparallel gravities*, *JCAP* **12** (2014) 048.
607. Dell J., deLya J.L., and Smolin L., *Quantization of a gauge theory with independent metric and connection fields*, *Phys. Rev.* **D34** (1986) 3012-3024.
608. Demattteis G., *Supersymmetric formulation of gravity theories with nontrivial Lorentz connection*, *Phys. Rev.* **D32** (1985) 3184-3189.

609. Demianski M., De Ritis R., Platania G., Scudellaro P., and Stornaiolo C., *Inflationary models in ECSK theory*, *Phys. Lett.* **A116** (1986) 13-16.
610. Demianski M., De Ritis R., Platania G., Scudellaro P., and Stornaiolo C., *Inflation in a Bianchi type-I Einstein-Cartan cosmological model*, *Phys. Rev.* **D35** (1987) 1181-1184.
611. Demianski M. and Proszynski M., *Static spherical configurations of cold matter in the Einstein-Cartan theory of gravitation*, *Astrophys. and Space Sci.* **53** (1978) 173-179.
612. Demichev A.P. and Nelipa N.F., *Method of construction of gauge invariant Lagrangians for arbitrary Lie groups*, *Progr. Theor. Phys.* **76** (1986) 715-725.
613. Demichev A.P. and Nelipa N.F., *Gauge-invariant Lagrangian for the Poincaré group*, *Moscow University Physics Bulletin*, **27**, n. **3** (1987) 3-7 (in Russian).
614. Denardo G. and Spallucci E., *Curvature and torsion as induced quantum effects*, In: "General relativity and gravitational physics". *Proc. 7th Italian Conf. Rapallo, Italy, September 3-7, 1986*, Eds. U. Bruzzo, R. Cianci, E. Massa (Singapore: World Scientific, 1987) 179-188.
615. Denardo G. and Spallucci E., *Bosonization in a two dimensional Riemann-Cartan geometry*, *Nuovo Cim.* **B98** (1987) 25-36.
616. Denardo G. and Spallucci E., *Curvature and torsion from matter*, *Class. Quantum Grav.* **4** (1987) 89-99.
617. Dent J.B., Dutta S., and Saridakis E.N.,  *$f(T)$  gravity mimicking dynamical dark energy. Background and perturbation analysis*, *JCAP* **01** (2011) 009.
618. Dereli T., Özdemir N., and Sert Ö., *Einstein-Cartan-Dirac theory in (1 + 2)-dimensions*, *Eur. Phys. J.* **C73** (2013) 2279.
619. Dereli T. and Tucker R.W., *Exact neutrino solutions in the presence of torsion*, *Phys. Lett.* **A82** (1981) 229-231.
620. Dereli T. and Tucker R.W., *A gravitational model for a matter-free torsion ball*, *J. Phys.* **A14** (1981) 2957-2967.
621. Dereli T. and Tucker R.W., *Weyl scaling and spinor matter interactions in scalar-tensor theories of gravitation*, *Phys. Lett.* **B110** (1982) 206-210.
622. Dereli T. and Tucker R.W., *Variational methods and effective actions in string models*, *Class. Quantum Grav.* **4** (1987) 791-797.
623. Dereli T. and Tucker R.W., *Gravitational interactions in 2+1 dimensions*, *Class. Quantum Grav.* **5** (1988) 951-959.
624. Dereli T. and Tucker R.W., *Non-metricity induced by dilaton gravity in two dimensions*, *Class. Quantum Grav.* **11** (1994) 2575-2583.
625. Dereli T. and Tucker R.W., *An Einstein-Hilbert action for axi-dilaton gravity in four dimensions*, *Class. Quantum Grav.* **12** (1995) L31-L36.
626. Dereli T. and Tucker R.W., *On the detection of scalar field induced space-time torsion*, *Mod. Phys. Lett.* **A17** (2002) 421.
627. Dereli T. and Tucker R.W., *A broken gauge approach to gravitational mass and charge*, *JHEP* **03** (2002) 041.
628. Dereli T. and Tucker R.W., *On the energy-momentum density of gravitational plane waves*, *Class. Quantum Grav.* **21** (2004) 1459.
629. Dereli T., Önder M., and Tucker R.W., *Solutions for neutral axi-dilaton gravity in four dimensions*, *Class. Quantum Grav.* **12** (1995) L25-L30.
630. Dereli T., Önder M., Schray J., Tucker R.W., and Wang C., *Non-Riemannian gravity and the Einstein - Proca system*, *Class. Quantum Grav.* **13** (1996) L103-L109.

631. Dereli T. and Ücoluk G., *Kaluza-Klein reduction of generalised theories of gravity and non-minimal couplings*, *Class. Quantum Grav.* **7** (1990) 1109-1115.
632. Dereli T. and Verçin, *A gauge model of amorphous solids containing defects*, *Phil. Mag.* **B56** (1987) 625-631.
633. Dereli T. and Verçin, *A gauge model of amorphous solids containing defects II. Chern-Simons free energy*, *Phil. Mag.* **B64** (1991) 509-513.
634. Dereli T. and Yetişmişoğlu C., *New improved massive gravity and three-dimensional space-times of constant curvature and constant torsion*, *Phys. Rev.* **D94** (2016) 064067 (8 pages).
635. De Ritis R., Lavorgna M., Platania G., and Stornaiolo C., *Spin fluid in Einstein-Cartan theory: a variational principle and an extension of the velocity potential representation*, *Phys. Rev.* **D28** (1983) 713-717.
636. De Ritis R., Lavorgna M., Platania G. and Stornaiolo C., *Properties of a spin fluid in a Riemann-Cartan space-time*, *Phys. Lett.* **A95** (1983) 25-28.
637. De Ritis R., Lavorgna M., Platania G., and Stornaiolo C., *Charged spin fluid in the Einstein-Cartan theory*, *Phys. Rev.* **D31** (1985) 1854-1859.
638. De Ritis R., Lavorgna M., and Stornaiolo C., *Geometric optics in a Riemann-Cartan space-time*, *Phys. Lett.* **A98** (1983) 411-413.
639. De Ritis R., Platania G., Scudellaro P., and Stornaiolo C., *Einstein-Cartan theory for a nonminimally coupled scalar field*, In: "General relativity and gravitational physics" *Proc. 8th Italian Conf., Cavalese, Italy, August 30 - September 3, 1988*, Eds. M. Cerdonio, M. Toller, R. Cianci, and M. Francaviglia (Singapore: World Scientific, 1989), 161-170.
640. De Ritis R., Scudellaro P., and Stornaiolo C., *A non-minimal coupled theory of gravitation with torsion*, *Phys. Lett.* **A126** (1988) 389-312.
641. DeSabbata V., *A finite classical model of the electron with torsion and strong gravity*, *Ann. der Phys.* **47** (1990) 511-513.
642. DeSabbata V., *The importance of spin and torsion in the early universe*, *Nuovo Cim.* **A107** (1994) 363-374.
643. DeSabbata V., *Twistors, torsion, and tensor-spinors space-time*, in: "Proc. of the 14th Course of Internat. School of Cosmology and Gravitation: Quantum Gravity" (Erice, 1995), Eds. P.G. Bergmann, V. de Sabbata, and H.-J. Treder (World Scientific: Singapore, 1996) 80-93.
644. DeSabbata V., *Evidence for torsion in gravity?*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?" *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. de Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 51-85.
645. DeSabbata V. and Gasperini M., *Is weak interaction a Cartan force between the elementary particles?*, *Lett. Nuovo Cim.* **21** (1978) 328-332.
646. DeSabbata V. and Gasperini M., *External torsion and the electromagnetic force*, *Lett. Nuovo Cim.* **23** (1978) 657-659.
647. DeSabbata V. and Gasperini M., *Strong gravity with torsion: Some cosmological deduction*, *Lett. Nuovo Cim.* **25** (1979) 489-492.
648. DeSabbata V. and Gasperini M., *Strong gravity with torsion and the Cabbibo angle*, *Gen. Relat. Grav.* **10** (1979) 825-831.
649. DeSabbata V. and Gasperini M., *Strong gravity and weak interactions*, *Gen. Relat. Grav.* **10** (1979) 731-741.
650. DeSabbata V. and Gasperini M., *A cosmological interpretation of the magnetic field of the celestial bodies*, *Lett. Nuovo Cim.* **27** (1980) 133-139.
651. DeSabbata V. and Gasperini M., *On the propagation of electromagnetic waves in a cosmological neutrino sea*, *Lett. Nuovo Cim.* **28** (1980) 181-185.

652. DeSabbata V. and Gasperini M., *Torsion contributions to the magnetic field of a spinning charge*, *Lett. Nuovo Cim.* **28** (1980) 234-236.
653. DeSabbata V. and Gasperini M., *A "semi-minimal" coupling principle for the electromagnetic field in a space with torsion*, *Lett. Nuovo Cim.* **28** (1980) 229-233.
654. DeSabbata V. and Gasperini M., *On Maxwell equations in a Riemann-Cartan space*, *Phys. Lett.* **A77** (1980) 300-302.
655. DeSabbata V. and Gasperini M., *Electromagnetic fields in spaces with torsion*, *Nukleonika* **25** (1980) 1383-1396.
656. DeSabbata V. and Gasperini M., *Torsion and strong gravity in the realm of elementary particles and cosmological physics*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 139-171.
657. DeSabbata V. and Gasperini M., *Gauge invariance, semi-minimal coupling, and propagating torsion*, *Phys. Rev.* **D23** (1981) 2116-2120.
658. DeSabbata V. and Gasperini M., *Propagating torsion and electromagnetic gauge invariance*, *Phys. Lett.* **A83** (1981) 115-117.
659. DeSabbata V. and Gasperini M., *A gauge invariant interaction between charged spinor field and a propagating torsion*, *Lett. Nuovo Cim.* **30** (1981) 1933-198.
660. DeSabbata V. and Gasperini M., *Torsion production by electromagnetic fields*, *Lett. Nuovo Cim.* **30** (1981) 363-366.
661. DeSabbata V. and Gasperini M., *Macroscopical consequences of a propagating torsion potential*, *Lett. Nuovo Cim.* **30** (1981) 503-506.
662. DeSabbata V. and Gasperini M., *A theory of gravity with two metric and two torsion tensor*, *Lett. Nuovo Cim.* **32** (1981) 69-74.
663. DeSabbata V. and Gasperini M., *Neutrino oscillations in the presence of torsion*, *Nuovo Cim.* **A65** (1981) 479-500.
664. DeSabbata V. and Gasperini M., *Neutron spin rotation induced by torsion in vacuum*, *Lett. Nuovo Cim.* **33** (1982) 363-366.
665. DeSabbata V. and Gasperini M., *The anthropic principle and the actual size of the universe*, *Lett. Nuovo Cim.* **37** (1983) 605-608.
666. DeSabbata V. and Gasperini M., *Introduction to gravitation* (World Scientific: Singapore, 1985) 346 pp.
667. DeSabbata V., Melnikov V.N., and Pronin P.I., *Theoretical treatment of non-Newtonian forces*, *Progr. Theor. Phys.* **88** (1992) 623-661.
668. DeSabbata V., Pronin P.I., and Sivaram C., *Neutron interferometry in gravitational field with torsion*, *Int. J. Theor. Phys.* **30** (1991) 1671-1678.
669. DeSabbata V. and Ronchetti L., *A Hamiltonian formulation of gravitational theory that allows one to consider curvature and torsion as conjugate variables*, *Found. Phys.* **29** (1999) 1099-1117.
670. DeSabbata V. and Sivaram C., *The need for an energy-dependent torsion-coupling constant in the early Universe*, *Astrophys. and Space Sci.* **158** (1989) 347-352.
671. DeSabbata V. and Sivaram C., *Strong spin torsion interaction between spinning protons*, *Nuovo Cim.* **A101** (1989) 273-283.
672. DeSabbata V. and Sivaram C., *Torsion effects on neutrino emission from dense collapsing cores*, *Ann. der Phys.* **46** (1989) 627-630.
673. DeSabbata V. and Sivaram C., *A finite classical model of the electron with torsion and strong gravity*, *Ann. der Phys.* **47** (1990) 511-513.

674. DeSabbata V. and Sivaram C., *Fifth force as a manifestation of torsion*, *Int. J. Theor. Phys.* **29** (1990) 1-6.
675. DeSabbata V. and Sivaram C., *Torsion and the cosmological constant problem*, *Astropys. and Space Sci.* **165** (1990) 51-55.
676. DeSabbata V. and Sivaram C., *Torsion and inflation*, *Astropys. and Space Sci.* **176** (1991) 141-144.
677. DeSabbata V. and Sivaram C., *Torsion, wormholes and the problem of the cosmological constant*, *Int. J. Theor. Phys.* **30** (1991) 123-127.
678. DeSabbata V. and Sivaram C., *Gravimagnetic field, torsion and gravitational shielding*, *Nuovo Cim.* **B106** (1991) 883-888.
679. DeSabbata V. and Sivaram C., *Torsion and quantum effects*, in: *Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko*, Eds.P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 143-174.
680. DeSabbata V. and Sivaram C., *Torsion and magnetism*, in: *Perspectives of the unified theory*, Eds. D.V.Gal'tsov, L.S.Kuz'menkov and P.I.Pronin (Moscow Univ. Publ. House: Moscow, 1991) 98-115 (in Russian).
681. DeSabbata V. and Sivaram C., *Torsion as the basis for string tension*, *Ann.der Phys.* **48** (1991) 1195-1203.
682. DeSabbata V. and Sivaram C., *The central role of spin in black hole evaporation*, in: "Black hole physics", *Proc. of the NATO Adv. Study Inst. (12th Course of the Int. School of Cosmology and Gravitation of the E. Majorana Centre, Erice, Italy, 12-22 May 1991)*, Eds. V. DeSabbata and Z. Zhang (Kluwer Academic Publ.: Dordrecht, 1992) 225-282.
683. DeSabbata V. and Sivaram C., *Anyons and torsion*, *Nuovo Cim.* **B107** (1992) 1201-1205.
684. DeSabbata V. and Sivaram C., *Torsion, minimum time, string tension and its physical implications in cosmology*, *Astropys. and Space Sci.* **187** (1992) 149-154.
685. DeSabbata V. and Sivaram C., *A minimal time and time-temperature uncertainty principle*, *Found. Phys. Lett.* **5** (1992) 183-189.
686. DeSabbata V. and Sivaram C., *Curvature from spin*, *Found. Phys. Lett.* **5** (1992) 579-584.
687. DeSabbata V. and Sivaram C., *Universality of charge arising from torsion*, *Found. Phys. Lett.* **6** (1993) 201-206.
688. DeSabbata V. and Sivaram C., *Gravity and CP violation*, *Found. Phys. Lett.* **7** (1994) 475-482.
689. DeSabbata V. and Sivaram C., *Spin and torsion in gravitation* (World Scientific: Singapore, 1994) 313 pp.
690. DeSabbata V. and Sivaram C., *Torsion, string tension, and topological origin of charge and mass*, *Found. Phys. Lett.* **8** (1995) 375-380.
691. DeSabbata V. and Sivaram C., *Twistors, torsion and quantization of space-time*, *Nuovo Cim.* **A109** (1996) 377-390.
692. DeSabbata V., Sivaram C., Borzeszkowski H.-H. von, and Treder H.-J. *Quantum general relativity, torsion and uncertainty relations*, *Ann. der Phys.* **48** (1991) 497-502.
693. DeSabbata V., Sivaram C., and Garcia de Andrade L., *Torsion and gauge-invariant massive electrodynamics*, *Int. J. Theor. Phys.* **32** (1993) 1523-1530.
694. DeSabbata V., Dingxiong Wang, and Sivaram C., *Torsion effects in black hole evaporation*, *Ann. d. Phys.* **47** (1990) 508-510.
695. DeSabbata V. and Xin Y., , *Twistors and Einstein-Cartan theory*, *Int. J. Mod. Phys.* **A10** (1995) 3663-3674.

696. Desai S. and Poplawski N., *Non-parametric reconstruction of an inflaton potential from Einstein-Cartan-Sciama-Kibble gravity with particle production*, *Phys. Lett.* **B755** (2016) 183-189.
697. De Sousa G.P., *On spin and (quantum) gravity in (2+1) dimensions*, *Nucl. Phys.* **B346** (1990) 440-472.
698. Destri C. and Connella G., *One-loop spontaneous compactification with torsion and spin*, *Nuovo Cim.* **A97** (1987) 343-358.
699. Diakonov D., Tumanov A.G., and Vladimirov A.A., *Low-energy general relativity with torsion: A systematic derivative expansion*, *Phys. Rev.* **D84** (2011) 124042 (16 pages).
700. Dimakis A., *The initial value problem of the Poincaré gauge theory in vacuum. I. Second order formalism*, *Ann. Inst. H.Poincaré* **A51** (1989) 371-388.
701. Dimakis A., *The initial value problem of the Poincaré gauge theory in vacuum. II. First order formalism*, *Ann. Inst. H.Poincaré* **A51** (1989) 389-417.
702. Dimakis A. and Müller-Hoisen F., *Massive ghost Dirac fields in Einstein-Cartan theory*, *Phys. Lett.* **A92** (1982) 431-432.
703. Dimakis A. and Müller-Hoisen F., *Solutions of the Einstein-Cartan-Dirac equations with vanishing energy-momentum tensor*, *J. Math. Phys.* **26** (1985) 1040-1048.
704. Dineykhon M., *Spinors in quantum space with torsion*, *Int. J. Theor. Phys.* **26** (1987) 715-723.
705. Dineykhon M. and Namsrai Kh., *Quantum space-time, torsion and magnetic monopoles*, *Int. J. Theor. Phys.* **24** (1985) 1197-1215.
706. Ding Shu-Xue, *Topology changing solutions of (2+1)-dimensional Einstein-Cartan theory*, *Class. Quantum Grav.* **10** (1993) 851-858.
707. Ding Shu-Xue, Song He-Shan, and An Ing *Anyon solution of an abelian gauge field coupling with torsion*, *Commun. Theor. Phys.* **19** (1993) 357-360.
708. DiStefano R., and Rauch R.T., *New approach the first-order canonical formulation of gravitation. Application to Einstein-Cartan-Sciama-Kibble theory*, *Phys. Rev.* **D26** (1982) 1242-1253.
709. Dobado A. and Maroto A.L., *Standard model anomalies in curved space-time with torsion*, *Phys. Rev.* **D54** (1996) 5185-5194.
710. Dobado A. and Maroto A.L., *Primordial torsion fields as an explanation of the anisotropy in cosmological electromagnetic propagation*, *Mod. Phys. Lett.* **A12** (1997) 3003-3007.
711. Dobado A. and Maroto A.L., *Non-local low-energy effective action for gravity with torsion*, *Class. Quantum Grav.* **16** (1999) 4057-4074.
712. Doccatelli D., Agostini W., and Festa P., *The motion of spinning particles. Post-Newtonian approximation in the Einstein-Cartan theory*, *Nuovo Cim.* **B49** (1979) 45-54.
713. Dolan B.P., *On elimination of Pauli couplings in Kaluza-Klein theories with torsion*, *Phys. Lett.* **B159** (1985) 279-283.
714. Dolan B.P., *Chiral fermions and torsion in the early Universe*, *Class. Quantum Grav.* **27** (2010) 095010.
715. Dong H., Wang Y., and Meng X., *Extended Birkhoff's theorem in  $f(T)$  gravity*, *Eur. Phys. J.* **C72** (2012) 2002.
716. Dong H., Wang Y., and Meng X., *Birkhoff's theorem in  $f(T)$  gravity up to the perturbative order*, *Eur. Phys. J.* **C72** (2012) 2201.
717. Donnelly H., *Heat equation asymptotics with torsion*, *Indiana Univ. Math. J.* **34** (1985) 105-113.

718. Doran C., Lasenby A., Challinor A., and Gull S., *Effects of spin-torsion in gauge theory of gravity*, *J. Math. Phys.* **39** (1998) 3303-3321.
719. Dorofeev V.Yu., *Riemann-Cartan space in O-theory*, *Grav. Cosmol.* **18** (2012) 100-103.
720. Downes R.J., Levitin M., and Vassiliev D., *Spectral asymmetry of the massless Dirac operator on a 3-torus*, *J. Math. Phys.* **54** (2013) 111503.
721. Drechsler W., *Wave equations on a de Sitter fiber bundle*, *Fortschr. Phys.* **23** (1975) 607-647.
722. Drechsler W., *Group contraction in a fiber bundle with Cartan connection*, *J. Math. Phys.* **18** (1977) 1358-1366.
723. Drechsler W., *Lagrange formulation for a gauge theory of strong and electromagnetic interactions defined on a Cartan bundle*, *Nuovo Cim.* **A41** (1977) 597-620.
724. Drechsler W., *Field and particle equations for a gauge theory based on the (4,1) de Sitter group*, *Nuovo Cim.* **A50** (1979) 405-422.
725. Drechsler W., *Poincaré gauge field theory and gravitation*, *Ann. Inst. H.Poincaré* **A37** (1982) 155-184.
726. Drechsler W., *Topological invariants and the dynamics of an axial vector torsion field*, *Gen. Relat. Grav.* **15** (1983) 703-723.
727. Drechsler W., *Poincaré gauge theory, gravitation, and transformation of matter fields*, *Fortschr. Phys.* **32** (1984) 449-472.
728. Drechsler W., *Linearly and nonlinearly transforming fields on homogeneous spaces of the (4,1) de Sitter group*, *J. Math. Phys.* **26** (1985) 41-54.
729. Drechsler W., *Generalized spinor fields and gravitation*, *Class. Quantum Grav.* **4** (1987) 291-318.
730. Drechsler W., *U(4) spin gauge theory over a Riemann-Cartan space-time*, *Z. Phys.* **C41** (1988) 197-205.
731. Drechsler W., *Modified Weyl theory and extended elementary objects*, *Found. Phys.* **19** (1989) 1479-1497.
732. Drechsler W., *Gauge theory for extended elementary objects*, *Class. Quantum Grav.* **6** (1989) 623-657.
733. Drechsler W., *Geometric formulation of gauge theories*, *Z. Naturforsch.* **A46** (1991) 645-654.
734. Drechsler W., *Quantized fiber dynamics for extended elementary objects involving gravitation*, *Found. Phys.* **22** (1992) 1041-1077.
735. Drechsler W., *Classical versus quantum gravity*, *Found. Phys.* **23** (1993) 261-276.
736. Drechsler W., *Quantized matter in a de Sitter gauge theory with classical metric and axial torsion*, in: "Proc. of the 14th Course of Internat. School of Cosmology and Gravitation: Quantum Gravity" (Erice, 1995), Eds. P.G. Bergmann, V. DeSabbata, and H.-J. Treder (World Scientific: Singapore, 1996) 94-118.
737. Drechsler W., *Geometro-stochastically quantized fields with internal spin variables*, *J. Math. Phys.* **38** (1997) 5531-5558.
738. Drechsler W. and Mayer M.E., *Fiber bundle techniques in gauge theories*, *Lect. Notes Phys.* **67** (Springer: Heidelberg, 1977) 102 p.
739. Drechsler W. and Prugovecki E., *Geometro-stochastic quantization of a theory for extended elementary objects*, *Found. Phys.* **21** (1991) 513-546.
740. Drechsler W. and Thacker W., *Generalised spinor fields and gravitation*, *Class. Quantum Grav.* **4** (1987) 291-318.

741. Drummond I.T., *Regge-Palatini calculus*, Nucl. Phys. **B273** (1986) 125-136.
742. Duan Yishi, *Gauge theories of gravitation*, Commun. Theor. Phys. **4** (1985) 661-674.
743. Duan Y.S. and Jiang Y., *Can torsion play a role in angular momentum conservation law?*, Gen. Rel. Grav. **31** (1999) 3-12.
744. Duan Y.S. and Jiang Y., *Strings and the gauge theory of spacetime defects*, Int. J. Theor. Phys. **38** (1999) 563-573.
745. Duan Y.S. and Li S., *The topological structure of the space-time disclination*, J. Math. Phys. **39** (1998) 6696-6705.
746. Duan Y.S., Yang G.H., and Jiang Y., *The origin and bifurcation of the space-time defects in the early universe*, Gen. Rel. Grav. **29** (1997) 715-725.
747. Duan Y.S., Yang G.H., and Jiang Y., *The quantization of the space-time defects in the early universe*, Helv. Phys. Acta **70** (1997) 565-577.
748. Duan Y.S., Yang G.H., and Jiang Y., *The quantization and production of the space-time defects in the early universe*, Int. J. Mod. Phys. **A13** (1998) 2001-2012.
749. Duan Y.S., Zhang S.L., and Feng S.S., *Gauge potential decomposition, space-time defects, and Planck's constant*, J. Math. Phys. **35** (1994) 4463-4468.
750. Dürr H.P., *Poincaré gauge invariant spinor theory and the gravitational field I*, Nuovo Cim. **A4** (1971) 187-197.
751. Dürr H.P., *Poincaré gauge invariant spinor theory and the gravitational field II*, Gen. Relat. Grav. **4** (1973) 29-52.
752. Duncan M.J., Kaloper N., and Olive K.A., *Axion hair and dynamical torsion from anomalies*, Nucl. Phys. **B387** (1992) 215-235.
753. Duplij S., *Towards gauge principle for semigroups*, Acta Phys. Pol. **B23** (1992) 733-743.
754. Dzhunushaliev V.D., *Spherically symmetric solution for torsion and the Dirac equation in 5D spacetime*, Int. J. Mod. Phys. **D7** (1998) 909-915.
755. Dzhunushaliev V.D., Folomeev V., Kleihaus B., and Kunz J., *Quantum torsion with non-zero standard deviation: Non-perturbative approach for cosmology*, Phys. Lett. **B719** (2013) 5-8.
756. Dzhunushaliev V. and Singleton D., *Einstein-Cartan-Heisenberg theory of gravity with dynamical torsion*, Phys. Lett. **A257** (1999) 7-13.

## E

757. Eddington A.S., *A generalization of Weyl's theory of the electromagnetic and gravitational field*, Proc. Roy. Soc. **A99** (1921) 104-122.
758. Eddington A.S., *The mathematical theory of relativity* (Cambridge Univ. Press: Cambridge, 1924).
759. Edelen D.G.B., *The structure of field space* (University of California Press: Berkeley, 1962).
760. Edelen D.G.B., *Operator-valued connections, Lie-connections and gauge field theory*, Int. J. Theor. Phys. **23** (1984) 949-985.
761. Edelen D.G.B., *Direct gauging of the Poincaré group*, Int. J. Theor. Phys. **24** (1985) 659-673.
762. Edelen D.G.B., *Direct gauging of the Poincaré group II*, Int. J. Theor. Phys. **24** (1985) 1091-1111.

763. Edelen D.G.B., *Direct gauging of the Poincaré group III. Interaction with internal symmetries*, *Int. J. Theor. Phys.* **24** (1985) 1133-1141.
764. Edelen D.G.B., *Direct gauging of the Poincaré group IV. Curvature, holonomy, spin, and gravity*, *Int. J. Theor. Phys.* **24** (1985) 1173-1190.
765. Edelen D.G.B., *Direct gauging of the Poincaré group V. Group scaling, classical gauge theory, and gravitational corrections*, *Int. J. Theor. Phys.* **25** (1986) 671-684.
766. Edelen D.G.B., *Static vacuum solution of direct Poincaré gauge theory in ten-dimensions with four external*, *Int. J. Theor. Phys.* **27** (1988) 641-648.
767. Edelen D., *Gravity and gauge. A new perspective*, *Int. J. Theor. Phys.* **28** (1989) 633-649.
768. Edgar S.B., *Necessary and sufficient conditions for (I) Weyl, (II) Riemann-Cartan connections*, *J. Geom. Phys.* **13** (1994) 259-274.
769. Einstein A., *The meaning of relativity* (Princeton Univ. Press: Princeton, 1955).
770. Eisenhart L.P., *Non-Riemannian geometry* (American Math. Soc.: Providence RI, 1990) ninth printing.
771. Eguchi T., Gilkey P.B., and Hanson A.J., *Gravitation, gauge theories and differential geometry*, *Phys. Rep.* **66** (1980) 213-393.
772. El Hanafy W. and Nashed G.G.L., *Exact teleparallel gravity of binary black holes*, *Astrophys. Space Sci.* **361** (2016) 68 (11 pages).
773. Elizalde E. and Odintsov S.D., *Higher derivative quantum gravity with torsion in the conformally selfdual limit*, *Int. J. Mod. Phys.* **D2** (1993) 51-58.
774. Elizalde E. and Odintsov S.D., *Effective potential for the conformal sector of quantum gravity with torsion*, *Phys. Lett.* **B315** (1993) 245-250.
775. Elizalde E. and Odintsov S.D.,  *$R^2$  gravity with torsion in the conformally selfdual limit*, *Yad. Fiz.* **56** (1993) 215-219 (in Russian); English translation: *Phys. Atomic Nucl.* **56** (1993) 409-411.
776. Elyasi N. and Boroojerdian N., *Affine metrics: an structure for unification of gravitation and electromagnetism*, *Int. J. Theor. Phys.* **50** (2011) 850-860.
777. Epstein M., Burton D.A., and Tucker R., *Relativistic anelasticity*, *Class. Quantum Grav.* **23** (2006) 3545.
778. Epstein M. and de Leon M., *Geometrical theory of uniform Cosserat media*, *J. Geom. Phys.* **26** (1998) 127-170.
779. Ertl M.F., Katanaev M.O., and Kummer W., *Generalized supergravity in two dimensions*, *Nucl. Phys.* **B530** (1998) 457-486.
780. Espiro J.L. and Vásquez, Y., *Some cosmological models coming from gravitational theories having torsional degrees of freedom*, *Gen. Relat. Grav.* **48** (2016) 117 (27 pages).
781. Esposito G., *Hamiltonian structure of Friedman-Robertson-Walker Universe with torsion*, *Nuovo Cim.* **B104** (1989) 199-212.
782. Esposito G., *The singularity problem for the space-time with torsion*, *Nuovo Cim.* **B105** (1990) 75-90.
783. Esposito G., *Mathematical structures of space-time*, *Fortschr. Phys.* **40** (1992) 1-30.
784. Esposito G., *Quantum gravity, quantum cosmology and Lorentzian geometries* (Berlin et al: Springer, 1992), *Lect. Notes Phys.* **m 12**, 326 p.
785. Esposito G.,  *$\alpha$ -Surfaces for complex space-times with torsion*, *Nuovo Cim.*, **B108** (1993) 123-125.
786. Esposito G., *Complex general relativity* (Amsterdam: Kluwer Academic, 1995), *Fundamental Theories of Physics* **v 69**.
787. Esposito G., Gionti G., Marmo G., and Stornaiolo C., *Lagrangian theory of constraint systems: cosmological application*, *Nuovo Cim.* **B109** (1994) 1259-1273.

*F*

788. Fabbri L., *On a completely antisymmetric Cartan torsion tensor*, *Annales de la Fondation L. de Broglie* **32** (2007) 215-228.
789. Fabbri L., *On the problem of unicity in Einstein-Sciama-Kibble theory*, *Annales de la Fondation L. de Broglie* **33** (2008) 365-378.
790. Fabbri L., *Leptonic electroweak spin-torsion interactions*, *Mod. Phys. Lett.* **A26** (2011) 1697.
791. Fabbri L., *The spin-torsion coupling and causality for the Standard Model*, *Mod. Phys. Lett.* **A26** (2011) 2091.
792. Fabbri L., *From the torsion tensor for spinors to the weak forces for leptons*, *Int. J. Theor. Phys.* **50** (2011) 3616-3620.
793. Fabbri L., *Metric-torsional conformal gravity*, *Phys. Lett.* **B707** (2012) 415-417.
794. Fabbri L., *On the consistency of constraints in matter field theories*, *Int. J. Theor. Phys.* **51** (2012) 954-963.
795. Fabbri L., *Massless fermion mixing for semispinorial torsional interaction*, *Annales de la Fondation L. de Broglie* **37** (2012) 33-39.
796. Fabbri L., *Hadronic electroweak spin-torsion interactions*, *Annales de la Fondation L. de Broglie* **37** (2012) 83-90.
797. Fabbri L., *On geometrically unified fields and universal constants*, *Gen. Rel. Grav.* **45** (2013) 1285-1296.
798. Fabbri L., *A discussion on Dirac field theory, no-go theorems and renormalizability*, *Int. J. Theor. Phys.* **52** (2013) 634-643.
799. Fabbri L., *Metric solutions in torsionless gauge for vacuum conformal gravity*, *J. Math. Phys.* **54** (2013) 062501.
800. Fabbri L., *Conformal gravity with Dirac matter*, *Annales de la Fondation L. de Broglie* **38** (2013) 155-165.
801. Fabbri L., *Least-order torsion-gravity for chiral-spinor fields, induced self-interacting potentials and parity conservation*, *Gen. Rel. Grav.* **46** (2014) 1663.
802. Fabbri L., *On a purely geometric approach to the Dirac matter field and its quantum properties*, *Int. J. Theor. Phys.* **53** (2014) 1896-1909.
803. Fabbri L., *On the geometry of the Dirac matter with the fermionic potentials and its quantum properties*, *Int. J. Theor. Phys.* **53** (2014) 3744-3755.
804. Fabbri L., *Torsion-gravity for Dirac fields and their effective phenomenology*, *Mod. Phys. Lett.* **A29** (2014) 1450133 (12 pages).
805. Fabbri L., *Least-order torsion-gravity for Dirac fields, and their non-linearity terms*, *Gen. Rel. Grav.* **47** (2015) 1837.
806. Fabbri L., *Torsionally-gravitating charged matter fields and quanta*, *Gen. Rel. Gravit.* **47** (2015) 119.
807. Fabbri L., *Torsion gravity for Dirac fields*, *Int. J. Geom. Methods Mod. Phys.* **14** (2017) 1750037 [17 pages].
808. Fabbri L. and Mannheim P.D., *Continuity of the torsionless limit as a selection rule for gravity theories with torsion*, *Phys. Rev.* **D90** (2014) 024042 (6 pages).
809. Fabbri L. and Vignolo S., *Dirac fields in  $f(R)$ -gravity with torsion*, *Class. Quantum Grav.* **28** (2011) 125002.

810. Fabbri L. and Vignolo S., *A modified theory of gravity with torsion and its applications to cosmology and particle physics*, *Int. J. Theor. Phys.* **51** (2012) 3186-3207.
811. Fabbri L. and Vignolo S., *ELKO and Dirac spinors seen from torsion*, *Int. J. Mod. Phys. D* **23** (2014) 1444001 (19 pages).
812. Fabbri L., Vignolo S., and Carloni S., *Renormalizability of the Dirac equation in torsion gravity with nonminimal coupling*, *Phys. Rev.* **D90** (2014) 024012 (16 pages).
813. Fairchild E.E., Jr., *Gauge theory of gravitation*, *Phys. Rev.* **D14** (1976) 384-391; (E) 3439.
814. Fairchild E.E., Jr., *Yang-Mills formulation of gravitational dynamics*, *Phys. Rev.* **D16** (1977) 2438-2447.
815. Fariborz A.H. and McKeon D.G.C., *Spinors in Weyl geometry*, *Class. Quantum Grav.* **14** (1997) 2517-2525.
816. Farfán F., Garca-Salcedo R., Loaiza-Brito O., Moreno C., and Yakhno A., *Spherically symmetric solution in a space-time with torsion*, *Gen. Rel. Grav.* **44** (2012) 535-553.
817. Farrugia D. and Levi Said J., *Stability of the flat FLRW metric in  $f(T)$  gravity*, *Phys. Rev.* **D94** (2016) 124054 (13 pages).
818. Fatibene L. and Garruto S., *Extended theories of gravitation*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650070 (5 pages).
819. Fazlipoor, B., *Interacting quintessence in a new scalar-torsion gravity*, *Gen. Relativ. Gravit.* **48** (2016) 159 (19 pages).
820. Fedorov F.I., Kudin V.I., and Minkevich A.V., *Nonsingular cosmology and the gravitational Lagrangians in the gauge theory of gravity*, *Acta Phys. Pol.* **B15** (1984) 107-115.
821. Fedorov F.I., Minkevich A.V., and Chuong Nguen Hong, *Spontaneous symmetry breaking by gravitational field and regular isotropic models with torsion*, *Class. Quantum Grav.* **5** (1988) 515-519.
822. Feng C.-J., Ge F.-F., Li X.-Z., Lin R.-H., and Zhai X.-H., *Towards realistic  $f(T)$  models with nonminimal torsion-matter coupling extension*, *Phys. Rev.* **D92** (2015) 104038 (12 pages).
823. Fennelly A.J., Krisch J.P., Ray J.R., and Smalley L.L., *Including spin in the Raychaudhuri equation*, *J. Math. Phys.* **32** (1991) 485-487.
824. Fennelly A.J. and Pavelle R., *Homogeneous cosmological models in Yang's gravitational theory*, *J. Phys.* **A12** (1979) 227-228.
825. Fennelly A.J., Bradas J.C., and Smalley L.L., *Inflation in the Einstein-Cartan theory with energy-momentum tensor with spin*, *Phys. Lett.* **A129** (1988) 195-200.
826. Fernández V.A. and Rodrigues W.A. Jr., *Gravitation as a plastic distortion of the Lorentz vacuum* (Springer: Berlin, 2010).
827. Ferrández A., Guerrero J., Javaloyes M.A., and Lucas P., *Particles with curvature and torsion in three-dimensional pseudo-Riemannian space forms*, *J. Geom. Phys.* **56** (2006) 1666-1687.
828. Ferrari A.F., Neto J.A.S., and da Rocha R., *The role of singular spinor fields in a torsional gravity, Lorentz-violating, framework*, *Gen. Relat. Grav.* **49** (2017) 70.
829. Ferraris M. and Kijowski J., *Unified geometric theory of electromagnetic and gravitational interactions*, *Gen. Relat. Grav.* **14** (1982) 37-47.
830. Ferraris M., Francaviglia M., Palese M., and Winterroth E., *Gauge-natural noether currents and connection fields*, *Int. J. Geom. Methods Mod. Phys.* **08** (2011) 177-186.
831. Ferraris M., Francaviglia M., and Raiteri M., *Conserved quantities from the equations of motion: with applications to natural and gauge natural theories of gravitation*, *Class. Quantum Grav.* **20** (2003) 4043.

832. Ferraris M., Francaviglia M., and Volovich I., *A model of affine gravity in two dimensions and plurality of topology*, *Int. J. Mod. Phys.* **A12** (1997) 5067-5080.
833. Ferraro R. and Fiorini F., *Modified teleparallel gravity: Inflation without an inflaton*, *Phys. Rev.* **D75** (2007) 084031 (5 pages).
834. Ferraro R. and Fiorini F., *Born-Infeld gravity in Weitzenböck spacetime*, *Phys. Rev.* **D78** (2008) 124019 (8 pages).
835. Ferraro R. and Fiorini F., *Spherically symmetric static spacetimes in vacuum  $f(T)$  gravity*, *Phys. Rev.* **D84** (2011) 083518 (8 pages).
836. Ferraro R. and Fiorini F., *Non-trivial frames for  $f(T)$  theories of gravity and beyond*, *Phys. Lett.* **B702** (2011) 75-80.
837. Ferraro R. and Guzmán M.J., *Hamiltonian formulation of teleparallel gravity*, *Phys. Rev.* **D94** (2016) 104045 (14 pages).
838. Ferreira A.C., *Einstein four-manifolds with skew torsion*, *J. Geom. Phys.* **61** (2011) 2341-2351.
839. Ferreira C.N., Godinho C.F.L., and Helayël Neto J.A., *A Landau-Lifshitz-Gilbert-type equation and torsion effects on the dynamics of magnetization*, *Ann. Phys. (Berlin)* **529** (2017) 1600186 (11 pages).
840. Festuccia G., Hansen D., Hartong J., and Obers N.A., *Torsional Newton-Cartan geometry from the Noether procedure*, *Phys. Rev.* **D94** (2016) 105023 (16 pages).
841. Figueredo B.D.B., *Gödel-type spacetimes and the motions of charged particles*, *Class. Quantum Grav.* **15** (1998) 3849-3862.
842. Figueredo B.D.B., Damiao S.I., and Tiomno J., *On the detection of matter vorticity and space-time torsion*, *Phys. Lett.* **A137** (1989) 99-102.
843. Figueredo B.D.B., Damiao S.I., and Tiomno J., *Gravitational coupling of Klein-Gordon and Dirac particles to matter vorticity and spacetime torsion*, *Class. Quantum Grav.* **9** (1992) 1593-1617.
844. Finkelstein D. and Gibbs J.M., *Quantum relativity*, *Int. J. Theor. Phys.* **32** (1993) 1801-1813.
845. Finkelstein R., *Spacetime of the elementary particles*, *J. Math. Phys.* **1** (1960) 440-451.
846. Finkelstein R., *Spinor fields in spaces with torsion*, *Ann. Phys. (USA)* **12** (1961) 200-221.
847. Finkelstein R., *Elementary interactions in spaces with torsion*, *Ann. Phys. (USA)* **15** (1961) 223-249.
848. Finkelstein R. and Ramsay W., *The strong couplings in a space with torsion*, *Ann. Phys. (USA)* **17** (1962) 379-403.
849. Finkelstein R. and Ramsay W., *The weak and the strong couplings and general covariance*, *Ann. Phys. (USA)* **21** (1963) 408-433.
850. Fiziev P., *Spinless matter in transposed-equi-affine theory of gravity*, *Gen. Rel. Grav.* **30** (1998) 1341-1370.
851. Fiziev P. and Kleinert H., *New action principle for classical particle trajectories in spaces with torsion*, *Europhys. Lett.* **35** (1996) 241-246.
852. Fiziev P. and Kleinert H., *Comment on path integral derivation of Schrödinger equation in spaces with curvature and torsion*, *J. Phys. A: Math. Gen.* **29** (1996) 7619-7624.
853. Fiziev P. and Yazadjiev S., *The transposed-equi-affine theory of gravity and solar-system experiments*, *Class. Quantum Grav.* **16** (1999) 3133-3136.
854. Flaherty F.J., and Kerlick G.D., *Inhomogeneous local gauge transformations in spacetimes with torsion*, *J. Math. Phys.* **20** (1979) 1281-1284.

855. Flanagan É.É., and Rosenthal E., *Can Gravity Probe B usefully constrain torsion gravity theories?* *Phys. Rev.* **D75** (2007) 124016 (13 pages).
856. Floreanini R. and Percacci R., *Canonical algebra of  $GL(4)$ -invariant gravity*, *Class. Quantum Grav.* **7** (1990) 975-984.
857. Floreanini R. and Percacci R., *Palatini formalism and new canonical variables for  $GL(4)$ -invariant gravity*, *Class. Quantum Grav.* **7** (1990) 1805-1818.
858. Floreanini R. and Percacci R., *Mean-field quantum gravity*, *Phys. Rev.* **D46** (1992) 1566-1579.
859. Floreanini R. and Percacci R., *Average effective potential for the conformal factor*, *Nucl. Phys.* **B436** (1995) 141-160.
860. Floreanini R., Percacci R., and Spallucci E., *Coleman-Weinberg effect in quantum gravity*, *Class. Quantum Grav.* **8** (1991) L193-L197.
861. Fock V., *Sur les equations de Dirac dans la théorie de relativité générale*, *C. R. Acad. Sci. (Paris) Phys. Théor.* **189** (1929) 25.
862. Fock V., *Geometrisierung der Diracschen Theorie des Elektrons*, *Zs. f. Phys.* **57** (1929) 261-277.
863. Fock V. and Ivanenko D., *Géométrie quantique linéaire et déplacement parallèle*, *C. R. Acad. Sci. (Paris) Phys. Théor.* **188** (1929) 1470-1473.
864. Fock V. and Ivanenko D., *Über eine mögliche geometrische Deutung der relativistischen Quantentheorie*, *Zs. f. Phys.* **54** (1929) 798-802.
865. Folland G.B., *Weyl manifolds*, *J. Differ. Geom.* **4** (1970) 145-153.
866. Fonseca-Neto J.B., and Riveles V.O., *A geometric formulation of Einstein-Maxwell theory in Einstein-Cartan space-time*, *Rev. Bras. Fis.* **16** (1986) 521.
867. Fonseca-Neto J.B. and Rebouças M.J., *On a class of Riemann-Cartan space-times of Gödel type*, *Gen. Rel. Grav.* **30** (1998) 1301-1317.
868. Fonseca-Neto J.B., Rebouças M.J., and MacCallum, M. A. H., *Algebraic computing in torsion theories of gravitation*, *Math. Comput. Simulation* **42** (1996) 739-748.
869. Fonseca-Neto J.B., Rebouças M.J., and Teixeira A.F.F., *The equivalence problem in torsion theories of gravitation*, *J. Math. Phys.* **33** (1992) 2574-2577.
870. Fonseca-Neto J.B., Romero C., Martinez S.P.G., *Scalar torsion and a new symmetry of general relativity*, *Gen. Rel. Grav.* **45** (2013) 1579-1601.
871. Formiga J.B., *Comment on "Metric-affine approach to teleparallel gravity"*, *Phys. Rev.* **D88** (2013) 068501 (3 pages).
872. Formiga J.B., *Equivalence between an extension of teleparallelism to a Weyl geometry and general relativity*, *Int. J. Theor. Phys.* **53** (2014) 1971-1977.
873. Foster J., Kostelecký V.A., and Xu R., *Constraints on nonmetricity from bounds on Lorentz violation*, *Phys. Rev.* **D95** (2017) 084033 (6 pages).
874. Fournel C., François J., Lazzarini S., and Masson T., *Gauge invariant composite fields out of connections, with examples*, *Int. J. Geom. Methods Mod. Phys.* **11** (2014) 1450016 (28 pages).
875. Fradkin E.S. and Shvartsman Sh.M., *Effective action of a relativistic spinning particle in a gravitational field with torsion*, *Class. Quantum Grav.* **9** (1992) 17-26.
876. Francaviglia M. and Kijowski J., *Integrability conditions for a gravitational theory with nonmetric connection*, *Gen. Relat. Grav.* **12** (1980) 279-286.

877. Frąckiewicz K., *Gauge group, degeneration, and degrees of freedom in the ECSK theory of gravity. I. The symplectic structure of the ECSK theory*, *Gen. Relat. Grav.* **17** (1985) 939-952.
878. Frąckiewicz K., *Gauge group, degeneration, and degrees of freedom in the ECSK theory of gravity. II. The Moncrief decomposition*, *Gen. Relat. Grav.* **17** (1985) 953-970.
879. Frąckiewicz K. and Szczyrba W., *A partly reduced system of Hamiltonian equations and the initial value problem in the ECSK theory of gravity*, *Gen. Relat. Grav.* **18** (1986) 411-431.
880. Freedman D.Z. and Khuri R.R., *The Wu-Yang ambiguity revisited*, *Phys. Lett.* **B329** (1994) 263-270.
881. Freidel L., Minic D., and Takeuchi T., *Quantum gravity, torsion, parity violation, and all that*, *Phys. Rev.* **D72** (2005) 104002 (6 pages).
882. Frolov B.N., *Principle of local invariance and Noether theorem*, *Vestnik Mosk. Univ., Ser. fiz., astron.*, n. 6 (1963) 48-58 (in Russian).
883. Frolov B.N., *On the gravitational field equations in the quadratic theory of gravity*, *Sov. J. Phys.* **20** (1977) 411-412 [*Izvestia VUZov, Fizika*, n. 3 (1977) 154-155 (in Russian)].
884. Frolov B.N., *Schwarzschild problem in the quadratic theory of gravity*, *Sov. J. Phys.* **20** (1977) 413-414 [*Izvestia VUZov, Fizika*, n. 3 (1977) 155-156 (in Russian)].
885. Frolov B.N., *Friedmann problem in quadratic theory of gravity*, *Sov. J. Phys.* **21** (1978) 676-678 [*Izvestia VUZov, Fizika*, n. 5 (1978) 148-149 (in Russian)].
886. Frolov B.N., *Tetrad Palatini formalism and quadratic Lagrangians in the gravitational field theory*, *Acta Phys. Pol.* **B9** (1978) 823-829.
887. Frolov B.N., *Generalized conformal invariance and gauge theory of gravity*, in: "Gravity, Particles and Space-time", Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996) 113-144.
888. Frolov B.N., *Poincaré-gauge theory of gravity* (Prometej Publ. House: Moscow, 2003) 160 pp. (in Russian).
889. Frolov B.N., *On foundations of Poincaré-gauge theory of gravity*, *Grav. Cosmol.* **10** (2004) 116-120.
890. Frolov B.N. and Sardanashvily G.A., *Gravitation and gauge fields*, *Sov. J. Phys.* **17** (1974) 1228-1231 [*Izvestia VUZov, Fizika*, n. 9 (1974) 47-51 (in Russian)].
891. Frolov B.N. and Karbanovski V., *The asymptotical solutions near the singular points for perfect fluid configurations in the conformal Poincaré gauge theory of gravity*, *Phys. Lett.* **A169** (1992) 1-4.
892. Frolov I.E. and Zhukovsky V.Ch., *Synchrotron radiation in the standard model extension*, *J. Phys. A: Math. Theor.* **40** (2007) 10625 (16 pages).
893. Fuchs H., *Equations of motion for test particles in a Poincaré gauge field*, *Ann. der Phys.* **43** (1986) 76-82.
894. Fuchs H., *A general form of the equations of motion for test particles with internal structure*, *Ann. der Phys.* **44** (1987) 471-472.
895. Fuchs H., *First integrals for the equations of motion for test particles with internal structure*, *J. Math. Phys.* **32** (1991) 2473-2477.
896. Fujishiro T.T., Hayashi M.J., and Takeshita S., *What is the role of torsion in the universe?*, *Progr. Theor. Phys., Suppl.* **110** (1992) 179-190.
897. Fujishiro T.T., Hayashi M.J., and Takeshita S., *Exact cosmic string solutions based on the continuum theory of dislocations*, *Int. J. Mod. Phys.* **A9** (1994) 4101-4127.
898. Fujishiro T.T., Takeshita S., and Hayashi M.J., *Are the cosmic strings seen as the dislocations?*, *Mod. Phys. Lett.* **A6** (1991) 2237-2242.

899. Fujishiro T.T., Takeshita S., and Hayashi M.J., *The cosmic strings generated from the torsion*, *Mod. Phys. Lett.* **A8** (1993) 491-501.
900. Fukuda M., Yajima S., Higashida Y., Kubota S.-I., Tokuo S., and Kamo Y., *An irreducible form of gamma matrices for HMDS coefficients of the heat kernel in higher dimensions*, *Prog. Theor. Phys.* **121** (2009) 983-1002.
901. Fukui M., *Massless torsion fields. I. The case  $\alpha + \frac{2a}{3} \neq 0$* , *Progr. Theor. Phys.* **71** (1984) 633-647.
902. Fukui M., Hayashi K., and Shirafuji T., *Conformal rescalings applied to Poincaré gauge theory*, *Progr. Theor. Phys.* **74** (1985) 852-865.
903. Fukui M. and Masukawa J., *Massless torsion fields. II. The case  $\alpha + \frac{2a}{3} = 0$* , *Progr. Theor. Phys.* **73** (1985) 75-83.
904. Fukui M. and Masukawa J., *Weak field approximation of new general relativity*, *Progr. Theor. Phys.* **73** (1985) 973-978.
905. Fukui M. and Shirafuji T., *The homogeneous, isotropic universe in new general relativity*, *Progr. Theor. Phys.* **71** (1984) 1063-1073.
906. Fukui T., *Energy-momentum and angular momentum in Poincaré gauge theory of gravity and physical multiplets of torsion fields*, *Progr. Theor. Phys.* **82** (1989) 183-206.
907. Fukui T., Fukuma K., Miyamoto S., Nakano T., Ohtani T., and Tamura Y., *Level shift of hyperfine structure due to the Lorentz gauge field*, *Progr. Theor. Phys.* **78** (1987) 1470-1486.
908. Fukui T., Kawai T., and Saitoh H., *Special relativistic limits of  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **79** (1988) 1420-1430.
909. Fukuma K., *Massless modes of Lorentz gauge fields in Poincaré gauge theory of gravity: The case with  $\alpha + 2a/3 = \beta - 2a/3 = \gamma + 3a/2 = 0$* , *Progr. Theor. Phys.* **107** (2002) 191-210.
910. Fukuma K., Miyamoto S., Nakano T., Ohtani T., and Tamura Y., *Massless Lorentz gauge field consistent with Einstein's gravitational theory*, *Progr. Theor. Phys.* **73** (1985) 874-883.
911. Fukuyama T., *Gauge theory of gravitation and nodal singularity*, *Nuovo Cim.* **B68** (1982) 287-301.
912. Fukuyama T. and Kamimura K., *Hamiltonian formulation of the gauge theory of gravitation. Pure gravity case*, *Nuovo Cim.* **B74** (1983) 93-111.
913. Fulp R.O., Davis W.R., and Norris L.K., *Projective and volume-preserving bundle structures involved in the formulation of  $A(4)$  gauge theories*, *Gen. Relat. Grav.* **18** (1986) 1-29.
914. Fustero X., Gambini R., and Trias A., *Einstein's gravitation as a gauge theory of the Lorentz group*, *Phys. Rev.* **D31** (1985) 3144-3149.

## G

915. Gad R.M., *Energy and momentum densities of cosmological models, with equation of state  $\rho = \mu$ , in general relativity and teleparallel gravity*, *Int. J. Theor. Phys.* **46** (2007) 3263-3274.
916. Gad R.M., *Torsion vector and torsion axial-vector in van Stockum spacetime*, *Mod. Phys. Lett.* **A27** (2012) 1250099 (12 pages).
917. Gad R.M., *Torsion axial-vector and gravitational energy for a stiff fluid model in the theory of teleparallel gravity*, *Int. J. Theor. Phys.* **53** (2014) 307-319.
918. Gaete P. and Heláyel-Neto J.A., *Confining potential from interacting magnetic and torsion fields*, *Mod. Phys. Lett.* **A25** (2010) 471.
919. Gaete P. and Heláyel-Neto J.A., *Aspects of interacting electromagnetic and torsion fields*, *Int. J. Theor. Phys.* **50** (2011) 88-94.

920. Galiakhmetov A.M., *Some exact solutions in self-consistent systems of Einstein-Cartan-Vlasov equations*, *Sov. J. Phys.* **28** (1985) 777-780 [*Izvestia VUZov, Fizika*, n. 10 (1985) 12-16 (in Russian)].
921. Galiakhmetov A.M., *Relativistic statistics of particles with scalar interaction in a space with torsion*, *Sov. J. Phys.* **31** (1988) 848-851 [*Izvestia VUZov, Fizika*, n. 10 (1988) 92-96 (in Russian)].
922. Galiakhmetov A.M., *Kinetics of classical charged particles in a space with torsion*, *Sov. J. Phys.* **32** (1989) 408-412 [*Izvestia VUZov, Fizika*, n. 5 (1989) 103-108 (in Russian)].
923. Galiakhmetov A.M., *Relativistic collisionless gas in an isotropic Universe in Einstein-Cartan theory*, *Sov. J. Phys.* **32** (1989) 461-464 [*Izvestia VUZov, Fizika*, n. 6 (1989) 62-67 (in Russian)].
924. Galiakhmetov A.M., *Cosmological consequences of two sources of torsion in the Einstein-Cartan theory*, *Russ. Phys. J.* **44** (2001) 1316-1322.
925. Galiakhmetov A.M., *On Einstein-Cartan theory with two sources of torsion*, *Grav. Cosmol.* **7** (2001) 33-36.
926. Galiakhmetov A.M., *Cosmology in the Einstein-Cartan theory with two sources of torsion*, *Grav. Cosmol.* **7** (2001) 327-331.
927. Galiakhmetov A.M., *The countable number of models in the cosmology with nonminimal coupling and torsion*, *Russ. Phys. J.* **46** (2003) 662-668.
928. Galiakhmetov A.M., *Exact solutions in Einstein-Cartan cosmology with scalar field*, *Grav. Cosmol.* **10** (2004) 300-304.
929. Galiakhmetov A.M., *Scalar field potential in Einstein-Cartan cosmology*, *Grav. Cosmol.* **12** (2006) 147-150.
930. Galiakhmetov A.M., *The role of a scalar-field potential in the Einstein-Cartan cosmology*, *Russ. Phys. J.* **49** (2006) 642-647.
931. Galiakhmetov A.M., *Exact anisotropic scalar field cosmologies in Einstein-Cartan theory*, *Grav. Cosmol.* **13** (2007) 217-223.
932. Galiakhmetov A.M., *Bianchi-I cosmologies with a scalar field and an ultrarelativistic gas in Einstein-Cartan theory*, *Grav. Cosmol.* **14** (2008) 190-196.
933. Galiakhmetov A.M., *Exact rotating and expanding cosmologies in Einstein-Cartan theory*, *Grav. Cosmol.* **15** (2009) 250-255.
934. Galiakhmetov A.M., *Exact isotropic scalar field cosmologies in Einstein-Cartan theory*, *Class. Quantum Grav.* **27** (2010) 055008.
935. Galiakhmetov A.M., *Cosmology with polynomial potentials of the fourth degree in Einstein-Cartan theory*, *Class. Quantum Grav.* **28** (2011) 105013.
936. Galiakhmetov A.M., *Nonminimally coupled scalar field and perfect fluid in Einstein-Cartan cosmology*, *Gen. Rel. Grav.* **44** (2012) 1043-1056.
937. Galiakhmetov A.M., *Einstein-Cartan and general relativity cosmologies with a nonminimally coupled ghost scalar field*, *Gen. Rel. Grav.* **45** (2013) 275-303.
938. Galiakhmetov A.M., *Multi-component Einstein-Cartan cosmologies*, *Int. J. Theor. Phys.* **52** (2013) 765-778.
939. Galiakhmetov A.M., *Spatially flat FRW models with torsion*, *Grav. Cosmol.* **20** (2014) 90-98.
940. Galiakhmetov A.M., *Multi-component Bianchi I cosmologies with torsion*, *Int. J. Mod. Phys.* **D23** (2014) 1450034 (19 pages).
941. Galiakhmetov A.M., *Flat Friedmann cosmologies with stiff fluid in Einstein-Cartan theory*, *Grav. Cosmol.* **22** (2016) 36-43.

942. Galiakhmetov A.M., *Oscillating universe with two sources of torsion*, in: "Gravitation, Astrophysics, and Cosmology", Proc. Twelfth Asia-Pacific Internat. Conf., Moscow, 28 June–5 July 2015, Eds. V. Melnikov and J.-P. Hsu (World Scientific: Singapore, 2016) p. 262-263.
943. Galitsky V.S., *Cosmology in Einstein-Cartan gravity*, Sov. J. Phys. **31** (1988) 194-198 [Izvestia VUZov, Fizika, n. 3 (1988) 25-29 (in Russian)].
944. Galitsky V.S. and Nurgaliev I.S., *To the problem of stability of spherically symmetric configurations in the Einstein-Cartan theory*, Izvestia VUZov, Fizika, n. 8 (1985) 118-119 (in Russian).
945. Galitsky V.S. and Ponomarev V.N., *Cosmological anisotropic solutions of the Einstein-Cartan equations with cosmological term*, Izvestia VUZov, Fizika, n. 3 (1987) 116 (in Russian).
946. Gambini R. and Herrera L., *Einstein-Cartan theory in the spin formalism*, J. Math. Phys. **21** (1980) 1449-1454.
947. Gambini R. and Trias A., *Path dependent formulation of gravitation as a gauge theory*, Nuovo Cim. **A80** (1984) 488-498.
948. Gangopadhyay D. and Sengupta S., *Duality invariance of cosmological solutions with torsion*, Int. J. Mod. Phys. **A14** (1999) 4953-4966.
949. Garcia De Andrade L.C., *Divergence law in Riemann-Cartan space-time*, Rev. bras. fis. **18** (1988) 89-92.
950. Garcia De Andrade L.C., *Quasi-Maxwellian fields in Riemann-Cartan space-time*, Int. J. Theor. Phys. **28** (1989) 79-81.
951. Garcia De Andrade L.C., *The structure of singularities in space-time with torsion*, Found. Phys. **20** (1990) 403-416.
952. Garcia De Andrade L.C., *Recurrent Riemann-Cartan space-times*, Int. J. Theor. Phys. **29** (1990) 189-192.
953. Garcia De Andrade L.C., *Singularities in spacetimes with torsion*, Int. J. Theor. Phys. **29** (1990) 996-1001.
954. Garcia De Andrade L.C., *Nonlinear electrodynamics and torsion*, Nuovo Cim. **B105** (1990) 1297-1303.
955. Garcia De Andrade L.C., *Nonlinear electrodynamics in space-time with torsion*, Gen. Relat. Grav. **22** (1990) 883-887.
956. Garcia De Andrade L.C., *Ricci recurrent spacetimes with torsion*, Int. J. Theor. Phys. **30** (1991) 235-238.
957. Garcia De Andrade L.C., *Physical structure of the energy-momentum tensor of a dust of spinning particles*, Int. J. Theor. Phys. **30** (1991) 1117-1120.
958. Garcia De Andrade L.C., *Torsion, massive electrodynamics and gravitation induced by scalar fields*, Astropys. and Space Sci. **198** (1992) 79-83.
959. Garcia De Andrade L.C., *Torsion, superconductivity, and massive electrodynamics*, Int. J. Theor. Phys. **31** (1992) 1221-1227.
960. Garcia De Andrade L.C., *The massive photon-torsion interaction and the geometrization of electromagnetic fields*, Nuovo Cim. **B107** (1992) 1407-1411.
961. Garcia De Andrade L.C., *Spin-torsion effects in the Stern-Gerlach experiment*, Nuovo Cim. **B108** (1993) 947-948.
962. Garcia De Andrade L.C., *Torsion and maximum acceleration in the early universe*, Nuovo Cim. **B109** (1994) 433-434.
963. Garcia De Andrade L.C., *Electron gyroscopes to test torsion gravity?*, Nuovo Cim. **B109** (1994) 1123-1127.

964. Garcia De Andrade L.C., *Non-Riemannian stress space structure of gravitation and ferromagnetism from distant parallelism*, in: "Continuum models and discrete systems"(Proc. of 8th Int. Symp, June 11-16, 1995, Varna, Bulgaria), Ed. K.Z. Markov (World Scientific: Singapore, 1996) 437-443.
965. Garcia De Andrade L.C., *Placing limits on Cartan's torsion from non-Riemannian crystals*, in: "Gravity, Particles and Space-time" Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996), 267-274.
966. Garcia De Andrade L.C., *Space-time defects: Torsion walls*, *Mod. Phys. Lett.* **A12** (1997) 2005-2009.
967. Garcia De Andrade L.C., *Spacetime defects: domain walls and torsion*, *J. Math. Phys.* **39** (1998) 372-379.
968. Garcia De Andrade L.C., *On non-Riemannian domain walls*, *Gen. Rel. Grav.* **30** (1998) 1629-1637.
969. Garcia De Andrade L.C., *Non-Riemannian cosmic walls as boundaries of spinning matter with torsion*, *Mod. Phys. Lett.* **A13** (1998) 1179-1184.
970. Garcia De Andrade L.C., *On the stability of wormholes in Weitzenböck space-time*, *Mod. Phys. Lett.* **A13** (1998) 2045-2047.
971. Garcia De Andrade L.C., *Distributional torsion of cosmic strings with polarized spinning matter*, *Mod. Phys. Lett.* **A13** (1998) 2227-2230.
972. Garcia De Andrade L.C., *Negative energy density in spinning matter sources in wormhole space-times with torsion*, *Mod. Phys. Lett.* **A13** (1998) 3069-3072.
973. Garcia De Andrade L.C., *On time dislocation solution of Einstein field equations*, *Mod. Phys. Lett.* **A14** (1999) 93-97.
974. Garcia De Andrade L.C., *Distributional torsion of hybrid defects*, *Mod. Phys. Lett.* **A14** (1999) 1719-1723.
975. Garcia De Andrade L.C., *Inflating torsion defects and dilatonic domain walls*, *Class. Quantum Grav.* **16** (1999) 2097-2103.
976. Garcia De Andrade L.C., *On dilaton solutions of de Sitter inflation and primordial spin-torsion density fluctuations*, *Phys. Lett.* **B468** (1999) 28-30.
977. Garcia De Andrade L.C., *Geodesics on non-Riemannian geometric theory of planar defects*, *Phys. Lett.* **A256** (1999) 291-293.
978. Garcia De Andrade L.C., *Averaged weak energy condition violation for wormhole geometries with torsion*, *Mod. Phys. Lett.* **A15** (2000) 1321-1327.
979. Garcia De Andrade L.C., *Inflaton potential reconstruction in Einstein-Cartan cosmology*, *Mod. Phys. Lett.* **A16** (2001) 1353-1360.
980. Garcia De Andrade L.C., *Astrophysical and terrestrial probes to test Einstein-Cartan gravity*, *Class. Quantum Grav.* **18** (2001) 3907-3915.
981. Garcia De Andrade L.C., *Placing limits on spin-torsion densities from FRW cosmological model and astronomical data*, *Grav. Cosmol.* **8** (2002) 241-242.
982. Garcia De Andrade L.C., *Superfluids in non-Riemannian spacetime background*, *Europhys. Lett.* **63** (2003) 649.
983. Garcia De Andrade L.C., *Holonomy and torsion defects in  $(2 + 1)D$  spacetime*, *Europhys. Lett.* **66** (2004) 296.
984. Garcia De Andrade L.C., *Non-Riemannian geometry of vortex acoustics*, *Phys. Rev.* **D70** (2004) 064004 (5 pages).
985. Garcia De Andrade L.C., *Limits to global rotation and spin density in Einstein-Cartan Cosmology*, *Astrophys. and Space Sci.* **289** (2004) 161-168.

986. Garcia De Andrade L.C., *On spin-driven inflation from inflation fields in General Relativity and COBE data*, *Astropys. and Space Sci.* **294** (2004) 213-217.
987. Garcia De Andrade L.C., *Non-Riemannian vortex geometry of rotational viscous fluids and breaking of the acoustic Lorentz invariance*, *Phys. Lett.* **A339** (2005) 188-193.
988. Garcia De Andrade L.C., *On the necessity of non-Riemannian acoustic spacetime in fluids with vorticity*, *Phys. Lett.* **A346** (2005) 327-329.
989. Garcia De Andrade L.C., *Constraints on torsion from the bosonic sector of Lorentz violation and magnetogenesis data*, *Nucl. Phys.* **B847** (2010) 485-490.
990. Garcia De Andrade L.C., *Galactic dynamos and slow decay of magnetic fields from torsion modes of Lorentz violation*, *Phys. Lett.* **B696** (2011) 1-4.
991. Garcia De Andrade L.C., *Torsion bounds from CP violation  $\alpha^2$ -dynamo in axion-photon cosmic plasma*, *Mod. Phys. Lett.* **A26** (2011) 2863.
992. Garcia De Andrade L.C., *Faraday rotation and primordial magnetic fields constraints on ultraviolet Lorentz violation with spacetime torsion*, *Mod. Phys. Lett.* **A27** (2012) 1250059 (5 pages).
993. Garcia De Andrade L.C., *Primordial magnetic fields and dynamos from parity violated torsion*, *Phys. Lett.* **B711** (2012) 143-146.
994. Garcia De Andrade L.C., *Cosmological magnetic fields as string dynamo seeds and axion fields in torsioned spacetime*, *JCAP* **08** (2014) 023.
995. Garcia De Andrade L.C., *Lorentz violation from torsion trace and non-minimal coupling in radio galactic dynamos*, *Phys. Rev.* **D84** (2011) 127504 (5 pages).
996. Garcia De Andrade L.C., *Primordial magnetic fields and dynamos from parity violated torsion*, *Phys. Lett.* **B711** (2012) 143-146.
997. Garcia De Andrade L.C., *Torsion constraints from cosmological magnetic field and QCD domain walls*, *Mod. Phys. Lett.* **A29** (2014) 1450171 (9 pages).
998. Garcia De Andrade L.C., *Cosmological and astrophysical consequences from the magnetic dynamo equation in torsioned spacetime and teleparallel gravity*, *Class. Quantum Grav.* **23** (2016) 025006.
999. Garcia De Andrade L.C. and Hammond R., *Einstein-Cartan-Proca geometry*, *Gen. Relat. Grav.* **27** (1995) 1259-1263.
1000. Garcia De Andrade L.C. and de Oliveira J.U.C.L., *Nonlinear electromagnetic effects induced by torsion*, in: "General relativity and gravitational physics". *Proc. of the 9th Ital. Conf. (Capri, Italy, September 25-28, 1990)*, Eds. R. Cianci, R. de Ritis, G. Marmo, C. Rubano, P. Scudellaro, and M. Francaviglia (World Scientific: Singapore, 1991) 446-449.
1001. Garcia De Andrade L.C. and Lima C.A.S., *Non-Riemannian geometry of macroscopic spin distributions*, *Int. J. Theor. Phys.* **36** (1997) 1227-1230.
1002. Garcia De Andrade L.C. and Lima C.A.S., *Spectrum of density fluctuations in Einstein-Cartan cosmology*, *Grav. Cosmol.* **7** (2001) 169-170.
1003. Garcia De Andrade L.C. and Lopes M., *Detecting torsion from massive electrodynamics*, *Gen. Relat. Grav.* **25** (1993) 1101-1106.
1004. Garcia De Andrade L.C. and de Oliveira J.U.C.L., *Nonlinear electromagnetism effects induced by torsion*, *Int. J. Theor. Phys.* **30** (1991) 1259-1262.
1005. Garcia De Andrade L.C. and Sivaram C., *Torsion, massive electrodynamics and primordial magnetic fields in the early Universe*, *Astropys. and Space Sci.* **209** (1993) 109-111.
1006. Garcia De Andrade L.C., Oguri V., Lopes M., and Hammond R., *On the energy splitting of spectral lines induced by the torsion field*, *Nuovo Cim.* **B107** (1992) 1167-1171.

1007. García A., Hehl F.W., Heinicke C., and Macías A., *Exact vacuum solution of a (1 + 2)-dimensional Poincaré gauge theory: BTZ solution with torsion*, *Phys. Rev.* **D67** (2003) 124016 (7 pages).
1008. García A., Hehl F.W., Lämmerzahl C., Macías A., and Socorro J., *Plebanski-Demianski-like solutions in metric-affine gravity*, *Class. Quantum Grav.* **15** (1998) 1793-1799.
1009. García A., Lämmerzahl C., Macías A., Mielke E.W., and Socorro J., *Colliding waves in metric-affine gravity*, *Phys. Rev.* **D57** (1998) 3457-3462.
1010. García A., Macías A., and Socorro J., *Electrovacuum sector of the MAG theories*, *Class. Quantum Grav.* **16** (1999) 93-100.
1011. García A., Macías A., D. Puetzfeld, and Socorro J., *Plane-fronted waves in metric-affine gravity*, *Phys. Rev.* **D62** (2000) 044021 (7 pages).
1012. Garcia G.Q., de S. Oliveira J.R., Bakke K., and Furtado C., *Fermions in Gödel-type background space-times with torsion and the Landau quantization*, *Eur. Phys. J. Plus* **132** (2017) 123 (15 pages).
1013. Garecki J., *The superenergy tensors*, *Acta Phys. Pol.* **B8** (1977) 159-167.
1014. Garecki J., *The total superenergy tensor in general relativity and in Einstein-Cartan theory and the global superenergetic quantities of a closed system*, *Acta Phys. Pol.* **B9** (1978) 291-302.
1015. Garecki J., *Superenergy tensors in the Einstein-Cartan theory of gravitation*, *Acta Phys. Pol.* **B12** (1981) 739-746.
1016. Garecki J., *On the gravitational theory with quadratic Lagrangian*, *Acta Phys. Pol.* **B14** (1983) 713-722.
1017. Garecki J., *Gauge theory of gravitation with quadratic Lagrangian  $L_g = \alpha \Omega^i_j \wedge \eta_i^j + \beta \Omega^i_j \wedge * \Omega_i^j + \gamma \Theta^i \wedge * \Theta_i$  containing Einsteinian term  $\alpha \Omega^i_j \wedge \eta_i^j$  and spherically symmetric solutions to its field equations*, in: "On Relativity Theory. Proc. of Sir A.Eddington Centenary Symp.", Eds. Y. Choquet-Bruhat and T.M. Karade (World Scientific: Singapore, 1985) vol. **2**, 232-260.
1018. Garecki J., *New physical interpretation of the Bel-Robinson tensor*, *Class. Quantum Grav.* **2** (1985) 403-408.
1019. Garecki J., *Homogeneous and isotropic cosmology in gauge gravitational theory with gravitational lagrangian*, *Acta Phys. Pol.* **B18** (1987) 147-161.
1020. Garecki J., *Revised gauge theory of gravitation with quadratic Lagrangian  $L_g = \alpha(\Omega^i_k \wedge \eta_i^k + Q^i \wedge * Q_i) + \beta \Omega_k^i \wedge * \Omega_i^k$* , *Acta Phys. Pol.* **B20** (1989) 997-1014.
1021. Garecki J., *A gauge theory of gravitation based on a specific Lagrangian*, *Gen. Relat. Grav.* **22** (1990) 111-119.
1022. Garecki J., *New cosmological models in the framework of a gauge theory of gravity*, *Gen. Relat. Grav.* **25** (1993) 257-265.
1023. Garecki J., *Energy-momentum and angular momentum of isotropic homogeneous cosmological models in a gauge theory of gravity*, *Int. J. Theor. Phys.* **34** (1995) 2307-2319.
1024. Garkun A.S., Kudin V.I., and Minkevich A.V., *Analysis of regular inflationary cosmological models with two torsion functions in Poincaré gauge theory of gravity*, *Int. J. Mod. Phys.* **A25** (2010) 2005-2022.
1025. Garkun A.S., Kudin V.I., and Minkevich A.V., *To theory of asymptotically stable accelerating Universe in Riemann-Cartan spacetime*, *JCAP* **12** (2014) 027.
1026. Gasperini M., *Gravitation and the Einstein-Sciama force*, *Lett. Nuovo Cim.* **21** (1978) 285-287.
1027. Gasperini M., *Lie-admissible theory of gravity*, *Nuovo Cim.* **B81** (1984) 7-20.

1028. Gasperini M., *On the geometrical structure of a Lie-isotopic gravity*, *Hadronic J.* **7** (1984) 650-658.
1029. Gasperini M., *Lie-isotopic theory of gravity*, *Nuovo Cim.* **A83** (1984) 309-326.
1030. Gasperini M., *Lie-isotopic gravity with matter. An effective theory of gravity for testing local Lorentz symmetry*, *Hadronic J.* **8** (1985) 52-59.
1031. Gasperini M., *Spin-dominated inflation in the Einstein-Cartan theory*, *Phys. Rev. Lett.* **56** (1986) 2873-2876.
1032. Gasperini M., *Spin contributions to an inflationary phase in the very early universe*, In: "General relativity and gravitational physics". Proc. 7th Italian Conf. Rapallo, Italy, September 3-7, 1986, Eds. U. Bruzzo, R. Cianci, E. Massa (Singapore: World Scientific, 1987) 89-93.
1033. Gasperini M., *Short-range interactions in gravitational theories with torsion and quadratic Lagrangians*, *Phys. Lett.* **B205** (1988) 517-520.
1034. Gasperini M., *A thermal interpretation of the cosmological constant*, *Class. Quantum Grav.* **5** (1988) 521-531.
1035. Gasperini M., *Repulsive gravity in the very early Universe*, *Gen. Rel. Grav.* **30** (1998) 1703-1709.
1036. Geng C.-Q., Lai C., Luo L.-W., and Tseng H.-H., *Kaluza-Klein theory for teleparallel gravity*, *Phys. Lett.* **B737** (2014) 248-250.
1037. Geng C.-Q., Lee C.-C., and Saridakis E.N., *Observational constraints on teleparallel dark energy*, *JCAP* **01** (2012) 002.
1038. Geng C.-Q., Lee C.-C., Saridakis E.N., and Wu Y.-P., "Teleparallel" dark energy, *Phys. Lett.* **B704** (2011) 384-387.
1039. Geng C.-Q., Lee C.-C., and Tseng H.-H., *Asymptotic cosmological behavior of scalar-torsion mode in Poincaré gauge theory*, *Phys. Rev.* **D87** (2013) 027301 (5 pages).
1040. Geng C.-Q., Luo L.-W., and Tseng H.-H., *Teleparallel gravity in five-dimensional theories*, *Class. Quantum Grav.* **31** (2014) 185004.
1041. Geng W.T. and Duan Y.S., *Space-time torsion and the quantization of spin*, *Acta Phys. Sinica (Overseas Edition)* **7** (1998) 249-257.
1042. Gerbert Ph. de Sousa, *On spin and (quantum) gravity in (2+1) dimensions*, *Nucl. Phys.* **B346** (1990) 440-472.
1043. Germán G., *Brans-Dicke-type models with torsion*, *Phys. Rev.* **D32** (1985) 3307-3308.
1044. Germán G., *On Kaluza-Klein theory with torsion*, *Class. Quant. Grav.* **2** (1985) 455-460.
1045. Germán G., Macías, and Obregón O., *Kaluza-Klein approach in higher-dimensional theories of gravity with torsion*, *Class. Quantum Grav.* **10** (1993) 1045-1053.
1046. Geronimi C., *Connexion et théorie quantique des champs sur un espace-temps courbe*, *C. R. Acad. Sci. Paris, Ser. A* **272** (1971) 1764-1766.
1047. Geyer B., Gitman D., and Shapiro I.L., *Path integral and pseudoclassical action for spinning particle in external electromagnetic and torsion fields*, *Int. J. Mod. Phys.* **A15** (2000) 3861-3876.
1048. Giachetta G., *Nonlinear realizations of the diffeomorphism group in metric-affine gauge theory of gravity*, *J. Math. Phys.* **40** (1999) 939-954.
1049. Giachetta G., Giambó R., and Mangiarotti L., *Conservation laws in metric-affine gravitation theories: superpotentials*, *Nuovo Cim.* **B112** (1997) 1163-1174.
1050. Giachetta G. and Mangiarotti L., *Affine theories of gravitation with torsion*, in: "Gravity, Particles and Space-time", Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996) 255-266.

1051. Giachetta G. and Mangiarotti L., *Projective invariance and Einstein's equations*, *Gen. Rel. Grav.* **29** (1997) 5-18.
1052. Giachetta G., Mangiarotti L., and Saltarelli A., *Metric-affine gravitation theory and superpotentials*, *Nuovo Cim.* **B112** (1997) 751-760.
1053. Giachetta G. and Sardanashvily G., *Stress-energy-momentum of affine-metric gravity. Generalized Komar superpotential*, *Class. Quantum Grav.* **13** (1996) L67-L71.
1054. Giachetta G., and Sardanashvily G., *Energy-momentum superpotential in gravitation theory*, in: "Gravity, Particles and Space-time", Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996) 471-506.
1055. Giachetti R., Ricci R., and Sorace E., *Tetrad fields and metric tensors in gauge theory of gravitation*, *Lett. Math. Phys.* **5** (1981) 85-91.
1056. Giachetti R., Ricci R., and Sorace E., *Hamiltonian formulation for the gauge theory of the gravitational coupling*, *J. Math. Phys.* **22** (1981) 1703-1710.
1057. Giacomini A., Troncoso R., and Willison S., *Junction conditions in general relativity with spin sources*, *Phys. Rev.* **D73** (2006) 104014 (11 pages).
1058. Giannopoulos A., *Generalized torsion and curvature*, *Class. Quantum Grav.* **8** (1991) 571-585.
1059. Gibbons G.W. and Gielen S., *Deformed general relativity and torsion*, *Class. Quantum Grav.* **26** (2009) 135005.
1060. Gieres F., *BRS algebras of Poincaré and conformal supergravity*, *Gen. Relat. Grav.* **21** (1989) 237-256.
1061. Gies H. and SLippoldt S., *Fermions in gravity with local spin-base invariance*, *Phys. Rev.* **D89** (2014) 064040 (19 pages).
1062. Gilkey P., Nikčević S., and Simon U., *Geometric realizations, curvature decompositions, and Weyl manifolds*, *J. Differ. Geom.* **61** (2011) 270-275.
1063. Gladchenko M.S., Ponomariev V.N., and Zhytnikov V.V., *PPN metric and PPN torsion in the quadratic Poincaré gauge theory of gravity*, *Phys. Lett.* **B241** (1990) 67-69.
1064. Gladchenko M.S., Zhytnikov V.V., and Ponomariev V.N., *Post-Newtonian approximation in the gauge theory of gravity*, in: "Gravitation and Electromagnetism" (University Publ. House: Minsk, 1989) **4**, 73-92 (in Russian).
1065. Gladchenko M.S. and Zhytnikov V.V., *Post-Newtonian effects in the quadratic Poincaré gauge theory of gravitation*, *Phys. Rev.* **D50** (1994) 5060-5071.
1066. Gökeler M. and Schücker T., *Differential geometry, gauge theory and gravity* (Cambridge Univ. Press: Cambridge, 1987).
1067. Godina M., Matteucci P., and Vickers J.A., *Metric-affine gravity and the Nester-Witten 2-form*, *J. Geom. Phys.* **39** (2001) 265-275.
1068. Godinho C.F.L. and Helayël Neto J.A., *On the role of a torsion-like field in a scenario for the Spin Hall Effect*, *Ann. Phys. (Berlin)* **528** (2016) 597-602.
1069. Godlewski P., *Generally-covariant and  $GL(N, R)$ -invariant models of self-interacting field of linear frames*, *Rept. Math. Phys.* **35** (1995) 77-99.
1070. Goenner H.F., *Unified field theories: from Eddington and Einstein up to now*, in: "Relativistic Astrophysics and Cosmology. Proc. of the Sir A.Eddington Centenary Symp.", Eds. V. DeSabbata and T.M. Karade (World Scientific: Singapore, 1984), vol. **1**, 137-169.
1071. Goenner H.F.M., *Exact solutions of the  $SO(3)$ -Yang-Mills-Higgs-system coupled to gravitation within Poincaré gauge theory*, *Wiss. Z. Friedrich-Schiller-Univ. Jena. Naturwiss.* **39** (1990) 499-554.

1072. Goenner H., *Alternative to the Palatini method: A new variational principle*, *Phys. Rev.* **D81** (2010) 124019 (6 pages).
1073. Goenner H.F. and Müller-Hoisen F., *Spatially homogeneous and isotopic spaces in theories of gravitation with torsion*, *Class. Quant. Grav.* **1** (1984) 651-672.
1074. Godazgar H., Godazgar M., and Nicolai H., *Einstein-Cartan calculus for exceptional geometry*, *JHEP* **06** (2014) 021 (27 pages).
1075. Gogala B., *Torsion and related concepts: an introductory overview*, *Int. J. Theor. Phys.* **19** (1980) 573-586.
1076. Gogala B., *Rotation and acceleration as parallel transport in a space-time with torsion*, *Int. J. Theor. Phys.* **19** (1980) 587-597.
1077. Gogala B., *Field equations of the gauge theory of gravitation originated from a quadratic Lagrangian with torsion*, *Int. J. Theor. Phys.* **22** (1983) 887-893.
1078. Goldthorpe W.H., *Spectral geometry and  $SO(4)$  gravity in a Riemann-Cartan spacetime*, *Nucl. Phys.* **B170[FS1]** (1980) 307-328.
1079. Golovnev A., Koivisto T., and Sandstad M., *Effectively nonlocal metric-affine gravity*, *Phys. Rev.* **D93** (2016) 064081 (10 pages).
1080. Golubyatnikov A.N. and Petrosyan P.A., *Curvature and torsion of space in the Dirac theory*, in: "Probl. Theor. Grav. and Elem. Particles", Ed. K.P. Stanyukovich (Atomizdat: Moscow) **5** (1974) 61-64 (in Russian).
1081. Gonçalves B., Dias M.M.Jr., and Ribeiro B.J., *The exact Foldy-Wouthuysen transformation for a Dirac theory with the complete set of CPT-Lorentz violating terms: The torsion field case*, *PoS FFP14* (2016) 112 (4 pages).
1082. Gonçalves B., Obukhov Yu.N., and Shapiro I.L., *Exact Foldy-Wouthuysen transformation for gravitational waves and magnetic field background*, *Phys. Rev.* **D75** (2007) 124023 (7 pages).
1083. Gonçalves B., Obukhov Yu.N., and Shapiro I.L., *Exact Foldy-Wouthuysen transformation for a Dirac spinor in torsion and other CPT and Lorentz violating backgrounds*, *Phys. Rev.* **D80** (2009) 125034 (7 pages).
1084. González P.A. and Vásquez Y., *Exact solutions in 3D gravity with torsion*, *JHEP* **8** (2011) 89.
1085. Gordeeva I.A., Panzhenskii V.I., and Stepanov S.E., *Riemann-Cartan manifolds*, *Journal of Mathematical Sciences*, **169** (2010) 342-361.
1086. Gottlieb D., Hojman R., Rodrigues L.H., and Zamorano N., *Yukawa potential in a Schwarzschild background*, *Nuovo Cim.* **B80** (1984) 62-70.
1087. Gotzes S., and Hishfeld A.C., *A geometric formulation of the  $SO(3,2)$  theory of gravity*, *Ann. Phys. (USA)* **203** (1990) 410-418.
1088. Gowdy R.H., *Affine projection tensor geometry: Decomposing the curvature tensor when the connection is arbitrary and the projection is tilted*, *J. Math. Phys.* **35** (1994) 1274-1301.
1089. Grachev A.V., Nesterov A.I., and Ovchinnikov S.G., *The gauge theory of point defects*, *Physica status solidi (b)* **156** (1989) 1521-3951.
1090. Gregorash D. and Papini G., *Weyl-Dirac theory with torsion*, *Nuovo Cim.* **B55** (1980) 37-51.
1091. Gregorash D. and Papini G., *Weyl-Dirac theory with torsion, II. Foundations and conservation equations*, *Nuovo Cim.* **B56** (1980) 21-38.
1092. Gregorash D. and Papini G., *Torsion in a multiply connected Weyl-Dirac geometry*, *Nuovo Cim.* **B64** (1981) 55-66.
1093. Gregorash D. and Papini G., *The equations of motion in a conformally invariant theory of gravitation and electromagnetism*, *Nuovo Cim.* **B70** (1982) 259-267.

1094. Gresning G., *Induced gravity for nonzero torsion*, *Phys. Lett.* **B169** (1986) 333-336.
1095. Gresning D. and Gresning G., *General relativity as a gauge theory of the Poincaré group, the symmetric momentum tensor of both matter and gravity, and gauge fixing conditions*, *Phys. Rev.* **D28** (1983) 286-296.
1096. Gresning D. and Gresning G., *How to fix the gauge in internal and external gauge theories*, *Gen. Relat. Grav.* **15** (1983) 985-996.
1097. Grib A.A. and Poberii E.A., *Gauge approach in gravity and strong interactions*, in: "Probl. Theor. Grav. and Element. Particles", Ed. K.P. Stanyukovich (Energoatizdat: Moscow, 1985) **16**, 4-17 (in Russian).
1098. Grib Lucas T., Obukhov Yu.N., and Pereira J.G., *Regularizing role of teleparallelism*, *Phys. Rev.* **D80** (2009) 064043 (11 pages).
1099. Grib Lucas T., and Pereira J.G., *A Hodge dual for soldered bundles*, *J. Phys. A: Math. Theor.* **42** (2009) 035402.
1100. Griffiths J.B., *Ghost neutrinos in Einstein-Cartan theory*, *Phys. Lett.* **A75** (1980) 441-442.
1101. Griffiths J.B., *Neutrino fields in Einstein-Cartan theory*, *Gen. Relat. Grav.* **13** (1981) 227-237.
1102. Griffiths J.B. and Jogia S., *A spin coefficient approach to Weyssenhoff fluids in Einstein-Cartan theory*, *Gen. Relat. Grav.* **14** (1982) 137-149.
1103. Grignani G. and Nardelli G., *Gravity and the Poincaré group*, *Phys. Rev.* **D45** (1992) 2719-2731.
1104. Grignani G. and Nardelli G., *Gravity in 2+1 dimensions coupled to point-like sources: a flat Chern-Simons gauge theory equivalent to Einstein's*, *Nucl. Phys.* **B370** (1992) 491-508.
1105. Grignani G. and Nardelli G., *Chern-Simons gravity from (3+1)-dimensional gravity*, *Phys. Lett.* **B300** (1993) 38-43.
1106. Grignani G. and Nardelli G., *Canonical analysis of Poincaré gauge theories for two-dimensional gravity*, *Class. Quantum Grav.* **10** (1993) 2569-2580.
1107. Grignani G. and Nardelli G., *Poincaré gauge theories for lineal gravity*, *Nucl. Phys.* **B412** (1994) 320-344.
1108. Gromov A. and Abanov A.G., *Thermal Hall effect and geometry with torsion*, *Phys. Rev. Lett.* **114** (2015) 016802 (6 pages).
1109. Gronwald F., *On non-Riemannian parallel transport in Regge calculus*, *Class. Quantum Grav.* **12** (1995) 1181-1189.
1110. Gronwald F., *Metric-affine gauge theory of gravity. I. Fundamental structure and field equations*, *Intern. J. Mod. Phys.* **D6** (1997) 263-303.
1111. Gronwald F., *A note on gauge covariant translations in the gauge approach to gravity*, *Acta Phys. Pol.* **B29** (1998) 1121-1129.
1112. Gronwald F., *BRST antifield treatment of metric-affine gravity*, *Phys. Rev.* **D57** (1998) 961-970.
1113. Gronwald F. and Hehl F.W., *Stress and hyperstress as fundamental concepts in continuum mechanics and in relativistic field theory*, in: "Advances in Modern Continuum Dynamics", *International Conference in Memory of Antonio Signorini, Isola d'Elba, June 1991*, Ed. G. Ferrarese (Bologna: Pitagora Editrice, 1993) 1-32.
1114. Gronwald F. and Hehl F.W., *On the gauge aspects of gravity*, in: *Proc. of the 14th Course of the International School of Cosmology and Gravitation "Quantum Gravity" (Erice, Italy, May 1995)*, Eds. P.G. Bergmann, V. De Sabbata and H.-J. Treder (World Scientific, 1996) 148-198.
1115. Gronwald F., Muench U., and Hehl F.W., *Nongravitational vacuum energy and the volume element of spacetime*, *Hadronic J.* **21** (1998) 3-17.

1116. Gronwald F., Muench U., Macías A., and Hehl F.W., *Volume elements of space-time and a quartet of scalar fields*, *Phys.Rev.* **D58** (1998) 084021 (4 pages).
1117. Grosse H., Kummer W., Prešnajder P., and Schwarz D.J., *Nonlinear gauge symmetry in  $R^2$  gravity in two-dimensions*, *Czech. J. Phys.* **42** (1992) 1325-1329.
1118. Grosse H., Kummer W., Prešnajder P., and Schwarz D.J., *Novel symmetry of non-Einsteinian gravity in two dimensions*, *J. Math. Phys.* **33** (1992) 3892-3900.
1119. Grundberg J. and Lindström U., *Some torsion potentials*, *Gen. Relat. Grav.* **18** (1986) 1037-1044.
1120. Gruver C., Hammond R., and Kelly P.F., *Tensor-scalar torsion*, *Mod. Phys. Lett.* **A16** (2001) 113-119.
1121. Gu J.-A., Lee C.-C., and Geng C.-Q., *Teleparallel dark energy with purely non-minimal coupling to gravity*, *Phys. Lett.* **B718** (2013) 722-726.
1122. Guendelman E.I. and Kaganovich A.B., *Principle of nongravitating vacuum energy and some of its consequences*, *Phys. Rev.* **D53** (1996) 7020-7025.
1123. Guendelman E.I. and Kaganovich A.B., *Gravitational theory without the cosmological constant problem*, *Phys. Rev.* **D55** (1997) 5970-5980.
1124. Guendelman E.I. and Kaganovich A.B., *Gravitational theory without the cosmological constant problem, symmetries of space-filling branes, and higher dimensions*, *Phys. Rev.* **D56** (1997) 3548-3554.
1125. Guendelman E.I. and Kaganovich A.B., *Gravitation and cosmology without the cosmological constant problem*, *Hadronic J.* **21** (1998) 19-74.
1126. Guilfoyle B.S. and Nolan B.C., *Yang's gravitational theory*, *Gen. Relat. Grav.* **30** (1998) 473-495.
1127. Gurtug O. and Halilsoy M., *Colliding wave solutions in a symmetric non-metric theory*, *Int. J. Theor. Phys.* **48** (2009) 139-149.
1128. Gusynin V.P., Gorbar E.V., and Romankov V.V., *Heat kernel expansions for nonminimal differential operators and manifold with torsion*, *Nucl. Phys.* **B362** (1991) 449-471.
1129. Gvozdev A.A. and Pronin P.I., *Quantum interaction of space-time torsion with fermions in mean field approximation*, in: "Fundamental Interactions", Ed. V.N. Ponomarev (Pedagogical Inst. Press: Moscow, 1984) 160-175 (in Russian).
1130. Gvozdev A.A. and Pronin P.I., *Quantum statistics and the temperature effects in the interaction of fermions with torsion*, *Vestnik Mosk. Univ., ser. fiz., astron.*, **26**, n. 1 (1985) 29-33 (in Russian).
1131. Gvozdev A.A. and Pronin P.I., *Four-fermion interaction as a quantum source of the torsion field*, *Sov. J. Phys.* **28** (1985) 4-8 [*Izvestia VUZov, Fizika*, n. 1 (1985) 7-12 (in Russian)].

## H

1132. Haghani Z., Harko T., Sepangi H.R., and Shahidi S., *Weyl-Cartan-Weitzenböck gravity as a generalization of teleparallel gravity*, *JCAP* **10** (2012) 061.
1133. Haghani Z., Harko T., Sepangi H.R., and Shahidi S., *Weyl-Cartan-Weitzenböck gravity through Lagrange multiplier*, *Phys. Rev.* **D88** (2013) 044024 (12 pages).
1134. Haghani Z., Khosravi N., and Shahidi S., *The Weyl-Cartan Gauss-Bonnet gravity*, *Class. Quantum Grav.* **32** (2015) 215016.
1135. Haider F. and Kummer W., *Quantum functional integration of noneinsteinian gravity in  $d=2$* , *Int. J. Mod. Phys.* **A9** (1994) 207-220.

1136. Hajicek P. and Stewart J., *Can spin avert singularities?*, *Nature Phys. Sci.* **244** (1973) 96.
1137. Halbwachs F., *Theorie relativiste des fluides à spin* (Gauthier-Villars: Paris, 1960).
1138. Halpern L., *A geometrical theory of spin motion*, *Int. J. Theor. Phys.* **23** (1984) 843-862.
1139. Halpern L., *Equivalence of spin and angular momentum in a unified theory*, In: "General relativity". *Proc. 4th Marcel Grossmann Meeting, Rome, Italy, June 17-21, 1985, Ed. R. Ruffini* (Amsterdam: North-holland, 1986) Pt. B, 1233-1245.
1140. Halpern L., *Geometrical structure of gravitation and matter fields*, *Int. J. Theor. Phys.* **33** (1994) 401-424.
1141. Hammad F., *An entropy functional for Riemann-Cartan space-times*, *Int. J. Theor. Phys.* **51** (2012) 362-373.
1142. Hammad F., *Erratum to: An entropy functional for Riemann-Cartan space-times*, *Int. J. Theor. Phys.* **52** (2013) 4592-4593.
1143. Hamamoto S., *Poincaré gauge theory with Weyl-type Lagrangian and gravitation*, *Progr. Theor. Phys.* **58** (1977) 710-712.
1144. Hamamoto S., *Invariant theoretical consideration on nonlinear realizations*, *Progr. Theor. Phys.* **60** (1978) 315-316.
1145. Hamamoto S., *Invariant theoretical consideration on the nonlinear realizations of general covariance group*, *Progr. Theor. Phys.* **60** (1978) 1910-1921.
1146. Hamamoto S., *Nonlinear realization of general covariance group. Theory with massless tensor graviton and massive vector torsion*, *Progr. Theor. Phys.* **61** (1979) 326-333.
1147. Hamamoto S., *The structure of general covariance group on superspace*, *Progr. Theor. Phys.* **62** (1979) 823-831.
1148. Hamamoto S., *Nonlinear realization of graded conformal group*, *Progr. Theor. Phys.* **63** (1980) 2095-2111.
1149. Hamamoto S., *Nonlinear realization of affine group in superspace*, *Progr. Theor. Phys.* **64** (1980) 1453-1465.
1150. Hammon K.S. and Norris L.K., *Space-time geometry and the bundle of biframe. I. Geometrical developments*, *Nuovo Cim.* **B107** (1992) 385-405.
1151. Hammon K.S. and Norris L.K., *Space-time geometry and the bundle of biframe. II. Applications to the RMW theory*, *Nuovo Cim.* **B107** (1992) 407-426.
1152. Hammon K.S. and Norris L.K., *The affine geometry of the Lanczos H-tensor formalism*, *Gen. Relat. Grav.* **25** (1993) 55-80.
1153. Hammond R.T., *Gravitation, torsion, and electromagnetism*, *Gen. Relat. Grav.* **20** (1988) 813-827.
1154. Hammond R.T., *Einstein-Maxwell theory from torsion*, *Class. Quantum Grav.* **6** (1989) L195-L198.
1155. Hammond R.T., *Dynamic torsion from a linear Lagrangian*, *Gen. Relat. Grav.* **22** (1990) 451-456.
1156. Hammond R.T., *Propagating torsion as a massive vector field*, *Class. Quantum Grav.* **7** (1990) 2107-2112.
1157. Hammond R.T., *Second order equations and quadratic Lagrangians*, *J. Math. Phys.* **31** (1991) 2221-2224.
1158. Hammond R.T., *Magnetic charge type equation from torsion*, *Gen. Relat. Grav.* **23** (1991) 973-980.

1159. Hammond R.T., *Gauge invariant electromagnetic coupling with torsion potential*, *Gen. Relat. Grav.* **23** (1991) 1195-1203.
1160. Hammond R.T., *Non-existence of static spherically symmetric solutions with torsion potential*, *Class. Quantum Grav.* **8** (1991) L175-L177.
1161. Hammond R.T., *Naked singularities with vector torsion gravity*, *Class. Quantum Grav.* **9** (1992) L171-L172.
1162. Hammond R.T., *Geometric foundation for monopoles*, *Nuovo Cim.* **B108** (1993) 725-738.
1163. Hammond R., *Automatic conservation law and gauge invariance in space-time with torsion potential*, *Nuovo Cim.* **B109** (1994) 315-317.
1164. Hammond R., *Stringy torsion*, *Nuovo Cim.* **B109** (1994) 319-323.
1165. Hammond R., *Nonlinear quantum equation from curved space*, *Phys. Lett.* **A184** (1994) 409-412.
1166. Hammond R., *Spin, torsion, forces*, *Gen. Relat. Grav.* **26** (1994) 247-263.
1167. Hammond R., *Tetrad formulation of gravity with a torsion potential*, *Gen. Relat. Grav.* **26** (1994) 1107-1116.
1168. Hammond R.T., *Upper limit on the torsion coupling constant*, *Phys. Rev.* **D52** (1995) 6918-6921.
1169. Hammond R.T., *Dirac coupling in gravity with the torsion potential*, *Class. Quantum Grav.* **12** (1995) 279-285.
1170. Hammond R.T., *New fields in general relativity*, *Contemp. Phys.* **36** (1995) 103-114.
1171. Hammond R.T., *Torsion gravity, the Brans - Dicke type generalization and string theory*, *Class. Quantum Grav.* **13** (1996) L73-L79.
1172. Hammond R.T., *Helicity flip cross section from gravity with torsion*, *Class. Quantum Grav.* **13** (1996) 1691-1697.
1173. Hammond R.T., *Gravitation, torsion, and string theory*, *Gen. Relat. Grav.* **28** (1996) 749-757.
1174. Hammond R.T., *Geometric foundation of the magnetic dipole moment*, *Mod. Phys. Lett.* **12** (1997) 2387-2390.
1175. Hammond R.T., *Torsion power*, *Gen. Rel. Grav.* **29** (1997) 727-731.
1176. Hammond R.T., *Spin gravity*, *Gen. Rel. Grav.* **31** (1999) 233-271.
1177. Hammond R.T., *Strings have spin*, *Gen. Rel. Grav.* **32** (2000) 347-351.
1178. Hammond R.T., *Strings in gravity with torsion*, *Gen. Rel. Grav.* **32** (2000) 2007-2019.
1179. Hammond R.T., *String, scalar field, and torsion interactions*, in: "Fluctuating Paths and Fields", *Festschrift Dedicated to Hagen Kleinert on the Occasion of His 60th Birthday*, Eds. W. Janke, A. Pelster, M. Bachmann, and H.-J. Schmidt (World Scientific: Singapore, 2001) 677-683.
1180. Hammond R.T., *Torsion gravity*, *Rep. Prog. Phys.* **65** (2002) 599.
1181. Hammond R.T., *The necessity of torsion in gravity*, *Int. J. Mod. Phys.* **D19** (2010) 2413.
1182. Hammond R.T., *The necessity of torsion in gravity*, *Gen. Rel. Grav.* **42** (2010) 2345-2348.
1183. Hammond R.T., *Spin flip probability of electron due to torsional wave*, *Phys. Rev.* **D90** (2014) 067501 (5 pages).

1184. Hammond R.T., Gruver C., and Kelly P.F., *Scalar field from Dirac coupled torsion*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 445-447.
1185. Hanisch F., Pfäffle F., and Stephan C.A., *The spectral action for Dirac operators with skew-symmetric torsion*, *Commun. Math. Phys.* **300** (2011) 877-888.
1186. Hanson A. and Regge T., *Torsion and quantum gravity*, *Lect. Notes Phys.* **94** (1979) 354-361.
1187. Hari Dass N.D., *A new spin test for equivalence principle*, *Gen. Relat. Grav.* **8** (1977) 89-93.
1188. Hari Dass N.D., *Experimental test for some quantum effects in gravitation*, *Ann.Phys. (NY)* **107** (1977) 337-359.
1189. Harko T., Mak M.K., Lu H.Q., and Cheng K.S., *Bianchi-type I and V cosmologies in Einstein-Cartan theory*, *Nuovo Cim.* **B114** (1999) 1389-1407.
1190. Harko T., Lobo F.S.N., Otalora G., and Saridakis E.N.,  *$f(T, T)$  gravity and cosmology*, *JCAP* **12** (2014) 021.
1191. Harko T., Lobo F.S.N., Otalora G., and Saridakis E.N., *Nonminimal torsion-matter coupling extension of  $f(T)$  gravity*, *Phys. Rev.* **D89** (2014) 124036 (12 pages).
1192. Harnad J.P., and Pettitt R.B., *Gauge theories of space-time symmetries I*, *J. Math. Phys.* **17** (1976) 1827-1837.
1193. Harnett G., *The GHP connection: a metric connection with torsion determined by a pair of null directions*, *Class. Quantum Grav.* **7** (1990) 1681-1705.
1194. Harst U. and Reuter M., *A new functional flow equation for Einstein-Cartan quantum gravity*, *Ann. Phys. (USA)* **354** (2015) 637-704.
1195. Hartley D., *Normal frames for non-Riemannian connections*, *Class. Quantum Grav.* **12** (1995) L103-L105.
1196. Hatzinikitas A., *Locally Weyl invariant massless bosonic and fermionic spin-1/2 action in the  $(Y_4, g)$  space-time*, *Gen. Relat. Grav.* **32** (2000) 2287-2294.
1197. Hashemi M., Jalalzadeh S., and Ziaie A.H., *Collapse and dispersal of a homogeneous spin fluid in Einstein-Cartan theory*, *Eur. Phys. J.* **C75** (2015) 53 (10 pages).
1198. Hayashi K., *Gauge theories of massive and massless tensor fields*, *Progr. Theor. Phys.* **39** (1968) 495-515.
1199. Hayashi K., *The gauge theory of the translation group and underlying geometry*, *Phys. Lett.* **B69** (1977) 441-444.
1200. Hayashi K., *Restrictions on gauge theory of gravitation*, *Phys. Lett.* **B65** (1976) 437-440.
1201. Hayashi K., *Extended translational invariance and associated gauge fields*, *Progr. Theor. Phys.* **38** (1967) 491-507.
1202. Hayashi K., *Minimal and nonminimal gravitational interactions, and the asymmetric energy momentum tensor*, *Gen. Relat. Grav.* **4** (1973) 1-11.
1203. Hayashi K. and Bregman A., *Poincaré gauge invariance and the dynamical role of spin in gravitational theory*, *Ann. Phys. (USA)* **75** (1973) 562-600.
1204. Hayashi K., Kasuya M., and Shirafuji T., *Elementary particles and Weyl's gauge field*, *Progr. Theor. Phys.* **57** (1977) 431-440; Erratum: *Progr. Theor. Phys.* **59** (1978) 681.
1205. Hayashi K., Kasuya M., and Shirafuji T., *Conformal phase group and electromagnetism*, *Progr. Theor. Phys.* **57** (1977) 1781-1794.
1206. Hayashi K. and Kugo T., *Remarks on Weyl's gauge field*, *Progr. Theor. Phys.* **61** (1979) 334-346.

1207. Hayashi K. and Nakano T., *Extended translation invariance and associated gauge fields*, *Progr. Theor. Phys.* **38** (1967) 491-507.
1208. Hayashi K., Nomura K., and Shirafuji T., *Spin precession in space-time with torsion*, *Progr. Theor. Phys.* **84** (1990) 1085-1099.
1209. Hayashi K. and Sasaki R., *Spinor electrodynamics in the Riemann-Cartan space and dynamical theory of axial vector torsion propagating in vacuum*, *Nuovo Cim.* **B45** (1978) 205-228.
1210. Hayashi K. and Shirafuji T., *Gravity, nonintegrable phase factor and motion associated with a closed curve*, *Progr. Theor. Phys.* **58** (1977) 353-361.
1211. Hayashi K. and Shirafuji T., *Spacetime structure explored by elementary particles: microscopic origin for the Riemann-Cartan geometry*, *Progr. Theor. Phys.* **57** (1977) 302-317.
1212. Hayashi K. and Shirafuji T., *New general relativity*, *Phys. Rev.* **D19** (1979) 3524-3553.
1213. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, I. General formulation*, *Progr. Theor. Phys.* **64** (1980) 866-882 [Errata: *Progr. Theor. Phys.* **65** (1981) 2079].
1214. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, II. Equations of motion for test bodies and various limits*, *Progr. Theor. Phys.* **64** (1980) 883-896 [Errata: *Progr. Theor. Phys.* **65** (1981) 2079].
1215. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, III. Weak field approximation*, *Progr. Theor. Phys.* **64** (1980) 1435-1452 [Errata: *Progr. Theor. Phys.* **66** (1981) 741].
1216. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, IV. Mass and energy of particle spectrum*, *Progr. Theor. Phys.* **64** (1980) 2222-2241.
1217. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, V. The extended Bach-Lanczos identity*, *Progr. Theor. Phys.* **65** (1981) 525-532.
1218. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, VI. Scattering amplitudes*, *Progr. Theor. Phys.* **66** (1981) 318-336.
1219. Hayashi K. and Shirafuji T., *Gravity from Poincaré gauge theory of fundamental particles, VII. The axial-vector model*, *Progr. Theor. Phys.* **66** (1981) 2258-2273.
1220. Hayashi K. and Shirafuji T., *Energy, momentum and angular momentum in Poincaré gauge theory*, *Progr. Theor. Phys.* **73** (1985) 54-74.
1221. Hayashi K. and Shirafuji T., *Static, isotropic spacetime in new general relativity*, *Progr. Theor. Phys.* **84** (1990) 36-40.
1222. Hayashi K. and Shirafuji T., *Frame-dragging precession of orbiting gyroscopes in new general relativity and possible violation of equivalence principle*, *Progr. Theor. Phys.* **84** (1990) 1074-1084.
1223. Hayward J., *Internal gauge theory of gravity*, *Phys. Rev.* **D20** (1979) 3039-3046.
1224. Hecht R.D., *On Toma Kerr metric solution in new general relativity*, *Phys. Lett.* **A165** (1992) 194-198.
1225. Hecht R.D., *Mass and spin in Poincaré gauge theory*, *Gen. Rel. Grav.* **27** (1995) 537-554.
1226. Hecht R.D. and Hehl F.W., *A metric affine framework for a gauge theory of gravity*, in: "General relativity and gravitational physics". *Proc. of the 9th Ital. Conf. (Capri, Italy, September 25-28, 1990)*, Eds. R. Cianci, R. de Ritis, G. Marmo, C. Rubano, P. Scudellaro, and M. Francaviglia (World Scientific: Singapore, 1991) 246-291.
1227. Hecht R.D., Hehl F.W., McCrea J.D., Mielke E.W., and Ne'eman Y., *Improved energy-momentum currents in metric-affine spacetimes*, *Phys. Lett.* **A172** (1992) 13-20.

1228. Hecht R.D., Lemke J., and Wallner R.P., *Tachyonic torsional shock waves in Poincaré gauge theory*, *Phys. Lett.* **A151** (1990) 12-14.
1229. Hecht R.D., Lemke J., and Wallner R.P., *Can Poincaré gauge theory be saved?*, *Phys. Rev.* **D44** (1991) 2442-2451.
1230. Hecht R.D. and Nester J.M., *A new evaluation of PGT mass and spin*, *Phys. Lett.* **A180** (1993) 324-331.
1231. Hecht R.D. and Nester J.M., *An evaluation of the mass and spin at null infinity for the PGT and GR gravity theories*, *Phys. Lett.* **A217** (1996) 81-89.
1232. Hecht R.D., Nester J.M., and Zhytnikov V.V., *Some Poincaré gauge theory Lagrangians with well-posed initial value problems*, *Phys. Lett.* **A222** (1996) 37-42.
1233. Heckel B.R., Terrano W.A., and Adelberger E.G., *Limits on exotic long-range spin-spin interactions of electrons*, *Phys. Rev. Lett.* **111** (2013) 151802 (4 pages).
1234. Hehl F.W., *Der Spindrehimpuls in der allgemeinen Relativitätstheorie*, *Abh. Braunsch. Wiss. Ges.* **18** (1966) 98-130.
1235. Hehl F.W., *How does one measure torsion of space-time?*, *Phys. Lett.* **A36** (1971) 225-226.
1236. Hehl F.W., *Spin and torsion in general relativity: I. Foundations*, *Gen. Relat. Grav.* **4** (1973) 333-349.
1237. Hehl F.W., *Spin and torsion in general relativity. II: Geometry and field equations*, *Gen. Relat. Grav.* **5** (1974) 491-516.
1238. Hehl F.W., *On the energy tensor of spinning massive matter in classical field theory and general relativity*, *Repts Math. Phys.* **9** (1976) 55-82.
1239. Hehl F.W., *Fermions and gravity*, in: "Einstein 1978-1955. Colloque du Centenaire" (College de France, 1979) (CNRS: Paris, 1980) 119-148.
1240. Hehl F.W., *Four lectures on Poincaré gauge field theory*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V.De Sabbata (Plenum: New York, 1980) 5-61.
1241. Hehl F.W., *On the kinematics of the torsion of space-time*, *Found. Phys.* **15** (1985) 451-471.
1242. Hehl F.W., *Note on the torsion tensor*, *Physics Today* **60**, n. 3 (2007) 16.
1243. Hehl F.W., *On reference frames in general relativity and in gauge theories of gravity*, in: "Semantical aspects of spacetime theories", Eds. U. Majer and H.J. Schmidt (Bl.-Wiss.-Verl.: Mannheim, 1994)103-112.
1244. Hehl F.W., *An assessment of Evans' unified field theory. I*, *Found. Phys.* **38** (2008) 7-37.
1245. Hehl F.W. and Datta B.K., *Nonlinear spinor equation and asymmetric connection in general relativity*, *J. Math. Phys.* **12** (1971) 1334-1339.
1246. Hehl F.W. and von der Heyde P., *Spin and the structure of space-time*, *Ann. Inst. H.Poincaré* **A19** (1973) 179-196.
1247. Hehl F.W., von der Heyde P., and Kerlick G.D., *General relativity with spin and torsion and its deviations from Einstein's theory*, *Phys. Rev.* **D10** (1974) 1066-1069.
1248. Hehl F.W., von der Heyde P., Kerlick G.D., and Nester J.M., *General relativity with spin and torsion: foundation and prospects*, *Revs. Mod. Phys.* **48** (1976) 393-416.
1249. Hehl F.W. and Kerlick G.D., *Metric-affine variational principles in general relativity. I. Riemannian space-time*, *Gen. Relat. Grav.* **9** (1978) 691-710.
1250. Hehl F.W., Kerlick G.D., and von der Heyde P., *On hypermomentum in general relativity I. The notion of hypermomentum*, *Z. Naturforsch.* **31a** (1976) 111-114.

1251. Hehl F.W., Kerlick G.D., and von der Heyde P., *On hypermomentum in general relativity II. The geometry of spacetime*, *Z. Naturforsch.* **31a** (1976) 524-527.
1252. Hehl F.W., Kerlick G.D., and von der Heyde P., *On hypermomentum in general relativity III. Coupling hypermomentum to geometry*, *Z. Naturforsch.* **31a** (1976) 823-827.
1253. Hehl F.W., Kerlick G.D., and von der Heyde P., *On a new metric affine theory of gravitation*, *Phys. Lett.* **B63** (1976) 446-448.
1254. Hehl F.W., Kerlick G.D., Lord E.A., and Smalley L.L., *Hypermomentum and the microscopic violation of the Riemann-Cartan constraints in general relativity*, *Phys. Lett.* **B70** (1977) 70-72.
1255. Hehl F.W., Kopczyński W., McCrea J.D., and Mielke E.W., *Chern-Simons terms in metric-affine spacetime: Bianchi identities as Euler-Lagrange equations*, *J. Math. Phys.* **32** (1991) 2169-2180.
1256. Hehl F.W. and Kröner E., *Zum Materialgesetz eines elastischen Medium mit Momentenspannungen*, *Z. Naturforsch.* **20a** (1965) 336-350.
1257. Hehl F.W. and Kröner E., *Über den Spin in der allgemeinen Relativitätstheorie: Eine notwendige Erweiterung der Einsteinschen Feldgleichungen*, *Z. Physik* **187** (1965) 478-489.
1258. Hehl F.W., Lemke J., and Mielke E.W., *Two lectures on fermions and gravity*, in: *Geometry and Theoretical Physics (Bad Honeff School, 1990)*, Eds. J. Debrus, and A.C.Hirshfeld (Springer: Berlin, 1992) 56-140.
1259. Hehl F.W., Lord E.A., and Ne'eman Y., *Hadron dilatation, shear and spin as components of the intrinsic hypermomentum current and metric-affine theory of gravitation*, *Phys. Lett.* **B71** (1977) 432-434.
1260. Hehl F.W., Lord E.A., and Ne'eman Y., *Hypermomentum in hadron dynamics and in gravitation*, *Phys. Rev.* **D17** (1978) 428-433.
1261. Hehl F.W., Lord E.A., and Smalley L.L., *Metric affine variational principles in general relativity. II. Relaxation of the Riemann constraint*, *Gen. Relat. Grav.* **13** (1981) 1037-1056.
1262. Hehl F.W. and Macías A., *Metric-affine gauge theory of gravity II. Exact solutions*, *Int. J. Mod. Phys.* **D8** (1999) 399-416.
1263. Hehl F.W. and Mashhoon B., *Nonlocal gravity simulates dark matter*, *Phys. Lett.* **B673** (2009) 279-282.
1264. Hehl F.W. and Mashhoon B., *Formal framework for a nonlocal generalization of Einstein's theory of gravitation*, *Phys. Rev.* **D79** (2009) 064028 (13 pages).
1265. Hehl F.W. and McCrea J.D., *Bianchi identities and the automatic conservation of energy-momentum and angular momentum in general-relativistic field theories*, *Found. Phys.* **16** (1986) 267-293.
1266. Hehl F.W., McCrea J.D., and Mielke E.W., *Weyl spacetimes, the dilaton current, and the creation of gravitating mass by symmetry breaking*, in: *Exact Sciences and their Philosophical Foundations (Vorträge des Internat. Hermann-Weyl-Kongresses, Kiel 1985)*, Eds. W. Deppert, K. Hübner, A. Oberschelp, and V. Weidemann (Verlag Peter Lang: Frankfurt, 1988) 241-310.
1267. Hehl F.W., McCrea J.D., Mielke E.W., and Ne'eman Y., *Progress in metric-affine theories of gravity with local scale invariance*, *Found. Phys.* **19** (1989) 1075-1100.
1268. Hehl F.W., McCrea J.D., Mielke E.W., and Ne'eman Y., *Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilaton invariance*, *Phys. Repts.* **258** (1995) 1-171.
1269. Hehl F.W. and Ne'eman Y., *Spacetime as a continuum with micro-structure and metric-affine gravity*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 31-52.
1270. Hehl F.W., Ne'eman Y., Nitsch J., and von der Heyde P., *Short-range confining component in a quadratic Poincaré gauge theory of gravitation*, *Phys. Lett.* **B78** (1978) 102-106.

1271. Hehl F.W., Nitsch J., and von der Heyde P., *Gravitation and the Poincaré gauge field theory with quadratic Lagrangians*, in: "General Relativity and Gravitation: One Hundred Years after the Birth of Albert Einstein", Ed. A.Held (Plenum: New York, 1980) vol. **1**, 329-355.
1272. Hehl F.W. and Obukhov Yu.N., *Is a 'hadronic' shear current one of the sources in metric-affine gravity?*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 448-450.
1273. Hehl F.W. and Obukhov Yu.N., *Foundations of classical electrodynamics: Charge, flux, and metric* (Birkhäuser: Boston, 2003) 410 p.
1274. Hehl F.W., and Obukhov Yu.N., *Élie Cartan's torsion in geometry and in field theory, an essay*, *Annales de la Fondation Louis de Broglie* **32**, n. 2-3 (2007) 157-194.
1275. Hehl F.W., and Obukhov Yu.N., *An assessment of Evans' unified field theory. II*, *Found. Phys.* **38** (2008) 38-46.
1276. Hehl F.W., Obukhov Yu.N., and Puetzfeld D., *On Poincaré gauge theory of gravity, its equations of motion, and Gravity Probe B*, *Phys. Lett.* **A377** (2013) 1775-1781.
1277. Hehl F.W., Obukhov Yu.N., Rubilar G.F., and Blagojević M., *On the theory of the skewon field: from electrodynamics to gravity*, *Phys. Lett.* **A347** (2005) 14-24.
1278. Hehl F.W. and Šijački D., *Towards a unified gauge theory of gravitational and strong interactions?*, *Gen. Relat. Grav.* **12** (1980) 83-90.
1279. Hehl F.W. and Socorro J., *Gauge theory of gravity: Electrically charged solutions within the metric-affine framework*, *Acta Phys. Pol.* **B29** (1998) 1113-1120.
1280. Heinicke C., *The Einstein 3-form  $G_\alpha$  and its equivalent 1-form  $L_\alpha$  in Riemann-Cartan space*, *Gen. Relat. Grav.* **33** (2001) 1115-1130.
1281. Heinicke C., Baekler P., and Hehl F.W., *Einstein-aether theory, violation of Lorentz invariance, and metric-affine gravity*, *Phys. Rev.* **D72** (2005) 025012 (18 pages).
1282. Helayël-Neto J.A., Penna-Firme A., and Shapiro I.L., *Conformal symmetry, anomaly and effective action for metric-scaler gravity with torsion*, *Phys. Lett.* **B479** (2000) 411-420.
1283. Helayël-Neto J.A., Hernaski C.A., Pereira-Dias B., Vargas-Paredes A.A., and Vasquez-Otoya V.J., *Chern-Simons gravity with (curvature)<sup>2</sup> and (torsion)<sup>2</sup> terms and a basis of degree-of-freedom projection operators*, *Phys. Rev.* **D82** (2010) 064014 (9 pages).
1284. Henning J., and Nitsch J., *Gravity as an internal Yang-Mills gauge field theory of the Poincaré group*, *Gen. Relat. Grav.* **13** (1981) 947-962.
1285. Hernandez E., and Macías A., *Induced masses and spin self-interaction from five-dimensional Einstein-Cartan-Dirac theory*, *Int. J. Mod. Phys.* **A13** (1998) 169-182.
1286. Hernaski C.A., Vargas-Paredes A.A., and Helayël-Neto J.A., *Discussion on massive gravitons and propagating torsion in arbitrary dimensions*, *Phys. Rev.* **D80** (2009) 124012 (9 pages).
1287. Herrmann H., Muschik W., Rückner G., and von Borzeszkowski H.-H., *Spin axioms in different geometries of relativistic continuum physics*, *Found. Phys.* **34** (2004) 1005-1021.
1288. Hestenes D., *Gauge theory gravity with geometric calculus*, *Found. Phys.* **35** (2005) 903-970.
1289. Heyde P. von der, *The field equations of the Poincaré gauge theory of gravitation*, *Phys. Lett.* **A58** (1976) 141-143.
1290. Heyde P. von der, *Is gravitation mediated by the torsion of space-time?*, *Z. Naturforsch.* **31a** (1976) 1725-1726.
1291. Heyde P. von der, *The equivalence principle in the  $U_4$  theory of gravitation*, *Lett. Nuovo Cim.* **14** (1975) 250-252.

1292. Heyde P. von der, *A generalized Lovelock theorem for the gravitational field with torsion*, *Phys. Lett.* **A51** (1975) 381-382.
1293. Hirashima H., *A new gravitational energy-momentum pseudotensor in tetrad form*, *Progr. Theor. Phys.* **96** (1996) 347-375.
1294. Hlavatý V., *Geometry of Einstein's unified field theory* (Noordhoff: Groningen, 1957) 341 pp.
1295. Ho J.K., Chern D.C., and Nester J.M., *Some spherically symmetric exact solutions of the metric-affine gravity theory*, *Chinese J. Phys.* **35** (1997) 640-650.
1296. Ho J.K. and Nester J.M., *Poincaré gauge theory with even and odd parity dynamic connection modes: isotropic Bianchi cosmological models*, *J. Phys.: Conf. Ser.* **330** (2011) 012005.
1297. Ho J.K. and Nester J.M., *Poincaré gauge theory with coupled even and odd parity spin-0 modes: cosmological normal modes*, *Ann. d. Physik* **524** (2012) 97-106.
1298. Ho F.H. and Nester J.M., *Poincaré gauge theory with coupled even and odd parity dynamic spin-0 modes: dynamical equations for isotropic Bianchi cosmologies*, *Int. J. Mod. Phys.* **D20** (2011) 2125.
1299. Ho P.-M., *Generalized Yang-Mills theory and gravity*, *Phys. Rev.* **D93** (2016) 044062 (13 pages).
1300. Hochberg D. and Visser M., *Null energy condition in dynamic wormholes*, *Phys. Rev. Lett.* **81** (1998) 746-749.
1301. Hochberg D. and Visser M., *Dynamic wormholes, antitrapped surfaces, and energy conditions*, *Phys. Rev.* **D58** (1998) 044021 (14 pages).
1302. Hoff da Silva J.M. and Rocha R. da, *Braneworld remarks in Riemann-Cartan manifolds*, *Class. Quantum Grav.* **26** (2009) 055007.
1303. Hoff da Silva J.M., and Rocha R. da, *Reply to 'comment on 'Braneworld remarks in Riemann-Cartan manifolds''*, *Class. Quantum Grav.* **26** (2009) 178002.
1304. Hohmann M., *Extensions of Lorentzian spacetime geometry: From Finsler to Cartan and vice versa*, *Phys. Rev.* **D87** (2013) 124034 (16 pages).
1305. Hohmann M., *Symmetry in Cartan language for geometric theories of gravity*, in: "Gravitation, Astrophysics, and Cosmology", *Proc. Twelfth Asia-Pacific Internat. Conf., Moscow, 28 June–5 July 2015*, Eds. V. Melnikov and J.-P. Hsu (World Scientific: Singapore, 2016) p. 257-258.
1306. Hohmann M., *Spacetime and observer space symmetries in the language of Cartan geometry*, *J. Math. Phys.* **57** (2016) 082502 (28 pages).
1307. Hojman S., *Lagrangian theory of the motion of spinning particles in torsion gravitational theories*, *Phys. Rev.* **D18** (1978) 2741-2744.
1308. Hojman R., Mukku C., and Sayed W.A., *Parity violation in metric-torsion theories of gravitation*, *Phys. Rev.* **D22** (1980) 1915-1921.
1309. Hojman S., Rosenbaum M., Ryan M.P., and Shepley L.C., *Gauge invariance, minimal coupling, and torsion*, *Phys. Rev.* **D17** (1978) 3141-3146.
1310. Hojman S., Rosenbaum M., and Ryan M.P., Jr., *Propagating torsion and gravitation*, *Phys. Rev.* **D19** (1979) 430-437.
1311. Hojman S., Rosenbaum M., and Ryan M.P., Jr., *A torsion-generated unified theory of the gravitational and real-massless-scalar field*, in: "Relativity and Gravitation", *Proc. of the 3rd Latin-American Symp. SILARG III*, Eds. S. Hojman, M. Rosenbaum, and M.P. Ryan, Jr. (UNAM: Mexico, 1982) 183-195.
1312. Holten J.W. van, Waldron A., and Peeters K., *An index theorem for non-standard Dirac operators*, *Class. Quantum Grav.* **16** (1999) 2537-2544.
1313. Holz A., *Geometry and action of arrays of disclinations in crystals and relation to (2+1)-dimensional gravitation*, *Class. Quant. Grav.* **5** (1988) 1259-1282.

1314. Horak M. and Krupka D., *First-order invariant Einstein-Cartan variational structures*, *Int. J. Theor. Phys.* **17** (1978) 573-584.
1315. Horvat D., Ilijić S., Kirin A., and Narančić Z., *Nonminimally coupled scalar field in teleparallel gravity: boson stars*, *Class. Quantum Grav.* **32** (2015) 035023.
1316. Horvat D., Ilijić S., Kirin A., and Narančić Z., *Note on the charged boson stars with torsion-coupled field*, *Phys. Rev.* **D92** (2015) 024045 (7 pages).
1317. Hourī T., Kubizňák D., Warnick C.M., and Yasui Y., *Symmetries of the Dirac operator with skew-symmetric torsion*, *Class. Quantum Grav.* **27** (2009) 185019.
1318. Hourī T., Kubizňák D., Warnick C.M., and Yasui Y., *Local metrics admitting a principal Killing-Yano tensor with torsion*, *Class. Quantum Grav.* **29** (2012) 165001.
1319. Hsu J.P., *De Sitter and Poincaré gauge invariant fermion Lagrangians and gravity*, *Phys. Lett.* **B119** (1982) 328-330.
1320. Hsu J.P. and Fine D. (Eds), “*100 Years of Gravity and Accelerated Frames: The Deepest Insights of Einstein and Yang-Mills*”, Advanced Series on Theoretical Physical Science: Vol. 9 (World Scientific: Singapore, 2005) 664 pp.
1321. Hsu R.-R. and Yeung W.-B., *A Wu-Yang monopole in an asymptotically flat Riemann-Cartan spacetime*, *Phys. Lett.* **B151** (1985) 240-242.
1322. Hsu R.-R. and Yeung W.-B., *Kaluza-Klein unification in the Poincaré gauge theory of gravitation*, *Phys. Lett.* **B155** (1985) 143-146.
1323. Hsu R.-R. and Yeung W.-B., *Kaluza-Klein electric monopole in a six-dimensional Poincaré gauge theory of gravitation*, *Phys. Lett.* **B155** (1985) 147-150.
1324. Hsu R.-R. and Yeung W.-B., *Electromagnetic waves as geometrical ripples in a higher dimensional space-time with torsion*, *Class. Quantum Grav.* **4** (1987) 753-757.
1325. Huang C.-G. and Ma M.-S., *de Sitter spacetimes with torsion in the model of de Sitter gauge theory of gravity*, *Phys. Rev.* **D80** (2009) 084033 (9 pages).
1326. Huang C.-G. and Ma M.-S., *On torsion-free vacuum solutions of the model of de Sitter gauge theory of gravity (II)*, *Frontiers of Physics in China* **4** (2009) 525-529.
1327. Huang C.-G. and Ma M.-S., *A new solution with torsion in model of dS gauge theory of gravity*, *Commun. Theor. Phys. (China)* **55** (2013) 65-68.
1328. Huang C.-G., Zhang H.-Q., and Guo H.-Y., *Cosmological solutions with torsion in a model of the de Sitter gauge theory of gravity*, *JCAP* **10** (2008) 010.
1329. Huang C.-G., Zhang H.-Q., and Guo H.-Y., *A new gravitational model for dark energy*, *Chinese Phys.* **C32** (2008) 687.
1330. Huang P. and Yuan F.-F., *Generalized Hodge dual for torsion in teleparallel gravity*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650106 (11 pages).
1331. Huang Q.-H., Wu P.-X., and Yu H.-W., *Emergent scenario in the Einstein-Cartan theory*, *Phys. Rev.* **D91** (2015) 103502 (8 pages).
1332. Huang Y.C. and Lin B.L., *General Weitzenböck theory of crystals with different dislocation distributions*, *Phys. Lett.* **A299** (2002) 644-649.
1333. Huang Z.-G., Fang W., and Lu H.-Q., *Inflation and singularity in Einstein-Cartan theory*, *Chinese Phys. Lett.* **28** (2011) 029802.
1334. Huang Z.-G., Fang W., and Lu H.-Q., *Inflation and singularity of a Bianchi type-VIII universe with a Dirac field in the EinsteinCartan theory*, *Chinese Phys. Lett.* **28** (2011) 089801.
1335. Hughes T.L., Leigh R.G., and Parrikar O., *Torsional anomalies, Hall viscosity, and bulk-boundary correspondence in topological states*, *Phys. Rev.* **D88** (2013) 025040 (30 pages).

1336. Hunter L.R. and Ang D.G., *Using geoelectrons to search for velocity-dependent spin-spin interactions*, *Phys. Rev. Lett.* **112** (2014) 091803 (5 pages).
1337. Hurth T., Nieuwenhuizen P. van, Waldron A., and Preitschopf C., *Possible new  $R^2$  theory of supergravity*, *Phys. Rev.* **D55** (1997) 7593-7614.

## I

1338. Ikeda N., *2-Dimensional gravity and nonlinear gauge theory*, *Ann. Phys.* **235** (1994) 435-464.
1339. Ikeda N. and Izawa K.-I., *Quantum gravity with dynamical torsion in two dimensions*, *Progr. Theor. Phys.* **89** (1993) 223-230.
1340. Ikeda N. and Izawa K.-I., *Gauge theory based on quadratic Lie algebras and 2D gravity with dynamical torsion*, *Progr. Theor. Phys.* **89** (1993) 1077-1085.
1341. Ikeda S., *Some physico-geometrical remarks on the relationship between the micro- and macro-gravitational fields*, *Lett. Nuovo Cim.* **25** (1979) 21-25.
1342. Ikeda S., *A thermodynamical aspect underlying the concept of torsion*, *Progr. Theor. Phys.* **64** (1980) 2265-2266.
1343. Iliev B.Z., *Deviation equation and equivalence principle in space with affine connection*, *Dokl. Bulgar. Akad. Nauk* **37** (1984) 1187-1190.
1344. Iliev B.Z., *Normal frames and the validity of the equivalence principle: I. Cases in a neighbourhood and at a point*, *J. Phys. A: Math. and Gen.* **A29** (1996) 6895-6901.
1345. Iliev B.Z., *Normal frames and the validity of the equivalence principle: II. The case along paths*, *J. Phys. A: Math. and Gen.* **A30** (1997) 4327-4336.
1346. Iliev B.Z., *Normal frames and the validity of the equivalence principle: III. The case along smooth maps with separable points of self-intersection*, *J. Phys. A: Math. and Gen.* **A31** (1998) 1287-1296.
1347. Iliev B.Z., *Is the principle of equivalence a principle?*, *J. Geom. Phys.* **24** (1998) 209-222.
1348. Iliev B.Z., *Normal frames for derivations and linear connections and the equivalence principle*, *J. Geom. Phys.* **45** (2003) 24-53.
1349. Iliev B.Z., *Normal frames for general connections on differentiable fibre bundles*, *J. Geom. Phys.* **56** (2006) 780-812.
1350. Illge R., *Lagrange-Funktion für massive teilchen mit ganzzahligem Spin in beliebig gekrümmten Raum-Zeiten*, *Experimentelle Technik der Physik* **34** (1986) 429-432.
1351. Infeld L. and van der Waerden B.L., *Die Wellengleichungen des Elektrons in der allgemeinen Relativitätstheorie*, *Sitzungs. Akad. Berlin* (1933) 380-403.
1352. Inomata A., *Effect of the selfinduced torsion of the Dirac sources on gravitational singularities*, *Phys. Rev.* **D18** (1976) 3552-3556.
1353. Inomata A. and Trinkala M., *General relativity as a limit of the de Sitter gauge theory*, *Phys. Rev.* **D19** (1979) 1665-1668.
1354. Iochum B., Levy C., and Vassilevich D., *Spectral action for torsion with and without boundaries*, *Commun. Math. Phys.* **310** (2012) 367-382.
1355. Iochum B. and Schücker T., *Diffeomorphisms and orthonormal frames*, *J. Geom. Phys.* **56** (2006) 1496-1505.
1356. Iorio L., *Modified theories of gravity with nonminimal coupling and orbital particle dynamics*, *Class. Quantum Grav.* **31** (2014) 085003.

1357. Isham C.J., Salam A., and Strathdee J., *Nonlinear realizations of space-time symmetries. Scalar and tensor gravity*, *Ann. Phys. (USA)* **62** (1971) 98-119.
1358. Israelit M., *Massive electrodynamics and magnetic monopoles*, *Gen. Rel. Grav.* **29** (1997) 1411-1424.
1359. Israelit M., *Torsional Weyl-Dirac electrodynamics*, *Found. Phys.* **28** (1998) 205-229.
1360. Israelit M., *On measuring standards in Weyl's geometry*, *Found. Phys.* **35** (2005) 1769-1782.
1361. Itin Y., *Gravity on a parallelizable manifold. Exact solutions*, *Gen. Rel. Grav.* **31** (1999) 187-203.
1362. Itin Y., *A class of quasi-linear equations in coframe gravity*, *Gen. Rel. Grav.* **31** (1999) 1891-1920.
1363. Itin Y., *Coframe teleparallel models of gravity: exact solutions*, *Int. J. Mod. Phys.* **D10** (2001) 547-573.
1364. Itin Y., *Energy-momentum current for coframe gravity*, *Class. Quantum Grav.* **19** (2002) 173.
1365. Itin Y. and Hehl F.W., *Maxwell's field coupled nonminimally to quadratic torsion: Axion and birefringence*, *Phys. Rev.* **D68** (2003) 127701 (4 pages).
1366. Itin Y., Hehl F.W., and Obukhov Yu.N., *Premetric equivalent of general relativity: Teleparallelism*, *Phys. Rev.* **D95** (2017) 084020 (13 pages).
1367. Itin Y. and Kaniel S., *On a class of invariant coframe operators with application to gravity*, *J. Math. Phys.* **41** (2000) 6318-6340.
1368. Ivanenko D., *Tetradic and compensational theory of gravitation*, *Compt. Rendus de l'Acad. Bulgare des Sciences* **17** (1964) 801-804.
1369. Ivanenko D., *Gravitation and unified picture of matter*, in: "Atti Conv. sulla Relat. Gen.: Probl. Energ. Onde Grav." (Barbera Editore: Firenze, 1965) 3-19.
1370. Ivanenko D., *A program of unified theory*, *Tensor N.S.* **25** (1972) 161-170.
1371. Ivanenko D., *A new attempt at a unified field theory*, *Acta Physica Hungarica* **32** (1973) 341-352.
1372. Ivanenko D., *The cosmological term, compensation, and singularities*, *Sov. J. Phys.* **17** (1974) 1661-1666 [*Izvestia VUZov, Fizika*, n. 12 (1974) 35-42 (in Russian)].
1373. Ivanenko D., *Connection between cosmology and the micro-cosmos*, *Sov. J. Phys.* **19** (1976) 37-39 [*Izvestia VUZov, Fizika*, n. 1 (1976) 49-51 (in Russian)].
1374. Ivanenko D., *On the extensions of general relativity*, in: "Old and New Questions in Physics, Cosmology, Philosophy and Theoret. Biology", Ed. A. van der Merwe (Plenum: New York, 1983) 213-221.
1375. Ivanenko D.D., *Gravitation and elementary particles*, in: "Relativistic Astrophysics and Cosmology. Proc. of the Sir A.Eddington Centenary Symp.", Eds. V.DeSabbata and T.M.Karade (World Scientific: Singapore, 1984) vol. 1, 137-169.
1376. Ivanenko D., Budylin S., and Pronin P., *Stochastic quantization of Einstein-Cartan theory*, *Ann. der Phys.* **45** (1988) 191-199.
1377. Ivanenko D. and Frolov B.N., *Unified description of isotopic and spacetime properties of elementary particles*, *Vestnik Mosk. Univ. Fizika*, n. 3 (1967) 106-107.
1378. Ivanenko D.D. and Krechet V.G., *Hierarchical models of unified theories*, *Sov. J. Phys.* **23** (1980) 985-987 [*Izvestia VUZov, Fizika*, n. 12 (1987) 3-6 (in Russian)].
1379. Ivanenko D.D. and Krechet V.G., *Rotation of universe*, *Sov. J. Phys.* **30** (1987) 191-194 [*Izvestia VUZov, Fizika*, n. 3 (1987) 12-16 (in Russian)].

1380. Ivanenko D.D. and Obukhov Yu. N., *Gravitational interaction of fermion antisymmetric tensor fields*, *Ann. d. Phys.* **42** (1985) 59-70.
1381. Ivanenko D.D. and Obukhov Yu.N., *Cosmological model of an expanding and rotating universe*, in: "Perspectives of the unified theory", Eds. D.V. Gal'tsov, L.S. Kuz'menkov and P.I. Pronin (Moscow Univ. Publ. House: Moscow, 1991) 98-115 (in Russian).
1382. Ivanenko D.D., Pronin P.I., and Sardanashvily G.A., *Gauge theory of gravity* (Moscow University Publ. House: Moscow, 1985) 144 pp. (in Russian).
1383. Ivanenko D. and Sardanashvily G., *New aspects of compensation theory*, in: "Actual Problems of Theor. Physics. Prof. D.Ivanenko's Festschrift", Ed. A.A. Sokolov (Moscow Univ. Publ. House: Moscow, 1976) 97-116 (in Russian).
1384. Ivanenko D.D. and Sardanashvily G.A., *Extensions of Einstein theory of gravitation and perspectives of unified field theories*, *Sov. J. Phys.* **23** (1980) 118-128 [*Izvestia VUZov, Fizika*, n. 2 (1980) 54-66 (in Russian)].
1385. Ivanenko D. and Sardanashvily G., *On the relativity and equivalence principles in the gauge theory of gravitation*, *Lett. Nuovo Cim.* **30** (1981) 220-222.
1386. Ivanenko D. and Sardanashvily G., *Relativity principle and equivalence principles in the gauge gravitation theory*, *Comptes Rendus de l'Academie Bulgare des Sciences* **34** (1981) 1237-1239.
1387. Ivanenko D. and Sardanashvily G., *Relativity and equivalence principles in the gauge theory of gravity*, *Sov. J. Phys.* **24** (1981) 555-557 [*Izvestia VUZov, Fizika*, n. 6 (1981) 79-82 (in Russian)].
1388. Ivanenko D. and Sardanashvily G., *The gauge treatment of gravity*, *Phys. Repts.* **94** (1983) 3-45.
1389. Ivanenko D. and Sardanashvily G., *Goldstone type (non-Poincaré) supergravity*, *Progr. Theor. Phys.* **75** (1986) 969-976.
1390. Ivanenko D. and Sardanashvily G., *Goldstone theory of gravity*, in: "Problems of Gravity", Ed. D.V. Gal'tsov (Moscow Univ. Publ. House: Moscow, 1986) 112-127 (in Russian).
1391. Ivanenko D. and Sardanashvily G., *On the Goldstone gravitation theory*, *Pramana J. Phys.* **29** (1987) 21-37.
1392. Ivanov A.N., Höllwieser R., Jenke T., Wellenzohn M., and Abele H., *Influence of the chameleon field potential on transition frequencies of gravitationally bound quantum states of ultracold neutrons*, *Phys. Rev.* **D87** (2013) 105013 (17 pages).
1393. Ivanov A.N. and Pitschmann M., *Nonrelativistic approximation of the Dirac equation for slow fermions in static metric spacetimes*, *Phys. Rev.* **D90** (2014) 045040 (8 pages).
1394. Ivanov A.N. and Snow W.M., *Parity-even and time-reversal-odd neutron optical potential in spinning matter induced by gravitational torsion*, *Phys. Lett.* **B764** (2017) 186-189.
1395. Ivanov A.N. and Wellenzohn M., *Standard electroweak interactions in gravitational theory with chameleon field and torsion*, *Phys. Rev.* **D91** (2015) 085025 (24 pages).
1396. Ivanov A.N. and Wellenzohn M., *Nonrelativistic approximation of the Dirac equation for slow fermions coupled to the chameleon and torsion fields in the gravitational field of the Earth*, *Phys. Rev.* **D92** (2015) 065006 (12 pages).
1397. Ivanov A.N. and Wellenzohn M., *Effective low-energy potential for slow Dirac fermions in Einstein-Cartan gravity with torsion and chameleon field*, *Phys. Rev.* **D92** (2015) 125004 (16 pages).
1398. Ivanov A.N. and Wellenzohn M., *Spin precession of slow neutrons in Einstein-Cartan gravity with torsion, chameleon, and magnetic field*, *Phys. Rev.* **D93** (2016) 045031 (13 pages).
1399. Ivanov A.N. and Wellenzohn M., *Einstein-Cartan gravity with torsion field serving as an origin for the cosmological constant or dark energy density*, *Astrophys. J.* **829** (2016) 47 (9 pages).

1400. Ivanov E.A. and Niederle J., *Gauge formulation of gravitational theories I. The Poincaré, de Sitter and conformal cases*, *Phys. Rev.* **D25** (1982) 976-986.
1401. Ivanov E.A. and Niederle J., *Gauge formulation of gravitational theories II. The special conformal case*, *Phys. Rev.* **D25** (1982) 987-994.
1402. Ivanov G.G., *Kinetic equation for a system of supersymmetric Dirac particles in electromagnetic and gravitational fields*, *Sov. J. Phys.* **28** (1985) 13-17 [*Izvestia VUZov, Fizika*, n. 1 (1985) 16-21 (in Russian)].
1403. Izumi K. and Ong Y.-C., *Cosmological perturbation in  $f(T)$  gravity revisited*, *JCAP* **06** (2013) 029.

## J

1404. Jamil M., Momeni D., and Myrzakulov R., *Energy conditions in generalized teleparallel gravity models*, *Gen. Rel. Grav.* **45** (2013) 263-273.
1405. Järv L. and Toporensky A., *General relativity as an attractor for scalar-torsion cosmology*, *Phys. Rev.* **D93** (2016) 024051 (8 pages).
1406. Jennen H., *Cartan geometry of spacetimes with a nonconstant cosmological function  $\Lambda$* , *Phys. Rev.* **D90** (2014) 084046 (6 pages).
1407. Jha R., *A parity-violating  $U_4$ -gravitation theory*, *Gen. Relat. Grav.* **25** (1993) 281-290.
1408. Jha R., *A new theory of gravitation with parity violating propagating torsion*, *Int. J. Mod. Phys.* **A9** (1994) 3959-3968.
1409. Jha R., Lord E.A., and Sinha K.P., *A singular-free cosmological model in Einstein-Cartan-Sciama-Kibble theory*, *Gen. Relat. Grav.* **20** (1988) 565-571.
1410. Jiménez J.B., Golovnev A., Karčiauskas M., and Koivisto T.S., *Bimetric variational principle for general relativity*, *Phys. Rev.* **D86** (2012) 084024 (8 pages).
1411. Joglekar S. and Griffiths J.B., *A Newman-Penrose-type formalism for space-times with torsion*, *Gen. Relat. Grav.* **12** (1980) 597-617.
1412. Juan F. de, Cortijo A., and Vozmediano M.A.H., *Dislocations and torsion in graphene and related systems*, *Nucl. Phys.* **B828** (2010) 625-637.
1413. Julia B. and Silva S., *Currents and superpotentials in classical gauge theories: I. Local results with applications to perfect fluids and general relativity*, *Class. Quantum Grav.* **15** (1998) 2173-2215.
1414. Julia B. and Silva S., *Currents and superpotentials in classical gauge theories: II. Global aspects and the example of affine gravity*, *Class. Quantum Grav.* **17** (2000) 4733-4743.
1415. Julve J., López-Pinto A., Tiemblo A., and Tresguerres R., *Nonlinear 4D conformal gauge spacetime symmetry*, *Class. Quantum Grav.* **12** (1995) 1327-1336.
1416. Julve J., López-Pinto A., Tiemblo A., and Tresguerres R., *Nonlinear local dynamics and space-time symmetries*, *An. Fis. (Spain)* **91** (1995) 89-95.
1417. Julve J., López-Pinto A., Tiemblo A., and Tresguerres R., *Nonlinear gauge foundation of affine theories of gravity with fermionic matter*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 115-144.
1418. Julve J., López-Pinto A., Tiemblo A., and Tresguerres R., *Nonlinear gauge realization of space-time symmetries including translations*, *Gen. Relat. Grav.* **28** (1996) 759-768.
1419. Junior E.L.B., Rodrigues M.E., Salako I.G., and Houndjo M.J.S., *Reconstruction, thermodynamics and stability of the CDM model in  $f(T, \mathcal{T})$  gravity*, *Class. Quantum Grav.* **33** (2016) 125006 (21 pages).

*K*

1420. Kadic A. and Edelen D.G.B., *A gauge theory of dislocations and disclinations*, Lect. Notes Phys. **174** (Springer: Berlin, 1983) 290 pp.
1421. Kaempfer F.A., *On a possible unification of gravitational and weak interactions*, Gen. Relat. Grav. **7** (1976) 327-331.
1422. Kaempfer F.A., *Self-consistent torsion potentials*, Phys. Rev. **D18** (1978) 2727-2729.
1423. Kaempfer B., *GL(4,R) version of gauge theory of gravitation*, Acta. Phys. Pol. **B12** (1981) 419-428.
1424. Kaku M., Townsend P.K., and van Nieuwenhuizen P., *Gauge theory of conformal and superconformal group*, Phys. Lett. **B69** (1977) 304-308.
1425. Kalinowski M.W., *On generalization of the Einstein-Cartan theory and the Klein-Kaluza theory*, Lett. Math. Phys. **5** (1981) 489-500.
1426. Kalinowski M.W., *An Einstein-Cartan-Moffat theory*, Phys. Rev. **D26** (1982) 3419-3424.
1427. Kalinowski M.W., *Torsion and the Kaluza-Klein theory*, Acta Phys. Austr. **57** (1985) 45-63.
1428. Kalinowski M.W., *Minimal coupling scheme for Dirac's field in the non-symmetric theory of gravitation*, Int. J. Mod. Phys. **A1** (1986) 227-242.
1429. Kalinowski M.W., *On some developments in the nonsymmetric Kaluza-Klein theory*, Eur. Phys. J. **C74** (2014) 2742 (33 pages).
1430. Kalmykov M.Yu., *Affine-metric quantum gravity with extra local symmetries*, Class. Quantum Grav. **14** (1997) 367-378.
1431. Kalmykov M.Yu. and Pronin P.I., *One-loop effective action in gauge gravitational theory*, Nuovo Cim. **B106** (1991) 1401-1415.
1432. Kalmykov M.Yu. and Pronin P.I., *One-loop effective action in gauge gravitational theory*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 53-74.
1433. Kalmykov M.Yu. and Pronin P.I., *One-loop divergences of linear gravity with torsion terms*, Gen. Relat. Grav. **27** (1995) 873-886.
1434. Kalmykov M.Yu. and Pronin P.I., *Affine-metric gravity with the projective invariance*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 145-183.
1435. Kalmykov M.Yu. and Pronin P.I., *The one-loop divergences of the linear gravity with the torsion terms in tetrad approach*, Mod. Phys. Lett. **A13** (1998) 2827-2837.
1436. Kalmykov M.Yu., Pronin P.I., and Stepanyantz K.V., *Projective invariance and one-loop effective action in affine-metric gravity interacting with scalar field*, Class. Quantum Grav. **11** (1994) 2645-2652.
1437. Kalyanshetti S.B., and Waghmode B.B., *A static cosmological model in Einstein-Cartan theory*, Gen. Relat. Grav. **14** (1982) 821-830.
1438. Kalyanshetti S.B. and Waghmode B.B., *Analog of Birkhoff's theorem in Einstein-Cartan theory for Zeldovich fluid*, Indian J. Pure Ap. Mat. **13** (1982) 1208-1212.
1439. Kalyanshetti S.B. and Waghmode B.B., *Conformally flat static spherically symmetric perfect fluid distribution in Einstein-Cartan theory*, Phys. Rev. **D27** (1983) 2835-2838.
1440. Kaniel S. and Itin Y., *Gravity on a parallelizable manifold*, Nuovo Cim. **B113** (1998) 393-400.
1441. Kánnár J., *A note on the strengths of singularities in the Einstein-Cartan theory*, Gen. Relat. Grav. **27** (1995) 23-33.

1442. Kanno S., *Interaction between fermions and gravitational field in the very early universe*, *Progr. Theor. Phys.* **79** (1988) 1365-1377.
1443. Kao W.F., *Scale-invariant effective action with torsion*, *Can. J. Phys.* **70** (1992) 238-241.
1444. Kao W.F., *Spatial dependence of inflaton in the presence of the torsion field*, *Phys. Lett.* **A163** (1992) 155-158.
1445. Kao W.F., *Torsion, compactification, and inflation*, *Phys. Rev.* **D47** (1993) 3639-3642.
1446. Karahan C.N., Altaş A., and Demir D.A., *Scalars, vectors and tensors from metric-affine gravity*, *Gen. Rel. Grav.* **45** (2013) 319-343.
1447. Karahan C.N., Dogangün, O., and Demir D.A., *Conformal transformations in metric-affine gravity and ghosts*, *Ann. Phys. (Berlin)* **524** (2012) 461-469.
1448. Karami K. and Abdolmaleki A., *Generalized second law of thermodynamics in  $f(T)$  gravity*, *JCAP* **04** (2012) 007.
1449. Karananas G.K., *The particle spectrum of parity-violating Poincaré gravitational theory*, *Class. Quantum Grav.* **32** (2015) 055012 (38 pages).
1450. Karananas G.K., *Corrigendum: The particle spectrum of parity-violating Poincaré gravitational theory*, *Class. Quantum Grav.* **32** (2015) 089501 (2 pages).
1451. Karbanovski V.V., Maxioma R.I., Bogdanov M.S., Nefiodova O.V., Kolesnikova Yu.V., and Panait E.O., *On status of constant  $f$  in the conformal Poincaré gauge theory of gravitation*, *Acta Phys. Pol.* **B36** (2005) 2143-2147.
1452. Karbanovski V.V., Nefiodova O.V., Kolesnikova Yu.V., Limonnikova Yu.N., and Sorokin A.S., *Study of cosmological model Bianchi I in the conformal Poincaré-gauge theory of gravitation*, *Acta Phys. Pol.* **B38** (2007) 1979-1983.
1453. Kassandrov V.V. and Rizcallah J.A., *Regularized Lienard-Wiechert fields in a space with torsion*, *Gravitation and Cosmology* **21** (2015) 273-278.
1454. Kassandrov V.V. and Rizcallah J.A., *Regular charged formation on an arbitrary worldline in a space with torsion*, in: "Gravitation, Astrophysics, and Cosmology", *Proc. Twelfth Asia-Pacific Internat. Conf., Moscow, 28 Jun - 5 July 2015*, Eds. V. Melnikov and J.-P. Hsu (World Scientific: Singapore, 2016) p. 204-207.
1455. Kasuya M., *On the gauge theory in the Einstein-Cartan-Weyl space-time*, *Nuovo Cim.* **B28** (1975) 127-137.
1456. Kasuya M., *The Einstein-Cartan theory of gravitation in a Hamiltonian form*, *Progr. Theor. Phys.* **60** (1978) 167-177.
1457. Kasuya M., *On quantization of Einstein-Cartan theory with anticommuting variables in Dirac's Hamiltonian formalism*, *Nuovo Cim.* **B52** (1979) 244-250.
1458. Katanaev M.O., *On the gauge theory for the Poincaré group*, *Theor. Math. Phys.* **54** (1983) 248-252 [*Teor. Mat. Fiz.* **54** (1983) 381-387 (in Russian)].
1459. Katanaev M.O., *Linear connection in a theory of kaluza-Klein type*, *Theor. Math. Phys.* **56** (1983) 795-798 [*Teor. Mat. Fiz.* **56** (1983) 246-250 (in Russian)].
1460. Katanaev M.O., *Kinetic term for Lorentz connection*, *Theor. Math. Phys.* **65** (1985) 1043-1050 [*Teor. Mat. Fiz.* **65** (1985) 108-118 (in Russian)].
1461. Katanaev M.O., *Dynamical torsion and indefinite metric problem in general relativity*, *Yad. Fizika* **43** (1985) 772-774 (in Russian).
1462. Katanaev M.O., *Kinematic part in the dynamical torsion theory*, *Theor. Math. Phys.* **72** (1987) 735-741 [*Theor. Math. Phys.* **72** (1987) 79-88 (in Russian)].
1463. Katanaev M.O., *Large-scale limit of dynamic torsion theory*, *Sov. J. Phys.* **30** (1987) 392-396 [*Izvestiya VUZov, Fizika* **n 5** (1987) 39-44 (in Russian)].

1464. Katanaev M.O., *A new integrable model: two-dimensional gravity with dynamical torsion*, *Sov. Phys. Dokl.* **34** (1989) 982-983 [*Dokl. AN SSSR* **309** (1989) 591-593 (in Russian)].
1465. Katanaev M.O., *String with dynamical geometry: Hamiltonian analysis in conformal gauge*, *Theor. Math. Phys.* **80** (1989) 838-848 [*Theor. Math. Phys.* **80** (1989) 239-252 (in Russian)].
1466. Katanaev M.O., *Complete integrability of two dimensional gravity with dynamical torsion*, *J. Math. Phys.* **31** (1990) 882-891.
1467. Katanaev M.O., *Conformal invariance, extremals and geodesics in two-dimensional gravitation with torsion*, *J. Math. Phys.* **32** (1991) 2483-2496.
1468. Katanaev M.O., *All universal coverings of two-dimensional gravity with torsion*, *J. Math. Phys.* **34** (1993) 700-736.
1469. Katanaev M.O., *New constraints in dynamical torsion theory*, *Gen. Relat. Grav.* **25** (1993) 349-359.
1470. Katanaev M.O., *Canonical quantization of the string with dynamical geometry and anomaly free nontrivial string in two dimensions*, *Nucl. Phys.* **B416** (1994) 563-605.
1471. Katanaev M.O., *Euclidean two-dimensional gravity with torsion*, *J. Math. Phys.* **38** (1997) 946-980.
1472. Katanaev M.O., *Global solutions in gravity*, *Nucl. Phys. B, Proc. Suppl.* **88** (2000) 233-236.
1473. Katanaev M.O., *Rotational elastic waves in double wall tube*, *Phys. Lett.* **A379** (2015) 1544-1548.
1474. Katanaev M.O., Klosch T., and Kummer W., *Global properties of warped solutions in general relativity*, *Ann. Phys. (NY)* **276** (1999) 191-222.
1475. Katanaev M.O., Kummer W., and Liebl H., *Geometric interpretation and classification of global solutions in generalized dilaton gravity*, *Phys. Rev.* **D53** (1996) 5609-5618.
1476. Katanaev M.O. and Volovich I.V., *Higgs fields in Kaluza-Klein theory with dynamical torsion*, *Phys. Lett.* **B156** (1985) 327-330.
1477. Katanaev M.O. and Volovich I.V., *Two-dimensional gravity with dynamical torsion and strings*, *Ann. Phys. (USA)* **197** (1990) 1-32.
1478. Katanaev M.O. and Volovich I.V., *Theory of defects in solids and three-dimensional gravity*, *Ann. Phys. (USA)* **216** (1992) 1-28.
1479. Katanaev M.O. and Volovich I.V., *Scattering on dislocations and cosmic string in the geometric theory of defects*, *Ann. Phys. (USA)* **216** (1999) 203-232.
1480. Katkar L.N., *An exact spherically symmetric space-time in Einstein-Cartan theory*, *Astrophys. Space Sci.* **326** (2010) 19-26.
1481. Katkar L.N., *Einstein-Cartan theory of gravitation: kinematical parameters and Maxwell equations*, *Int. J. Theor. Phys.* **54** (2015) 951-971.
1482. Katkar L.N. and Patil V.K., *On spacelike congruences in Riemann-Cartan space-time*, *Int. J. Theor. Phys.* **48** (2009) 3035-3043.
1483. Katkar L.N. and Patil V.K., *Static conformally flat spherically symmetric space-time in Einstein-Cartan theory of gravitation*, *Int. J. Theor. Phys.* **51** (2012) 1050-1061.
1484. Kaul R.K. and Sengupta S., *Degenerate spacetimes in first order gravity*, *Phys. Rev.* **D93** (2016) 084026 (8 pages).
1485. Kaul R.K. and Sengupta S., *New solutions in pure gravity with degenerate tetrads*, *Phys. Rev.* **D94** (2016) 104047 (5 pages).
1486. Kawai T., *A five dimensional unification of the vierbein and electromagnetic field*, *Progr. Theor. Phys.* **67** (1982) 1946-1965.

1487. Kawai T., *A five dimensional unification of the vierbein and electromagnetic field. II*, *Progr. Theor. Phys.* **68** (1982) 1365-1376.
1488. Kawai T., *A five dimensional unification of the Poincaré gauge and electromagnetic field*, *Progr. Theor. Phys.* **69** (1983) 1272-1290.
1489. Kawai T., *A  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Gen. Relat. Grav.* **18** (1986) 995-1018.
1490. Kawai T., *(3+1)-structure of space-time in  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **76** (1986) 1166-1176.
1491. Kawai T., *Energy-momentum and angular momentum in  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **79** (1988) 920-935.
1492. Kawai T., *Extended spinor structure and exotic spinor field in  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **82** (1989) 850-862.
1493. Kawai T., *Group of paths, holonomy groups and nonintegrable phase factors in  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **84** (1990) 993-1007.
1494. Kawai T., *Teleparallel theory of (2+1)-dimensional gravity*, *Phys. Rev.* **D48** (1993) 5668-5675.
1495. Kawai T., *Poincaré gauge theory of (2+1)-dimensional gravity*, *Phys. Rev.* **D49** (1994) 2862-2871.
1496. Kawai T., *Black-hole-type solution in teleparallel theory of (2+1)-dimensional gravity*, *Progr. Theor. Phys.* **94** (1995) 1169-1174.
1497. Kawai T., *Energy-momentum and angular momentum densities in gauge theories of gravity*, *Phys. Rev.* **D62** (2000) 104014 (9 pages).
1498. Kawai T. and Saitoh H., *A note on  $\overline{\text{Poincaré}}$  gauge theory of gravity*, *Progr. Theor. Phys.* **81** (1989) 280-285.
1499. Kawai T. and Saitoh H., *A note on  $\overline{\text{Poincaré}}$  gauge theory of gravity. Additional comments*, *Progr. Theor. Phys.* **81** (1989) 1119-1123.
1500. Kawai T., Sakane E., and Tojo T., *Schwarzschild space-time in gauge theories of gravity*, *Progr. Theor. Phys.* **99** (1998) 971-992.
1501. Kawai T., Shibata K., and Tanaka I., *Generalized equivalence principle in extended new general relativity*, *Progr. Theor. Phys.* **104** (2000) 505-530.
1502. Kawai T. and Toma N., *Singularities of "Schwarzschild-like" space-time in new general relativity*, *Progr. Theor. Phys.* **83** (1990) 1-6.
1503. Kawai T. and Toma N., *An extended new general relativity as a reduction of Poincaré gauge theory of gravity*, *Progr. Theor. Phys.* **85** (1991) 901-926.
1504. Kawai T. and Toma N., *A charged Kerr metric solution in new general relativity*, *Progr. Theor. Phys.* **87** (1992) 583-598.
1505. Kawai T. and Yoshida H., *Special relativistic limit of gauge theory of gravitation*, *Progr. Theor. Phys.* **62** (1979) 793-801.
1506. Kawai T. and Yoshida H., *De Sitter gauge theory of gravitation*, *Progr. Theor. Phys.* **62** (1979) 266-277.
1507. Kawai T. and Yoshida H., *De Sitter gauge theory of gravitation. II*, *Progr. Theor. Phys.* **64** (1980) 1596-1607.
1508. Kawamoto N. and Nielsen H.M., *Lattice gauge gravity*, *Phys. Rev.* **D43** (1991) 1150-1156.
1509. Kaźmierczak M., *Nonuniqueness of gravity induced fermion interaction in the Einstein-Cartan theory*, *Phys. Rev.* **D78** (2008) 124025 (10 pages).

1510. Kaźmierczak M., *Einstein-Cartan gravity with Holst term and fermions*, *Phys. Rev.* **D79** (2009) 064029 (10 pages).
1511. Kaźmierczak M., *Modified coupling procedure for the Poincaré gauge theory of gravity*, *Phys. Rev.* **D79** (2009) 127501 (4 pages).
1512. Kelly P.F., *Topics in two-dimensional non-Riemannian gravity*, in: “Gravitation. Proc., Banff summer institute”, Banff, Canada, Aug 12-25, 1990, Ed. R. Mann (World Scientific: Singapore, 1991) 303-313.
1513. Kerlick G.D., *The effect of intrinsic spin on neutron stars in the Einstein-Cartan theory of gravity*, *Astrophys. J.* **185** (1973) 631-633.
1514. Kerlick G.D., *Cosmology and particle pair production via gravitational spin-spin interaction in the Einstein-Cartan-Sciama-Kibble theory of gravity*, *Phys. Rev.* **D12** (1975) 3004-3006.
1515. Kerlick G.D., “*Bouncing*” of simple cosmological models with torsion, *Ann. Phys. (USA)* **99** (1976) 127-141.
1516. Kerner R., *Geometrical background for the unified field theories: the Einstein-Cartan theory over a principal fibre bundle*, *Ann. Inst. H.Poincaré* **A34** (1981) 437-463.
1517. Ketsaris A.A., *Lie groups and gravitation theory*, *Sov. J. Phys.* **13** (1970) 1584-1590 [*Izvestia VUZov, Fizika*, n. 12 (1970) 40-47 (in Russian)].
1518. Khakshournia S., *Comment on ‘Braneworld remarks in Riemann-Cartan manifolds’*, *Class. Quantum Grav.* **26** (2009) 178001.
1519. Khatsymovsky V.M., *Affine connection form of Regge calculus*, *Mod. Phys. Lett.* **A31** (2016) 1650010 (8 pages).
1520. Khodunov A.V. and Zhytnikov V.V., *Gravitational equations in space-time with torsion*, *J. Math. Phys.* **33** (1992) 3509-3520.
1521. Khrapko R.I., *Classical spin in space with and without torsion*, *Grav. Cosmol.* **10** (2004) 91-97.
1522. Khriplovich I.B., *Gravitational four-fermion interaction on the Planck scale*, *Phys. Lett.* **B709** (2012) 111-113.
1523. Khriplovich I.B. and Pomeransky A.A., *Immirzi parameter, torsion, and discrete symmetries*, *Phys. Rev.* **D73** (2006) 107502 (2 pages).
1524. Khriplovich I.B. and Rudenko A.S., *Cosmology constrains gravitational four-fermion interaction*, *JCAP* **11** (2012) 040.
1525. Khriplovich I.B., and Rudenko A.S., *Gravitational four-fermion interaction and dynamics of the early Universe*, *JHEP* **11** (2013) 174.
1526. Kibble T.W.B., *Lorentz invariance and the gravitational field*, *J. Math. Phys.* **2** (1961) 212-221.
1527. Kibble T.W.B. and Stelle K.S., *Gauge theories of gravity and supergravity*, in: *Progress in Quantum Field Theory, Festschrift for H. Umezawa*, Eds. H. Ezawa and S. Kamefuchi (Elsevier Science Publ.: Amsterdam, 1986) 57-81.
1528. Kiehn R.M., *Topological torsion and topological spin*, *Annales de la Fondation L. de Broglie* **32** (2007) 389-408.
1529. Kikukawa H. and Kikukawa J., *Geometrical description of BRS and anti-BRS transformations for the gravitational fields*, *Progr. Theor. Phys.* **69** (1983) 1835-1836.
1530. Kikukawa H. and Kikukawa J., *BRS and anti-BRS transformations in  $GL(4, R)$  theory*, *Progr. Theor. Phys.* **70** (1983) 625-627.
1531. Kikukawa H. and Kikukawa J., *Geometric formulation of BRS and anti-BRS transformations in  $T_4$  theory*, *Progr. Theor. Phys.* **71** (1984) 174-182.

1532. Kikukawa H. and Kikukawa J., *Geometric description of BRS transformation for the fields in  $P_4$  theory*, *Progr. Theor. Phys.* **71** (1984) 875-877; Erratum: *Progr. Theor. Phys.* **72** (1984) 1060.
1533. Kikukawa H. and Kikukawa J., *Geometric description of BRS\* transformations for the fields in  $P_4$  theory*, *Progr. Theor. Phys.* **73** (1985) 979-986.
1534. Kikukawa H. and Kikukawa J., *Geometric approach to the BRS\* transformation in super-space Poincaré theory*, *Progr. Theor. Phys.* **73** (1985) 987-998.
1535. Kikukawa H., Kikukawa J., and Nakamura A., *Internal translation group, extended Cartan-Maurer condition and soldering forms*, *Progr. Theor. Phys.* **89** (1993) 523-530.
1536. Kikukawa H., Kikukawa J., and Nakamura A., *Internal  $P_{CI}^4$  group in spin space and gauge theory of gravity*, *Progr. Theor. Phys.* **89** (1993) 1063-1075.
1537. Kim E.J., Kouwn S., Oh P., and Park C.-G., *Dark aspects of massive spinor electrodynamics*, *JCAP* **07** (2014) 001.
1538. Kim S.-W., *Brans-Dicke theory in general space-time with torsion*, *Phys. Rev.* **D34** (1986) 1011-1013.
1539. Kim S.-W., *The cosmological model with scalar, spin and torsion field*, *Nuovo Cim.* **B112** (1997) 363-373.
1540. Kim S.-W., *The gravitational wave and the dipole moment in Einstein-Cartan theory*, *Hadronic J.* **21** (1998) 95-109.
1541. Kim S.-W., *The gravitational wave of the scalar-tensor theory*, *J. Korean Phys. Soc. Suppl.* **33** (1998) S560-S564.
1542. Kim S.-W. and Pak D.G., *Torsion as a dynamic degree of freedom of quantum gravity*, *Class. Quantum Grav.* **25** (2008) 065011.
1543. Kim J.K. and Yoon Y., *Yukawa-type potential besides Newtonian potential in Poincaré gauge theory of gravitation*, *Phys. Lett.* **A121** (1987) 213-216.
1544. Kim J.K. and Yoon Y., *The relationship between conformal and gravitational anomalies*, *Phys. Lett.* **B214** (1988) 98-104.
1545. Kim J., Park C.J., and Yoon Y., *Phase transition in conformally induced gravity with torsion*, *Phys. Rev.* **D51** (1995) 562-567.
1546. Kim J., Park C.J., and Yoon Y., *Constraint of a general effective potential in vector torsion coupled conformally induced gravity*, *Phys. Rev.* **D51** (1995) 4595-4597.
1547. Kim Sung-Won, *Brans-Dicke theory in general space with torsion*, *Phys. Rev.* **D34** (1986) 1011-1013.
1548. Kim S.-W., *The cosmological model with scalar, spin and torsion field*, *Nuovo Cim.* **B112** (1997) 363-373.
1549. Kim Sung-Won and Cho Byung He, *Unification of the Einstein-Cartan theory and the relativistic kinetic theory*, *Phys. Lett.* **A94** (1983) 210-212.
1550. Kim Sung-Won and Cho Byung He, *Static vacuum solutions of Brans-Dicke theory with torsion*, *Phys. Lett.* **A124** (1987) 243-244.
1551. Kim Sung-Won, and Cho Byung He, *Scalar-tensor theory in higher-dimensional space-time with torsion*, *Phys. Rev.* **D36** (1988) 2314-2314.
1552. Kim S.-W., Kim S.P., Lee J., and Park D.H., *Brans-Dicke cosmological model with torsion*, *J. Korean Phys. Soc. Suppl.* **28** (1995) S519-S524.
1553. Kim S.K., Soh K.S., and Yee J.H., *(1+1)-dimensional gravity through a dimensional reduction*, *Phys. Lett.* **B300** (1993) 223-226.

1554. Kim S.K., Soh K.S., and Yee J.H., *GL(2,R) gauge theory of (1+1)-dimensional gravity*, *Phys. Rev.* **D47** (1993) 4433-4437.
1555. Kimura T., *Expansion coefficient of heat kernel of Laplacian operator in Riemann-Cartan space*, *J. Phys.* **A14** (1981) L329-L331.
1556. Kimura T., *Conformal and axial anomalies in Riemann-Cartan space*, *Progr. Theor. Phys.* **66** (1981) 2011-2024.
1557. Kimura T., *Index theorems on torsional geometries*, *JHEP* **08** (2007) 048.
1558. Kimura T. and Nishioka T., *The chiral heat effect*, *Prog. Theor. Phys.* **127** (2012) 1009-1017.
1559. King A.D. and Vassiliev D., *Torsion waves in metric-affine field theory*, *Class. Quantum Grav.* **18** (2001) 2317-2329.
1560. Kirsch I., *Higgs mechanism for gravity*, *Phys. Rev.* **D72** (2005) 024001 (16 pages).
1561. Kirsch I. and Šijački Dj., *From Poincaré to affine invariance: how does the Dirac equation generalize?* *Class. Quantum Grav.* **19** (2002) 3157.
1562. Kiriushcheva N. and Kuzmin S.V., *Translational invariance of the Einstein-Cartan action in any dimension*, *Gen. Rel. Grav.* **42** (2010) 2613-2631.
1563. Kleinert H., *Gauge fields in condensed matter* (World Scientific: Singapore, 1989) vol. 1-2, pp. 1474.
1564. Kleinert H., *Quantum dynamics in spaces with curvature and torsion*, *Mod. Phys. Lett.* **A4** (1989) 2329.
1565. Kleinert H., *Non-Abelian bosonization as a nonholonomic transformation from a flat to a curved field space*, *Ann. Phys.* **253** (1997) 121-176.
1566. Kleinert H., *Nonholonomic mapping principle for classical mechanics in spacetimes with curvature and torsion. New covariant conservation law for energy-momentum tensor*, *Acta Phys. Pol.* **B29** (1998) 1033-1045.
1567. Kleinert H., *Spontaneous generation of torsion coupling of electroweak massive gauge bosons*, *Phys. Lett.* **B440** (1998) 283-286.
1568. Kleinert H., *Nonholonomic mapping principle for classical and quantum mechanics in spaces with curvature and torsion*, *Gen. Relat. Grav.* **32** (2000) 769-839.
1569. Kleinert H., *Torsion, a multivalued gauge degree of freedom in Einstein's gravity*, *J. Phys.: Conf. Ser.* **306** (2011) 012020.
1570. Kleinert H., *Universality principle for orbital angular momentum and spin in gravity with torsion*, *Gen. Relat. Grav.* **32** (2000) 1271-1280.
1571. Kleinert H., *Multivalued fields in condensed matter, electromagnetism, and gravitation* (World Scientific: Singapore, 2008) 524 pp.
1572. Kleinert H., *Path integrals in quantum mechanics, statistics, polymer physics, and financial markets*, 5th ed. (World Scientific: Singapore, 2009) 1624 pp.
1573. Kleinert H., *New gauge symmetry in gravity and the evanescent role of torsion*, in: "Quantum Mechanics, Elementary Particles, Quantum Cosmology and Complexity", *Proc. Conf. in Honour of Murray Gell-Mann's 80th Birthday*, Eds. H. Fritzsche and K.K. Phua (World Scientific: Singapore, 2010) 174-185.
1574. Kleinert H. and Pelster A., *Novel geometric gauge invariance of autoparallels*, *Acta Phys. Pol.* **B29** (1998) 1015-1023.
1575. Kleinert H. and Shabanov S.V., *Spaces with torsion from embedding, and the special role of autoparallel trajectories*, *Phys. Lett.* **B428** (1998) 315-321.
1576. Kleinert H. and Shabanov S.V., *Theory of Brownian motion of a massive particle in spaces with curvature and torsion*, *J. Phys. A: Math. Gen.* **31** (1998) 7005-7009.

1577. Kleinert H. and Zaanen J., *Nematic world crystal model of gravity explaining absence of torsion in spacetime*, *Phys. Lett.* **A324** (2004) 361-365.
1578. Kleman M. and Friedel J., *Disclinations, dislocations, and continuous defects: A reappraisal*, *Rev. Mod. Phys.* **80** (2008) 61-115.
1579. Klosch T. and Strobl T., *Classical and quantum gravity in  $(1+1)$ -dimensions. Part 1: A unifying approach*, *Class. Quantum Grav.* **13** (1996) 965-984.
1580. Kofinas G., Leon G., and Saridakis E.N., *Dynamical behavior in  $f(T, T_G)$  cosmology*, *Class. Quantum Grav.* **31** (2014) 175011.
1581. Kofinas G., Papantonopoulos E., and Saridakis E.N., *Self-gravitating spherically symmetric solutions in scalar-torsion theories*, *Phys. Rev.* **D91** (2015) 104034 (14 pages).
1582. Kofinas G. and Saridakis E.N., *Teleparallel equivalent of Gauss-Bonnet gravity and its modifications*, *Phys. Rev.* **D90** (2014) 084044 (11 pages).
1583. Kofinas G. and Saridakis E.N., *Cosmological applications of  $F(T, T_G)$  gravity*, *Phys. Rev.* **D90** (2014) 084045 (10 pages).
1584. Kofinti N.K., *On the effect of spin on the gravitational field*, *Int. J. Theor. Phys.* **12** (1975) 241-249.
1585. Kohler C., *Point particles in  $2+1$  dimensional gravity as defects in solid continua*, *Class. Quantum Grav.* **12** (1995) L11-L15.
1586. Kohler C., *Geometric classification of topological quantum phases*, *Phys. Lett.* **A237** (1998) 195-200.
1587. Kohler C., *Semi-teleparallel theories of gravitation*, *Gen. Relat. Grav.* **32** (2000) 1301-1317.
1588. Kohler C., *The Einstein-Cartan-Maxwell theory with scalar field through a five-dimensional unification*, *Int. J. Mod. Phys.* **A15** (2000) 1235-1243.
1589. Kokado A., Konisi G., Saito T., and Tada Y., *Brans-Dicke theory from  $M(4) \times Z(2)$  geometry*, *Progr. Theor. Phys.* **99** (1998) 293-303.
1590. Kondo K., *On the geometrical and physical foundations of the theory of yielding*, in: *Proc. 2nd Japan Nat. Congr. Appl. Mech.* (1952) 41-47.
1591. Kondo K., *Geometry of elastic deformation and incompatibility*, *Memoirs of the Unifying Study of the Basic Problems in Engineering Science by Means of Geometry* (ed. K. Kondo), vol. 1, Division C, (Gakujutsu Bunken Fukyo-Kai: 1955) pp. 5-17.
1592. Kondo K., *Non-Riemannian geometry of imperfect crystals from a macroscopic viewpoint*, *Memoirs of the Unifying Study of the Basic Problems in Engineering Science by Means of Geometry* (ed. K. Kondo), vol. 1, Division D-I, (Gakujutsu Bunken Fukyo-Kai: 1955) pp. 6-17.
1593. Kondo K., *Dimensional consistency in physics and the ordered construction of the world of elementary particles*, in: "On Relativity Theory. Proc. of Sir A. Eddington Centenary Symp.", Eds. Y. Choquet-Bruhat and T.M. Karade (World Scientific: Singapore, 1895) vol. 2, 57-121.
1594. Kong K.H., Cheung P.C.H., Lu H.Q., and Cheng K.S., *Inflation in Bianchi type-IX Einstein-Cartan cosmological model*, *Astrophys. and Space Sci.* **260** (1998) 521-529.
1595. Kong K.H., Cheng K.S., Fung P.C.W., and Lu H.Q., *Bianchi cosmology with spinning matter and magnetic field in the Einstein-Cartan theory*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 451-453.
1596. Konopleva N.P. and Popov V.N., *Gauge fields* (Harwood Academic Publ.: Chur, 1981) 264 pp.
1597. Kopczyński W., *A non-singular universe with torsion*, *Phys. Lett.* **A39** (1972) 219-220.

1598. Kopczyński W., *An anisotropic universe with torsion*, *Phys. Lett.* **A43** (1973) 63-64.
1599. Kopczyński W., *The Palatini principle with constraints*, *Bull. Acad. Polon. Sci., sér. sci. math., astr., phys.*, **23** (1975) 467-473.
1600. Kopczyński W., *The Einstein-Cartan theory of gravitation*, in: *Scripta Fac. Sci. natur UJEP brun. Phys.* **3-4** (1975) 255-263.
1601. Kopczyński W., *The role of spin in early universe*, in: "Energy and Physics", *Proc. of the 3rd Gen. Conf., Bucharest-1975* (1976) 243-247.
1602. Kopczyński W., *Metric-affine unification of gravity and gauge theories*, *Acta Phys. Pol.* **B10** (1979) 365-367.
1603. Kopczyński W., *A fibre bundle description of coupled gravitational and gauge fields*, *Lect. Notes Math.* **836** (1980) 462-483.
1604. Kopczyński W., *Problems with metric-teleparallel theories of gravitation*, *J. Phys.* bf A15 (1982) 493-506.
1605. Kopczyński W., *Lagrangian dynamics of particles and fluids with intrinsic spin in Einstein-Cartan space-time*, *Phys. Rev.* **D34** (1986) 352-356.
1606. Kopczyński W., *Variational principles for gravity and fluids*, *Ann Phys. (USA)* **203** (1990) 308-338.
1607. Kopczyński W., McCrea J.D., and Hehl F.W., *The Weyl group and its currents*, *Phys. Lett.* **A128** (1988) 313-317.
1608. Kopczyński W., McCrea J.D., and Hehl F.W., *The metric and the canonical energy-momentum currents in the Poincaré gauge theory of gravitation*, *Phys. Lett.* **A135** (1989) 89-91.
1609. Kopczyński W. and Trautman A., *Spacetime and gravitation* (Wiley, Chichester and PWN: Warszawa, 1992).
1610. Koppa S.S. and Patel L.K., *Interior fields of charged spheres in the Einstein-Cartan theory*, *Acta Phys. Pol.* **20** (1989) 165-173.
1611. Korniyak V.V., *Heat invariant  $E_2$  for nonminimal operator on manifolds with torsion*, in: "Computer Algebra in Scientific Computing" Eds. Ganzha V.G., Mayr E.W., and Vorozhtsov E.V. (Springer: Berlin, 2000) 273-284.
1612. Korotky V.A., *Primordial nucleosynthesis in the rotating and expanding universe*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds.P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 121-127.
1613. Korotky V.A. and Obukhov Yu.N., *Spin fluid in gauge gravity*, *Vestnik Mosk. Univ., ser. Fiz., Astr.*, **28**, n. 4 (1987) 6-11 (in Russian).
1614. Korotky V.A. and Obukhov Yu.N., *Cosmology in Einstein-Cartan theory I. Spin dynamics*, *Vestnik Mosk. Univ., ser. Fiz. Astr.*, **28**, n. 6 (1987) 3-8 (in Russian).
1615. Korotky V.A. and Obukhov Yu.N., *Cosmology in Einstein-Cartan theory II. Some exact solutions*, *Vestnik Mosk. Univ. ser. Fiz. Astr.* **29**, n. 1 (1988) 12-17 (in Russian).
1616. Korotky V.A. and Obukhov Yu.N., *Generalized model of fluid with spin*, *Izvestia VUZov, Fizika*, n. 2 (1991) 98-101 (in Russian).
1617. Korotky V.A. and Obukhov Yu.N., *Rotating and expanding cosmology in ECSK-theory*, *Astrophys. and Space Sci.* **198** (1992) 1-12.
1618. Korotky V.A. and Obukhov Yu.N., *On cosmic rotation*, in: "Gravity, Particles and Space-time", Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996) 421-439.
1619. Korotky V.A. and Obukhov Yu.N., *Rotating Cosmologies in Poincaré Gauge Gravity*, in: "The Universe Rotation: Pro and Contra", Ed. R. Myrzakulov (Nova Science Publishers: Hauppauge, NY, 2017) p. 157-206.

1620. Koryukin V.M., *Gravitation and unified covariant formalism of classical gauge fields in equi-affine space*, *Sov. J. Phys.* **24** (1981) 1108-1111 [*Izvestia VUZov, Fizika*, n. 12 (1981) 47-51 (in Russian)].
1621. Kostelecký V.A., ed. "CPT and Lorentz Symmetry", *Proc. 2nd Meeting Indiana University, Bloomington, USA, 15-18 August 2001* (World Scientific: Singapore, 2002) 368 pp.
1622. Kostelecký V.A., *Gravity, Lorentz violation, and the standard model*, *Phys. Rev.* **D69** (2004) 105009 (20 pages).
1623. Kostelecký V.A., ed. "CPT and Lorentz Symmetry", *Proc. 4th Meeting Indiana University, Bloomington, USA, 8-11 August 2007* (World Scientific: Singapore, 2007) 336 pp.
1624. Kostelecký V.A., ed. "CPT and Lorentz Symmetry", *Proc. 5th Meeting Indiana University, Bloomington, USA, 28 June-2 July 2010* (World Scientific: Singapore, 2010) 312 pages.
1625. Kostelecký V.A. and Mewes M., *Cosmological constraints on Lorentz violation in electrodynamics*, *Phys. Rev. Lett.* **87** (2001) 251304 (4 pages).
1626. Kostelecký V.A. and Mewes M., *Fermions with Lorentz-violating operators of arbitrary dimension*, *Phys. Rev.* **D88** (2013) 096006 (30 pages).
1627. Kostelecký V.A. and Russell R., *Data tables for Lorentz and CPT violation*, *Rev. Mod. Phys.* **83** (2011) 11-31.
1628. Kostelecký V.A., Russell R., and Tasson J.D., *Constraints on torsion from bounds on Lorentz violation*, *Phys. Rev. Lett.* **100** (2008) 111102 (4 pages).
1629. Kostelecký V.A. and Tasson J.D., *Prospects for large relativity violations in matter-gravity couplings*, *Phys. Rev. Lett.* **102** (2009) 010402 (4 pages).
1630. Kostelecký V.A. and Tasson J.D., *Matter-gravity couplings and Lorentz violation*, *Phys. Rev.* **D83** (2011) 016013 (59 pages).
1631. Kostkin R.S., *Solving problems of gravitation theory with the help of the Mathematica package for symbolic Cartan-Weyl calculations*, *Russ. Phys. J.* **52** (2009) 1000-1002.
1632. Kotler S., Ozeri R., and Jackson Kimball D.F., *Constraints on exotic dipole-dipole couplings between electrons at the micrometer scale*, *Phys. Rev. Lett.* **115** (2015) 081801 (5 pages).
1633. Kounieher J. and Barbachoux C., *Cartan's soldered spaces and conservation laws in physic*, *Int. J. Geom. Meth. in Mod. Phys.* **12** (2015) 1550089 (20 pages)
1634. Koun S., Lee J., Lee T.-H., and Oh P., *ELKO spinor model with torsion and cosmology*, *Mod. Phys. Lett.* **A28** (2013) 1350121 (14 pages).
1635. Krasinski A. and Perkowski M., *ORTOCARTAN - a new computer program for analytic calculations in general relativity*, *Gen. Relat. Grav.* **13** (1981) 67-77.
1636. Krechet V.G., *On the correspondence of gravitating wave fields of equal tensor dimension*, *Sov. J. Phys.* **21** (1978) 656-657 [*Izvestia VUZov, Fizika*, n. 5 (1978) 128-129 (in Russian)].
1637. Krechet V.G., *Spinor and electromagnetic fields in spacetime with torsion*, *Sov. J. Phys.* **21** (1978) 1412-1415 [*Izvestia VUZov, Fizika*, n. 11 (1978) 31-35 (in Russian)].
1638. Krechet V.G., *Free vector field in homogeneous space with torsion*, *Sov. J. Phys.* **22** (1979) 945-948 [*Izvestia VUZov, Fizika*, n. 9 (1979) 26-29 (in Russian)].
1639. Krechet V.G., *Configurations with a vector field in the Einstein-Cartan theory*, *Sov. J. Phys.* **22** (1979) 809-812 [*Izvestia VUZov, Fizika*, n. 8 (1979) 7-11 (in Russian)].
1640. Krechet V.G., *Spinor field and non-metricity of space-time*, *Sov. J. Phys.* **23** (1980) 506-509 [*Izvestia VUZov, Fizika*, n. 6 (1980) 52-55 (in Russian)].
1641. Krechet V.G., *Gravitational interaction of vector field and non-metricity of space-time*, *Sov. J. Phys.* **27** (1984) 671-674 [*Izvestia VUZov, Fizika*, n. 8 (1984) 50-54 (in Russian)].

1642. Krechet V.G., *Continuous medium dynamics in space with torsion*, *Sov. J. Phys.* **28** (1985) 957-961 [*Izvestia VUZov, Fizika*, n. 12 (1985) 9-14 (in Russian)].
1643. Krechet V.G., *Spinor analysis and the physical properties of fermions*, *Sov. J. Phys.* **29** (1986) 790-794 [*Izvestia VUZov, Fizika*, n. 10 (1986) 20-25 (in Russian)].
1644. Krechet V.G., *Spinor electrodynamics and space-time torsion*, *Izvestia VUZov, Fizika*, n. 3 (1988) 114-116 (in Russian).
1645. Krechet V.G., *Gauge theory of gravitational interaction of macroscopic matter*, *Russ. Phys. J.* **34** (1991) 630-632 [*Izvestia VUZov, Fizika*, n. 7 (1991) 81-83 (in Russian)].
1646. Krechet V.G., *Electrodynamics of leptons in curved space and weak interaction*, *Russ. Phys. J.* **37** (1994) 929-931 [*Izvestia VUZov, Fizika*, n. 10 (1994) 18-21 (in Russian)].
1647. Krechet V.G., *Topological and physical effects of rotation and spin in the general relativistic theory of gravitation*, *Russ. Phys. J.* **50** (2007) 1021-1025.
1648. Krechet V.G., Lapchinsky V.G., and Ponomarev V.N., *Gravitation, spin, torsion*, in: "Probl. Theor. Grav. and Elem. Particles", Ed. K.P. Stanyukovich (Atomizdat: Moscow) **6** (1975) 139-146 (in Russian).
1649. Krechet V.G., Lapchinsky V.G., Ponomarev V.N., and Stanyukovich K.P., *Problems of gravitational collapse*, in: "Probl. Theor. Grav. and Elem. Particles", Ed. K.P. Stanyukovich (Atomizdat: Moscow) **6** (1975) 129-139 (in Russian).
1650. Krechet V.G., Lapchisky V.G., and Ponomarev V.N., *Wave fields in the Einstein-Cartan theory of gravity*, in: "Relativity Theory and Gravitation", Ed. V.I. Rodichev (Nauka: Moscow, 1976) 57-63 (in Russian).
1651. Krechet V.G., Levkoeva M.V., and Sadovnikov D.V., *A five-dimensional model of the gravitational interaction of an electromagnetic field in an affine-metric space*, *Russ. Phys. J.* **43** (2000) 933-937 [*Izvestia VUZov, Fizika*, n. 11 (2000) 43-47 (in Russian)].
1652. Krechet V.G., Levkoeva M.V., and Sadovnikov D.V., *A five-dimensional model of the gravitational interaction of an electromagnetic field in an affine-metric space*, *Russ. Phys. J.* **44** (2001) 933-937.
1653. Krechet V.G. and Ponomarev V.N., *Nonlinearity and torsion*, *Theor. Math. Phys.* **24** (1975) 1036-1038 [*Teor. Mat. Fiz.* **24** (1975) 141-144 (in Russian)].
1654. Krechet V.G. and Ponomarev V.N., *Nonlinearity and torsion*, *Acta Phys. Pol.* **B7** (1976) 553-556.
1655. Krechet V.G. and Ponomarev V.N., *On the analogy between neutrino and non-linear spinor fields*, *Phys. Lett.* **A56** (1976) 14-15.
1656. Krechet V.G. and Ponomarev V.N., *Observable effects of space-time torsion (equations of motion)*, in: "Probl. Theor. Grav. Elem. Particles", Ed. K.P. Stanyukovich (Atomizdat: Moscow) **7** (1976) 174-184 (in Russian).
1657. Krechet V.G. and Ponomarev V.N., *Problems of gravitational collapse, neutrino dynamics and gravitational vacuum*, in: "Actual Problems of Theor. Physics. Prof. D.Ivanenko's Festschrift", Ed. A.A. Sokolov (Moscow Univ. Publ. House: Moscow, 1976) 165-182 (in Russian).
1658. Krechet V.G. and Ponomarev V.N., *Cosmology in the Einstein-Cartan theory*, in: "Advances in the relativity and gravity theory" (Nauka: Moscow, 1977) 66-70 (in Russian).
1659. Krechet V.G., Ponomarev V.N., and Pronin P.I., *Observable effects of spacetime torsion*, in: "Advances in the relativity and gravity theory" (Nauka: Moscow, 1977) 71-74 (in Russian).
1660. Krechet V.G. and Sandina I.V., *New features of the gravitational interaction of a spinor field*, *Russ. Phys. J.* **26** (1983) 448-450 [*Izvestia VUZov, Fizika*, n. 5 (1983) 33-36 (in Russian)].
1661. Kreimer D. and Mielke E.W., *Comment on "Topological invariants, instantons, and the chiral anomaly on spaces with torsion"*, *Phys. Rev.* **D63** (2001) 048501 (4 pages).

1662. Kreinovich V., *Only particles with spin  $\leq 2$  are mediators for fundamental forces: Why?*, *Phys. Essays* **5** (1992) 458-462.
1663. Krisch J.P., *Spin distributions for a spheroidal Cartan fluid*, *Phys. Rev.* **D32** (1985) 1369-1374.
1664. Krisch J.P., *Spin fluids in stationary axis-symmetric space-times*, *J. Math. Phys.* **28** (1987) 1544-1546.
1665. Krisch J.P., *A family of strings with spin*, *Gen. Rel. Grav.* **28** (1996) 69-76.
1666. Krisch J.P. and Smalley L.L., *Cartan and relativistic spin fluids in a rotating cylinder*, *J. Math. Phys.* **29** (1988) 1640-1644.
1667. Krisch J.P. and Smalley L.L., *Spin fluids in van Stockum cylinders*, *Class. Quantum Grav.* **7** (1990) 481-486.
1668. Krisch J.P. and Smalley L.L., *Spin fluid spacetimes from static general relativistic solutions*, *Class. Quantum Grav.* **10** (1993) 995-1003.
1669. Krisch J.P. and Smalley L.L., *A spin polarized disc*, *Gen. Rel. Grav.* **27** (1995) 247-255.
1670. Krisch J.P. and Smalley L.L., *A spin polarized disc. II*, *Gen. Rel. Grav.* **28** (1996) 287-297.
1671. Krori K.D., Borgahain P., Das K., and Sarma A., *Exact interior solutions in Einstein-Cartan-Yukawa theory theory*, *Can. J. Phys.* **66** (1988) 126-131.
1672. Krori K.D., Sheikh A.R., and Mahanta L., *A singularity-free solution for a static charged fluid sphere in Einstein-Cartan theory*, *Can. J. Phys.* **59** (1981) 425-427.
1673. Kröner E., *Algemeine Kontinuums theorie der Versetzungen und Eigenspannungen*, *Arch. Rat. Mech. Anal.* **4** (1960) 273-334.
1674. Kröner E., *Continuum theory of defects*, in: "Physique des defaults" *Les Houches, 1980, Session 35*, Eds. R.Balain et al. (North-Holland: Amsterdam, 1981) 215-315.
1675. Kröner E., *The role of differential geometry in the mechanics of solids*, in: "Proc. 5th Nat. Congr. Theor. Appl. Mech." Vol. **1** (Bulgarian Acad. Sci.: Sofia, 1985) 352-362.
1676. Kröner E., *On gauge theory in defect mechanics*, in: "Trends in Applications of pure Mathematics to Mechanics", *Lect. Notes Phys.* **249** (1986) 281-294.
1677. Kröner E., *The continuized crystal — a bridge between micro- and macromechanics*, *Z. angew. Math. Mech.* **66** (1986) T284-T292.
1678. Kröner E., *The differential geometry of elementary point and line defects in Bravais crystals*, *Int. J. Theor. Phys.* **29** (1990) 1219-1237.
1679. Kröner E., *The internal mechanical state of solids with defects*, *Int. J. Solids Struct.* **29** (1992) 1849-1857.
1680. Kröner E. and Anthony K.H., *Dislocations and disclinations in material structures: the basic topological concepts*, *Annual Review of Materials Science* **5** (1975) 43-72.
1681. Krššák M., *Holographic renormalization in teleparallel gravity*, *Eur. Phys. J. C* **77** (2017) 44 (5 pages).
1682. Krššák M. and Pereira J.G., *Spin connection and renormalization of teleparallel action*, *Eur. Phys. J. C* **75** (2015) 519 (8 pages).
1683. Krššák M. and Saridakis E.N., *The covariant formulation of  $f(T)$  gravity*, *Class. Quantum Grav.* **33** (2016) 115009 (15 pages).
1684. Kruger J., De Meyer H., and van den Berghe G., *Derivation of a model for symmetrized electromagnetism from a space-time with torsion*, *J. Phys.* **A9** (1976) 1323-1329.
1685. Kruglov S.I., *On torsion fields in higher derivative quantum gravity*, *Annales de la Fondation L. de Broglie* **32** (2007) 323-334.

1686. Kruglov S.I., *On possible light-torsion mixing in background magnetic field*, Eur. Phys. J. **C63** (2009) 343-347.
1687. Kubota S.-I., Yajima S., Higasida Y., and Kawano K., *Gravitational anomalies with torsion in four dimensions*, Class. Quantum Grav. **18** (2001) 433-440.
1688. Kubyshin Yu.A., Malysenko V.O., and Ricoy D.M., *Invariant connections with torsion on group manifolds and their application in Kaluza-Klein theories*, J. Math. Phys. **35** (1994) 310-320.
1689. Kubyshin Yu.A., Richter O., and Rudolph G., *On the equations of spontaneous compactification for Gauss-Bonnet extended Einstein-Cartan theory*, Repts. Math. Phys. **30** (1991) 355-361.
1690. Kubyshin Yu.A., Richter O., and Rudolph G., *Invariant connections on homogeneous spaces*, J. Math. Phys. **34** (1993) 5268-5282.
1691. Kubyshin Yu.A., Mourão J.M., Rudolph G., and Volobujev I.P., *Dimensional reduction of gauge theories, spontaneous compactification and model building* (Springer: Berlin, 1989) Lect. Notes Phys. **349** 80 p.
1692. Kuchowicz B., *Neutrinos in general relativity: four (?) levels of approach*, Gen. Relat. Grav. **5** (1974) 201-234.
1693. Kuchowicz B., *Two-component neutrino field and spherical symmetry in a space-time with curvature and torsion*, Phys. Lett. **A50** (1974) 267-268.
1694. Kuchowicz B., *Flat anisotropic models of the universe with torsion and without singularity*, J. Phys. **A8** (1975) L29-L31.
1695. Kuchowicz B., *Methods of deriving exact solutions of spherical symmetry in the Einstein-Cartan theory for a perfect fluid with a "classical description" of spin*, Acta Phys. Pol. **B6** (1975) 173-196.
1696. Kuchowicz B., *A static universe filled with spinning matter and magnetic field*, Current Sci. (India) **44** (1975) 537-538.
1697. Kuchowicz B., *Cosmology with spin and torsion. I. Physical and mathematical foundations*, Acta Cosmologica **Z3** (1975) 109-129.
1698. Kuchowicz B., *Cosmology with spin and torsion. II. Spatially homogeneous aligned spin models with the Weyssenhoff fluid*, Acta Cosmologica **Z4** (1976) 67-100.
1699. Kuchowicz B., *The Einstein-Cartan equations in astrophysically interesting situations. I. The case of spherical symmetry*, Acta Phys. Pol. **B6** (1975) 555-575.
1700. Kuchowicz B., *The Einstein-Cartan equations in astrophysically interesting situations. II. Homogeneous cosmological models of axial symmetry*, Acta Phys. Pol. **B7** (1976) 81-97.
1701. Kuchowicz B., *Some cosmological models with spin and torsion, I*, Astrophys. Space Sci. **39** (1976) 157-172.
1702. Kuchowicz B., *Some cosmological models with spin and torsion, II. Axially symmetric models with uniform magnetic field*, Astrophys. Space Sci. **40** (1976) 167-181.
1703. Kuchowicz B., *Spin and torsion provide the bouncing mechanism for a non-singular pulsating universe*, Current Sci. (India) **45** (1976) 281-282.
1704. Kuchowicz B., *Zapobieganie osobliwosciom w kosmologii Einsteina-Cartana*, Post. Astron. **25** (1977) 27-33 (in Polish).
1705. Kuchowicz B., *Friedmann-like cosmological models without singularity*, Gen. Relat. Grav. **9** (1978) 511-517.
1706. Kuchowicz B. and Zebrowski J., *The presence of torsion enables a metric to allow a gravitational field*, Phys. Lett. **A67** (1978) 16-18.

1707. Kuchumov A.Yu., *Hamiltonian formulation of gravitation theory with a scalar field in the space-time having the Riemann-Cartan geometric structure*, *Russ. Phys. J.* **52** (2009) 930-937.
1708. Kudin B.I., Minkevich A.V., and Fedorov F.I., *On homogeneous isotropic cosmological models with torsion*, *Izvestia AN BSSR ser. Fiz.-Math. nauk*, n. 4 (1981) 59-67 (in Russian).
1709. Kuhne R.W., *Gauge theory of gravity requires massive torsion field*, *Int. J. Mod. Phys.* **A14** (1999) 2531-2535.
1710. Kunin I.A. and Kunin B.I., *Gauge theories in mechanics*, in: *Trends in Applications of pure Mathematics to Mechanics*, *Lect. Notes Phys.* **249** (1986) 246-269.
1711. Kuhfuss R. and Nitsch J., *Propagating modes in gauge field theories of gravity*, *Gen. Relat. Grav.* **18** (1986) 1207-1227.
1712. Kummer W., *Deformed  $iso(2,1)$  symmetry and non-Einsteinian 2d-gravity with matter*, in: *Proc. of the Conf. "Hadron Structure 1992"*, Stará Lesná (Slovakia), 6-11 Sept. 1992, D. Brunsko and J. Urban, eds. (Košice University Publications, Košice 1992) 48-56.
1713. Kummer W., *2d quantum gravity and black hole formation*, *Nucl. Phys. B, Proc. Suppl.* **88** (2000) 281-282.
1714. Kummer W. and Lau S.R., *Boundary conditions and quasilocal energy in the canonical formulation of all 1+1 models of gravity*, *Ann. Phys.* **258** (1997) 37-80.
1715. Kummer W., Liebl H., and Vassilevich D.V., *Integrating geometry in general 2D dilaton gravity with matter*, *Nucl. Phys.* **B544** (1999) 403-431.
1716. Kummer W. and Schütz H., *Linear derivative Cartan formulation of general relativity*, *Eur. Phys. J.* **C42** (2005) 227-241.
1717. Kummer W. and Schwarz D.J., *Renormalization of  $R^2$  gravity with dynamical torsion in  $d=2$* , *Nucl. Phys.* **B382** (1992) 171-186.
1718. Kummer W. and Schwarz D.J., *General analytic solution of  $R^2$  gravity with dynamical torsion in two dimensions*, *Phys. Rev.* **D45** (1992) 3628-3635.
1719. Kummer W. and Schwarz D.J., *Two-dimensional  $R^2$ -gravity with torsion*, *Class. Quantum Grav. Suppl.* **10** (1993) S235-S238.
1720. Kummer W. and Schwarz D.J., *Comment on "Canonical description of a two-dimensional gravity"*, *Mod. Phys. Lett.* **A8** (1993) 2903.
1721. Kummer W. and Tieber G., *Universal conservation law and modified Noether symmetry in 2D models of gravity with matter*, *Phys. Rev.* **D59** (1999) 044001 (10 pages).
1722. Kummer W. and Widerin P., *Non-Einsteinian gravity in  $d = 2$ : symmetry and current algebra*, *Mod. Phys. Lett.* **A9** (1994) 1407-1414.
1723. Kummer W. and Widerin P., *Conserved quasilocal quantities and general covariant theories in two dimensions*, *Phys. Rev.* **D52** (1995) 6965-6975.

## L

1724. Lagraa M.-H. and Lagraa M., *On the generalized Einstein-Cartan action with fermions*, *Class. Quantum Grav.* **27** (2010) 095012.
1725. Lämmerzahl C., *Constraints on space-time torsion from Hughes-Drever experiments*, *Phys. Lett.* **A228** (1997) 223-231.
1726. Lämmerzahl C., *Minimal coupling and the equivalence principle in quantum mechanics*, *Acta Phys. Pol.* **B29** (1998) 1057-1070.

1727. Lämmerzahl C., *Quantum tests of space-time structure*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?", *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. De Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 91-117.
1728. Lämmerzahl C., *New constraints on space-time torsion from Hughes-Drever experiments*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 454-456.
1729. Lämmerzahl C., Puntigam R.A., and Hehl F.W., *Can the electromagnetic field couple to post-Riemannian structures?*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 457-459.
1730. Lasenby A., Doran C., and Gull S., *Gravity, gauge theories and geometric algebra*, *Philos. Trans. R. Lond.* **A355** (1998) 487-582.
1731. Lasenby A.N. and Hobson M.P., *Scale-invariant gauge theories of gravity: Theoretical foundations*, *J. Math. Phys.* **57** (2016) 092505 (77 pages).
1732. Lavrov P.M. and Moshin P.Yu., *Covariant quantization of the Volovich-Katanaev model*, *Theor. and Math. Phys.* **114** (1998) 313-321.
1733. Lavrov P.M. and Moshin P.Yu., *Quantization of two-dimensional gravity with dynamical torsion*, *Class. Quantum Grav.* **16** (1999) 2247-2258.
1734. Lazar M., *An elastoplastic theory of dislocations as a physical field theory with torsion*, *J. Phys. A: Math. Gen.* **35** (2002) 1983-2004.
1735. Lazar M., *Wedge disclination in the field theory of elastoplasticity*, *Phys. Lett.* **A311** (2003) 416-425.
1736. Lazar M., *On the Higgs mechanism and stress functions in the translational gauge theory of dislocations*, *Phys. Lett.* **A373** (2009) 1578-1582.
1737. Lazar M., *Dislocation field theory in 2D: Application to graphene*, *Phys. Lett.* **A377** (2013) 423-429.
1738. Lazar M. and Hehl F.W., *Cartan's spiral staircase in physics and, in particular, in the gauge theory of dislocations*, *Found. Phys.* **40** (2010) 1298-1325.
1739. Le K.C. and Stumpf H., *On the determination of the crystal reference in nonlinear continuum theory of dislocations*, *Proc. Roy. Soc. Lond.* **A452** (1996) 359-371.
1740. Lebedev O., *Torsion constraints in the Randall-Sundrum scenario*, *Phys. Rev.* **D65** (2002) 124008 (7 pages).
1741. Lecian O.M. and Montani G., *Electroweak model within the framework of Lorentz gauge theory: Ashtekar variables?*, *Nuovo Cim.* **B122** (2007) 207-212.
1742. Lecian O.M. and Montani G., *Is torsion a fundamental physical field?*, in: *Proc. of the 11th Marcel Grossmann Meeting "On recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories"*, Berlin, Germany, 23-29 July 2006, eds. H. Kleinert, R.T. Jantzen, and R. Ruffini (World Scientific: Singapore, 2008) pt. C, pp. 2668-2670.
1743. Lecian O.M. and Montani G., *Riemannian and non-Riemannian extensions of geometrodynamics versus Einsteinian gravity*, *J. Korean Phys.Soc.* **56** (2010) 1653-1657.
1744. Lecian O.M., Montani G., and Carlevaro N., *Novel analysis of spinor interactions and non-Riemannian geometry*, *Eur. Phys. J. Plus* **128** (2013) 19.
1745. Leclerc M., *Teleparallel limit of Poincaré gauge theory*, *Phys. Rev.* **D71** (2005) 027503 (4 pages).

1746. Leclerc M., *One-parameter teleparallel limit of Poincaré gravity*, *Phys. Rev.* **D72** (2005) 044002 (8 pages).
1747. Leclerc M., *Mathisson-Papapetrou equations in metric and gauge theories of gravity in a Lagrangian formulation*, *Class. Quantum Grav.* **22** (2005) 3203.
1748. Leclerc M., *The Higgs sector of gravitational gauge theories*, *Ann. Phys. (USA)* **321** (2006) 708-743.
1749. Lee C., *A spherically symmetric electro-vacuum solution of the Poincaré gauge field theory of gravitation*, *Phys. Lett.* **B130** (1983) 257-259.
1750. Lee C.-Y., *Renormalization of quantum gravity with local  $GL(4, R)$  symmetry*, *Class. Quantum Grav.* **9** (1992) 2001-2020.
1751. Lee C.-Y. and Ne'eman Y., *Renormalization of gauge-affine gravity*, *Phys. Lett.* **B242** (1990) 59-63.
1752. Lee C.-Y. and Ne'eman Y., *BRST-transformations for an affine gauge model of gravity with local  $GL(4, R)$  symmetry*, *Phys. Lett.* **B233** (1990) 286-290.
1753. Lee Hac Won, *New algorithm for asymptotic expansion of the heat kernel*, *Phys. Rev.* **D35** (1987) 2440-2447.
1754. Lehnert R., Snow W.M., and Yan H., *A first experimental limit on in-matter torsion from neutron spin rotation in liquid  $^4\text{He}$* , *Phys. Lett.* **B730** (2014) 353-356.
1755. Lemke J., *Shock waves in the Poincaré gauge theory of gravitation*, *Phys. Lett.* **A143** (1990) 13-16.
1756. Lemke J. and Mielke E.W., *Gravitational moments of spin one-half particles and of topologically massive photons and gravitons in 2 + 1 dimensions*, *Phys. Lett.* **A175** (1993) 277-281.
1757. Lemke J., Ne'eman Y., and Pecina-Cruz J., *Wigner analysis and Casimir operators of  $SA(4, R)$* , *J. Math. Phys.* **33** (1992) 2256-2259.
1758. Lenzen H.-J., *On space-time models with axial torsion: Some vacuum solutions of the Poincaré gauge field theory of gravity*, *Nuovo Cim.* **B82** (1984) 85-89.
1759. Lenzen H.-J., *On spherically symmetric fields with dynamic torsion in gauge theories of gravitation*, *Gen. Relat. Grav.* **17** (1985) 1137-1151.
1760. Letelier P.S., *Ghost neutrinos in the Einstein-Cartan theory of gravitation*, *Phys. Lett.* **A54** (1975) 351-352.
1761. Letelier P.S., *On ghost neutrinos in the Einstein-Cartan theory of gravitation*, *Phys. Lett.* **A79** (1980) 283-284.
1762. Letelier P.S., *Spinning strings as torsion line spacetime defects*, *Class. Quantum Grav.* **12** (1995) 471-478.
1763. Letelier P.S., *Spacetime defects: torsion loops*, *Class. Quantum Grav.* **12** (1995) 2221-2223.
1764. Li B., Sotiriou T.P., and Barrow J.D.  *$f(T)$  gravity and local Lorentz invariance*, *Phys. Rev.* **D83** (2011) 064035 (5 pages).
1765. Li B., Sotiriou T.P., and Barrow J.D. *Large-scale structure in  $f(T)$  gravity*, *Phys. Rev.* **D83** (2011) 104017 (13 pages).
1766. Li J.-T., Lee C.-C., Geng C.-G., *Einstein static universe in exponential  $f(T)$  gravity*, *Eur. Phys. J.* **C73** (2013) 2315.
1767. Li S., *The topological structure of Nieh-Yan form and the chiral anomaly in spaces with torsion*, *J. Phys. A: Math. and Gen.* **A32** (1999) 7153-7162.
1768. Li Yuanjie, *Cosmic critical density modified by the torsion*, *Acta Astrophysica Sinica* **11** (1991) 1.

1769. Li Yuanjie, *Theory of fiber bundle with local metric of internal space and gravitation with torsion*, *Int. J. Theor. Phys.* **32** (1993) 1619-1625.
1770. Li Yuanjie, Luo Shi-Jun, and Ma Wei-Chan, *Inflation in Brans-Dicke theory with torsion*, *Int. J. Theor. Phys.* **32** (1993) 2173-2176.
1771. Li Yuanjie, Lin S.Y., and Luo Shi-Jun, *Inflation in Poincaré gauge gravitation theory*, *Int. J. Theor. Phys.* **34** (1995) 2003-2008.
1772. Li X.-Z., Sun C.-B., and Xi P., *Torsion cosmological dynamics*, *Phys. Rev. D* **79** (2009) 027301 (4 pages).
1773. Liang W.-C. and Lee S.-C., *Vorticity, gyroscopic precession, and spin-curvature force*, *Phys. Rev. D* **87** (2013) 044024 (11 pages).
1774. Lichnerowicz A., *Theories relativistes de la gravitation et de l'électromagnétisme. Relativité générale et théories unitaires* (Masson: Paris, 1955) 298 p.
1775. Liebscher D.-E., *Generalized equations of motion. I. The equivalence principle and non-Riemannian space-time*, *Ann. der Phys.* **30** (1973) 309-320.
1776. Liebscher D.-E., *Generalized equations of motion. II. The integration of generalized systems of dynamical equations*, *Ann. der Phys.* **30** (1973) 321-328.
1777. Liebscher D.-E., *The motion of test-particles in non-Riemannian space-time*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979), Eds. P.G.Bergmann and V.De Sabbata (Plenum: New York, 1980) 125-137.
1778. Lindsrom L., *A comment on the geometry of some scalar-tensor theories*, *Gen. Relat. Grav.* **18** (1986) 845-848.
1779. Lindsrom L. and Grundberg J., *A torsion potential*, *Nuovo Cim.* **B74** (1983) 181-184.
1780. Liu D. and Rebouças M.J., *Energy conditions bounds on  $f(T)$  gravity*, *Phys. Rev. D* **86** (2012) 083515 (8 pages).
1781. Liu Y.-F., *The dynamical effects of the torsion field with gravitation*, *Acta Astrophysica Sinica* **2** (1982) 320.
1782. Liu Y.-F., *Dynamical effect of torsion in gravitational field*, *Chinese Astronomy and Astrophysics* **7** (1983) 58.
1783. Liu Y.-F. and Tsou C.-L., *The Lorentz gauge field on the Riemannian space and its spherically symmetric static exact solutions*, *Acta Phys. Pol.* **B10** (1979) 893-899.
1784. Liu Y.-F. and Wu Yong-Shi, *The effects of torsion fields on spinning top*, *Acta Phys. Pol.* **B11** (1980) 173-183.
1785. Lledó M.A. and Sommovigo L., *Torsion formulation of gravity*, *Class. Quantum Grav.* **27** (2010) 065014.
1786. Lochak G., *Twisted space, chiral gauge and magnetism*, *Annales de la Fondation L. de Broglie* **32** (2007) 125-136.
1787. Lompay R.R., *On the energy-momentum and spin tensors in the Riemann-Cartan space*, *Gen. Rel. Grav.* **46** (2014) 1692.
1788. Lompay R.R. and Petrov A.N., *Covariant differential identities and conservation laws in metric-torsion theories of gravitation. I. General consideration*, *J. Math. Phys.* **54** (2013) 062504.
1789. Lompay R.R. and Petrov A.N., *Covariant differential identities and conservation laws in metric-torsion theories of gravitation. II. Manifestly generally covariant theories*, *J. Math. Phys.* **54** (2013) 102504.
1790. Lopez C., *Gravitational energy from a quadratic Lagrangian with torsion*, *Int. J. Theor. Phys.* **16** (1977) 167-176.

1791. López-Pinto A., Tiemblo A., and Tresguerres R., *Ordinary matter in non-linear affine gauge theories of gravitation*, *Class. Quantum Grav.* **12** (1995) 1503-1516.
1792. López-Pinto A., Tiemblo A., and Tresguerres R., *Nonlinear affine embedding of the Dirac field from the multiplicity-free  $SL(4, R)$  unirreps*, *Class. Quantum Grav.* **13** (1996) 2255-2265.
1793. López-Pinto A., Tiemblo A., and Tresguerres R., *Hamiltonian Poincaré gauge theory of gravitation*, *Class. Quantum Grav.* **14** (1997) 549-575.
1794. Lord E.A., *The metric-affine gravitational theory as the gauge theory of the affine group*, *Phys. Lett.* **A65** (1978) 1-4.
1795. Lord E.A., *Gauge theory of a group diffeomorphism. II. The conformal and de Sitter groups*, *J. Math. Phys.* **27** (1987) 3051-3054.
1796. Lord E.A., *Poincaré gauge theory from self-coupling*, *Pramana J. Phys.* **29** (1987) 359-368.
1797. Lord E.A., *A unified approach to the gauging of space-time and internal symmetries*, *Gen. Relat. Grav.* **19** (1987) 983-1002.
1798. Lord E.A., *Fiber bundles in gravitational theory*, in: "Gravitation and Cosmology, Proc. XV IAGRG Conf., North Bengal Univ., November 1989", Eds. S. Mukherjee et al (Wiley Eastern: New Delhi, 1992) 30-42.
1799. Lord E.A. and Goswami P., *Gauging the conformal group*, *Pramana* **25** (1985) 635-640.
1800. Lord E.A., and Goswami P., *Gauge theory of a group of diffeomorphisms I. General principles*, *J. Math. Phys.* **27** (1986) 2415-2422.
1801. Lord E.A. and Goswami P., *Gauge theory of a group of diffeomorphisms II. The conformal and de Sitter groups*, *J. Math. Phys.* **27** (1986) 3051-3054.
1802. Lord E.A. and Goswami P., *Gauge theory of a group of diffeomorphisms III. The fiber bundle description*, *J. Math. Phys.* **29** (1988) 258-267.
1803. Lord E.A. and Goswami P., *Poincaré gauge theory from self-coupling*, *Pramana* **29** (1987) 359-368.
1804. Lord E.A. and Sinha K.P., *Spin and mass content of linearized Poincaré gauge theories*, *Pramana J. Phys.* **30** (1988) 511-519.
1805. Lorenz D., *Einstein-Cartan-Maxwell-Bianchi cosmologies (I, VII<sub>0</sub>, VIII<sub>2</sub>, IX - types)*, *Acta Phys. Pol.* **B12** (1981) 939-950.
1806. Lorenz D., *Kantowski-Sachs cosmological solution with torsion*, *J. Phys.: Math. and Gen.* **A15** (1982) 2809-2812.
1807. Lorenz D., *Einstein-Cartan-Maxwell-Bianchi type II solutions*, *Acta Phys. Pol.* **B13** (1982) 195-199.
1808. Lorenz D., *Einstein-Cartan-Maxwell-Bianchi type V cosmological models*, *Gen. Relat. Grav.* **14** (1982) 691-702.
1809. Lu H.Q. and Cheng K.S., *A Bianchi type-V cosmological model with spin matter*, *Class. Quantum Grav.* **12** (1995) 2755-2765.
1810. Lu H.Q., Harko T., and Mak M.K., *Cosmic no-hair conjecture in Einstein-Cartan theory*, *Int. J. Mod. Phys.* **D10** (2001) 315-324.
1811. Lu J.-A., *Cosmological meaning of the gravitational gauge group*, *Gen. Rel. Grav.* **46** (2014) 1780.
1812. Lu J.-A.,  *$R + S^2$  theories of gravity without big-bang singularity*, *Ann. Phys. (USA)* **354** (2015) 424-430.
1813. Lu J.-A. and Chee G., *An analytic cosmology solution of Poincaré gauge gravity*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650096 (26 pages).

1814. Lu J.-A. and Huang C.-G., *Kaluza-Klein-type models of de Sitter and Poincaré gauge theories of gravity*, *Class. Quantum Grav.* **30** (2013) 145004.
1815. Lu J.S., Li X.Z., and Zhang G.Y., *The  $SU(11)$  family unified model and the thermal history of the very early universe*, *Nuovo Cim.* **A104** (1991) 1687-1695.
1816. Lubkin E., *Geometric definition of gauge invariance*, *Ann. Phys. (USA)* **23** (1963) 233-283.
1817. Lucat S. and Prokopec T., *The role of conformal symmetry in gravity and the standard model*, *Class. Quantum Grav.* **33** (2016) 245002 (24 pages).
1818. Luehr C.P., Rosenbaum M., Ryan M.P., Jr., and Shepley L.C., *Nonstandard vector connection given by nonstandard spinor connection*, *J. Math. Phys.* **18** (1977) 965-970.
1819. Luehr C.P. and Rosenbaum M., *Gravitation as an internal gauge theory of the Poincaré group*, *J. Math. Phys.* **21** (1980) 1432-1438.
1820. Luehr C.P. and Rosenbaum M.P., *Twistor bundles and gauge action principles of gravitation*, *J. Math. Phys.* **23** (1982) 1471-1488.
1821. Luo S.J., *Geometry of torsion*, *Int. J. Theor. Phys.* **34** (1995) 2009-2014.

## M

1822. Ma M.-S. and Huang C.-G., *On the weak field approximation of the de Sitter gauge theory of gravity*, *Gen. Rel. Grav.* **45** (2013) 143-153.
1823. Ma M.-S., Liu F., and Zhao R., *Continuous phase transition and critical behaviors of 3D black hole with torsion*, *Class. Quantum Grav.* **31** (2014) 095001.
1824. Ma M.-S. and Zhao R., *Phase transition and entropy spectrum of the BTZ black hole with torsion*, *Phys. Rev.* **D89** (2014) 044005 (9 pages).
1825. Ma W., Shao C., and Li Y., *Exact solution of Poincaré gauge field equations of gravity with torsion*, *Int. J. Theor. Phys.* **33** (1994) 1941-1947.
1826. Ma W., Shao C., Li Y., and Tian X., *An exact solution of gauge field equations of Poincaré gravity with torsion and spin current*, *Gen. Relat. Grav.* **27** (1995) 143-152.
1827. MacDowell S.W., and Mansouri F., *Unified geometric theory of gravity and supergravity*, *Phys. Rev. Lett.* **38** (1977) 739-742.
1828. Macías A., *Chiral ( $N = 1$ ) supergravity*, *Class. Quantum Grav.* **13** (1996) 3163-3174.
1829. Macías A., Lämmerzahl C., and García A., *A class of colliding waves in metric-affine gravity, nonmetricity and torsion shock waves*, *J. Math. Phys.* **41** (2000) 6369-6380.
1830. Macías A., Lämmerzahl C., and Pimentel L.O., *Matching conditions in metric-affine gravity*, *Phys. Rev.* **D66** (2002) 104013 (9 pages).
1831. Macías A., Mielke E.W., and Morales-Técotl H.A., *Gravitational-geometric phases and translations*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 227-242.
1832. Macías A., Mielke E.W., and Morales-Técotl H.A., *Projectively invariant metric-affine models of gravity*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 243-255.
1833. Macías A., Mielke E.W., Morales-Técotl H.A., and Tresguerres R., *Torsion and Weyl covector in metric-affine models of gravity*, *J. Math. Phys.* **36** (1995) 5868-5876.
1834. Macías A., Mielke E.W., and Socorro J., *Solitonic monopole solution in metric-affine gauge theory carrying Weyl charges*, *Class. Quantum Grav.* **15** (1998) 445-452.

1835. Macías A. and Socorro J., *Generalized Reissner-Nordstrom solution in metric-affine gravity*, *Class. Quantum Grav.* **16** (1999) 2323-2333.
1836. Magnano G., *Nonlinear Gravitational Lagrangians revisited*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1640006 (17 pages).
1837. Magnon A.M.R., *Spin plane defects and emergence of Planck's constant in gravity*, *J. Math. Phys.* **32** (1991) 928-931.
1838. Magueijo J., Złońnik T.G., and Kibble T.W.B., *Cosmology with a spin*, *Phys. Rev.* **D87** (2013) 063504 (13 pages).
1839. Magueijo J., Rodríguez-Vázquez M., Westman H., and Złońnik T., *Cosmological signature change in Cartan gravity with dynamical symmetry breaking*, *Phys. Rev.* **D89** (2014) 063542 (18 pages).
1840. Mahanta M.N. and Rao A.S., *On the equivalence of the Einstein-Mayer and Einstein-Cartan theories for describing a spinning medium*, *J. Math. Phys.* **16** (1975) 381-383.
1841. Mahato P., *Gravitational constant and torsion*, *Mod. Phys. Lett.* **A17** (2002) 1991.
1842. Mahato P., *Torsion, scalar field, mass and FRW cosmology*, *Int. J. Theor. Phys.* **44** (2005) 79-93.
1843. Mahato P., *Torsion, Dirac field, dark matter and dark radiation*, *Mod. Phys. Lett.* **A22** (2007) 835.
1844. Mahato P., *Torsion, scalar field and  $f(R)$  gravity*, *Annales de la Fondation L. de Broglie* **32** (2007) 297-310.
1845. Mahato P. and Bandyopadhyay P., *Stochastic geometry, torsion and  $N=1$  supergravity*, *Nuovo Cim.* **B98** (1987) 53-62.
1846. Mahato P. and Bhattacharya P., *Torsion, Chern-Simons term and diffeomorphism invariance*, *Mod. Phys. Lett.* **A26** (2011) 415.
1847. Maier R., *Static vacuum solutions in non-Riemannian gravity*, *Gen. Rel. Grav.* **46** (2014) 1830.
1848. Maier R. and Falciano F.T., *Brane world in non-Riemannian geometry*, *Phys. Rev.* **D83** (2011) 064019 (7 pages).
1849. Majewski E., *Spinors and torsion in a Riemann-Cartan approach to elasticity with a continuous defect distribution and analogies to the Einstein-Cartan theory of gravitation*, in: "Physics of Asymmetric Continuum: Extreme and Fracture Processes", Eds. Teisseyre R., Nagahama H., and Majewski E. (Springer: Berlin, 2008) 249-272.
1850. Majumdar P. and SenGupta S., *Parity-violating gravitational coupling of electromagnetic fields*, *Class. Quantum Grav.* **16** (1999) L89-L94.
1851. Makaruk H., *Tetrad, connection, and metric as independent variables in Lagrangians of micromorphic continua models*, *Int. J. Theor. Phys.* **35** (1996) 1461-1471.
1852. Maluf J.W., *Conformal invariance and torsion in general relativity*, *Gen. Relat. Grav.* **19** (1987) 57-71.
1853. Maluf J.W., *The Stephenson-Kilmister-Yang Hamiltonian*, *Class. Quantum Grav.* **5** (1988) L81-L85.
1854. Maluf J.W., *Energy spectrum of quadratic theories of gravitation*, *Class. Quantum Grav.* **6** (1989) 1189-1195.
1855. Maluf J.W., *Positivity of energy of  $R + R^2$  theories of gravitation*, *Class. Quantum Grav.* **6** (1989) L151-L154.
1856. Maluf J.W., *The constraint algebra of tetrad gravity reexamined*, *Class. Quantum Grav.* **8** (1991) 287-295.

1857. Maluf J.W., *On the absence of physical degrees of freedom in the Stephenson-Kilmister-Yang theory*, *J. Math. Phys.* **32** (1991) 1556-1559.
1858. Maluf J.W., *Self-dual connections, torsion and Ashtekar's variables*, *J. Math. Phys.* **33** (1992) 2849-2854.
1859. Maluf J.W., *Hamiltonian formulation of the teleparallel description of general relativity*, *J. Math. Phys.* **35** (1994) 335-343.
1860. Maluf J.W., *Teleparallelism and geometrodynamics*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 257-268.
1861. Maluf J.W., *The Hamiltonian constraint in the teleparallel equivalent of general relativity*, *Gen. Relat. Grav.* **28** (1996) 1361-1376.
1862. Maluf J.W., *On the distribution of gravitational energy in the de Sitter space*, *J. Math. Phys.* **37** (1996) 6293-6301.
1863. Maluf J.W., *Sparling two-forms, the conformal factor and the gravitational energy density of the teleparallel equivalent of general relativity*, *Gen. Relat. Grav.* **30** (1998) 413-423.
1864. Maluf J.W., *Dirac spinor fields in the teleparallel gravity: Comment on "Metric-affine approach to teleparallel gravity"*, *Phys. Rev.* **D67** (2003) 108501 (3 pages).
1865. Maluf J.W., *The teleparallel equivalent of general relativity*, *Ann. d. Phys. (Berlin)* **525** (2013) 339-357.
1866. Maluf J.W., *The teleparallel equivalent of General Relativity and the gravitational centre of mass*, *Universe* **2**(3) (2016) 19 [13 pages].
1867. Maluf J.W. and Faria F.F., *Conformally invariant teleparallel theories of gravity*, *Phys. Rev.* **D85** (2012) 027502 (5 pages).
1868. Maluf J.W. and Faria F.F., *Teleparallel gauge theory of gravity*, *Ann. d. Phys. (Berlin)* **524** (2012) 366-370.
1869. Maluf J.W., Faria F.F., and Castello-Branco K.H., *The gravitational energymomentum flux*, *Class. Quantum Grav.* **20** (2003) 4683.
1870. Maluf J.W., Faria F.F., and Ulhoa S.C., *On reference frames in spacetime and gravitational energy in freely falling frames*, *Class. Quantum Grav.* **24** (2007) 2743.
1871. Maluf J.W., Faria F.F., Ulhoa S.C., and da Rocha-Neto J.F. *The angular momentum of the gravitational field and the Poincaré group*, *Class. Quantum Grav.* **23** (2006) 6245.
1872. Maluf J.W. and Goya A., *Space-time defects and teleparallelism*, *Class. Quantum Grav.* **18** (2001) 5143.
1873. Maluf J.W. and Kneip A., *Gravitational energy of conical defects*, *J. Math. Phys.* **38** (1997) 458-465.
1874. Maluf J.W., Martins E.F., and Kneip A., *Gravitational energy of rotating black holes*, *J. Math. Phys.* **37** (1996) 6302-6310.
1875. Maluf J.W. and Santos-Silva G., *Stability of  $R + T^2$  theories of gravitation*, *Class. Quantum Grav.* **25** (1993) 653-661.
1876. Maluf J.W. and da Rocha-Neto J.F., *General relativity on a null surface: Hamiltonian formulation in the teleparallel geometry*, *Gen. Relat. Grav.* **31** (1999) 173-185.
1877. Maluf J.W. and da Rocha-Neto J.F., *Static Bondi energy in the teleparallel equivalent of general relativity*, *J. Math. Phys.* **40** (1999) 1490-1503.
1878. Maluf J.W. and da Rocha-Neto J.F., *Hamiltonian formulation of general relativity in the teleparallel geometry*, *Phys. Rev.* **D64** (2001) 084014 (8 pages).

1879. Maluf J.W., da Rocha-Neto J.F., Toribio T.M.L., and Castello-Branco K.H., *Energy and angular momentum of the gravitational field in the teleparallel geometry*, *Phys. Rev.* **D65** (2002) 124001 (12 pages).
1880. Maluf J.W. and Ulhoa S.C., *The energy-momentum of plane-fronted gravitational waves in the teleparallel equivalent of GR*, *Phys. Rev.* **D78** (2008) 047502 (4 pages).
1881. Maluf J.W. and Ulhoa S.C., *On the gravitational angular momentum of rotating sources*, *Gen. Rel. Grav.* **41** (2009) 1233-1247.
1882. Maluf J.W., Ulhoa S.C., and Faria F.F., *Pound-Rebka experiment and torsion in the Schwarzschild spacetime*, *Phys. Rev.* **D80** (2009) 044036 (6 pages).
1883. Maluf J.W., Ulhoa S.C., and da Rocha-Neto J.F., *Gravitational pressure on event horizons and thermodynamics in the teleparallel framework*, *Phys. Rev.* **D85** (2012) 044050 (10 pages).
1884. Maluf J.W., Ulhoa S.C., and da Rocha-Neto J.F., *Gravitational waves and the breaking of parallelograms in space-time*, *Gen. Rel. Grav.* **45** (2013) 1163-1177.
1885. Maluf J.W., Veiga M.V.O., and da Rocha-Neto J.F., *Regularized expression for the gravitational energy-momentum in teleparallel gravity and the principle of equivalence*, *Gen. Rel. Grav.* **39** (2007) 227-240.
1886. Malyshev C., *Underlying algebraic and gauge structures of the theory of disclinations*, *Arch. Mech. (Warsaw)* **45** (1993) 93-105.
1887. Manna B., Sinha S., and Sahoo S., *Exact solution for a nonstatic cylindrically symmetric perfect fluid distribution in Einstein-Cartan theory*, *Gravitation and Cosmology* **22** (2016) 44-47.
1888. Manna B., Sinha S., and Sahoo S., *Extension of van Stockum cylinder in Einstein-Cartan theory*, *Can. J. Phys.* **94** (2016) 1061-1063.
1889. Manna B., Sinha S., and Sahoo S., *Rotating spin-polarized cylinder in Einstein-Cartan theory*, *Gravitation and Cosmology* **23** (2017) 90-94.
1890. Mannheim P.D., *Conformal invariance and the metrification of the fundamental forces*, *Int. J. Mod. Phys.* **D25** (2016) 1644003 (6 pages)
1891. Manoff S., *Invariant projection of the energy-momentum tensor for field theories in spaces with affine connection and metric*, *J. Math. Phys.* **32** (1991) 728-734.
1892. Manoff S., *Geodesic and autoparallel equations over differentiable manifolds*, *Int. J. Mod. Phys.* **A11** (1996) 3849-3874.
1893. Manoff S., *Fermi derivative and Fermi-Walker transports over  $(L_n, g)$ -spaces*, *Class. Quantum Grav.* **15** (1998) 465-477.
1894. Manoff S., *Fermi derivative and Fermi-Walker transports over  $(\overline{L}_n, g)$ -spaces*, *Int. J. Mod. Phys.* **A13** (1998) 4289-4308.
1895. Manoff S., *Spaces with contravariant and covariant affine connections and metrics*, *Part. Nucl.* **30** (1999) 517-549 [*Fiz. Elem. Chast. Atom. Yadra* **30** (1999) 1211-1269 (in Russian)].
1896. Manoff S., *Lagrangian theory of tensor fields over spaces with contravariant and covariant affine connections and metrics and its applications to Einstein's theory of gravitation in  $\overline{V}_n$ -spaces*, *Acta Appl. Math.* **55** (1999) 51-125.
1897. Manoff S., *Conformal derivative and conformal transports over  $(\overline{L}_n, g)$ -spaces*, *Int. J. Mod. Phys.* **A15** (2000) 679-696.
1898. Manoff S., *Auto-parallel equation as Euler-Lagrange's equation in spaces with affine connections and metrics*, *Gen. Relat. Grav.* **32** (2000) 1559-1582.
1899. Manoff S., *Frames of reference in spaces with affine connections and metrics*, *Class. Quantum Grav.* **18** (2001) 1111-1125.

1900. Manoff S., *Deviation equations of Synge and Schild over  $(\overline{L}_n, g)$ -spaces*, *Int. J. Mod. Phys. A* **16** (2001) 1109-1122.
1901. Manoff S., *Deviation operator and deviation equations over spaces with affine connections and metrics*, *J. Geom. Phys.* **39** (2001) 337-350.
1902. Mansouri F., *Conformal gravity as a gauge theory*, *Phys. Rev. Lett.* **42** (1979) 1021-1024.
1903. Mansouri F. and Chang L.N., *Gravitation as a gauge theory*, *Phys. Rev.* **D13** (1976) 3192-3200.
1904. Mao Y., Tegmark M., Guth A.H., and Cabi S., *Constraining torsion with Gravity Probe B*, *Phys. Rev.* **D76** (2007) 104029 (26 pages).
1905. March R., Bellettini G., Tauraso R., and Dell'Agnello S., *Constraining spacetime torsion with the Moon and Mercury*, *Phys. Rev.* **D83** (2011) 104008 (19 pages).
1906. Mardones A. and Zanelly J., *Lovelock-Cartan theory of gravity*, *Class. Quantum Grav.* **8** (1991) 1545-1558.
1907. Maroto A.L. and Shapiro I.L., *On the inflationary solutions in higher-derivative gravity with dilaton field*, *Phys. Lett.* **B414** (1997) 34-44.
1908. Martellini M., *Quantum gravity in the Eddington purely affine picture*, *Phys. Rev.* **D29** (1984) 2746-2755.
1909. Mashhoon B. and Obukhov Yu.N., *Spin precession in inertial and gravitational fields*, *Phys. Rev.* **D88** (2013) 064037 (10 pages).
1910. Massa E. and Vignolo S., *Variational techniques in general relativity: A metric-affine approach to Kaluza's theory*, *J. Math. Phys.* **48** (2007) 022501.
1911. Maulbetsch C. and Shabanov S.V., *The inverse variational problem for autoparallels*, *J. Phys. A: Math. and Gen.* **A32** (1999) 5355-5366.
1912. Mavromatos N.E. and Pilaftsis A., *Anomalous Majorana neutrino masses from torsionful quantum gravity*, *Phys. Rev.* **D86** (2012) 124038 (8 pages).
1913. McCrea J.D., *Static vacuum cylindrical and plane symmetrical solution of the quadratic Poincaré gauge field equation*, *J. Phys.* **A16** (1983) 997-1004.
1914. McCrea J.D., *The use of REDUCE in finding exact solutions of the quadratic Poincaré gauge field equations*, in: "Classical General Relativity", Eds. W.B. Bonnor, J.N. Islam, and M.A.H. MacCallum (Cambridge Univ. Press: Cambridge, 1984) 173-182.
1915. McCrea J.D., *A NUT-like solution of the quadratic Poincaré gauge field equations*, *Phys. Lett.* **A100** (1984) 397-399.
1916. McCrea J.D., *Poincaré gauge theory of gravitation: foundations, exact solutions and computer algebra*, in: "Proc. of the 14th Intern. Conf. on Differential Geometric Methods in Mathematical Physics, Salamanca, 1985", Eds. P.L. Garcia and A. Pérez-Rendón, *Lect. Notes Math.* **1251** (1987) 222-237.
1917. McCrea J.D., *Irreducible decompositions of non-metricity, torsion, curvature and Bianchi identities in metric-affine spacetimes*, *Class. Quantum Grav.* **9** (1992) 553-568.
1918. McCrea J.D., *REDUCE in general relativity and Poincaré gauge theory*, in: "Algebraic computing in general relativity", *Lect. Notes of the 1st Brasil. School on Comp. Algebra, Rio de Janeiro, July-August 1989*, Eds. M.J. Rebouças and W.L. Roque (Clarendon Press: Oxford, 1994) 173-263.
1919. McCrea J.D., Baekler P., and Gürses M., *A Kerr-like solution of the Poincaré gauge field equations*, *Nuovo Cim.* **B99** (1987) 171-177.
1920. McCrea J.D., Hehl F.W., and Mielke E.W., *Mapping Noether identities into Bianchi identities in general relativistic theories of gravity and in the field theory of static lattice defects*, *Int. J. Theor. Phys.* **29** (1990) 1185-1206.

1921. McCrea J.D., Mielke E.W., and Hehl F.W., *A remark on the axisymmetric Chen et al. solution of the Poincaré gauge theory*, *Phys. Lett.* **A127** (1988) 65-69.
1922. McInnes B.T., *On the significance of compatibility condition in gauge theories of Poincaré group*, *Class. Quantum Grav.* **1** (1984) 1-5.
1923. McInnes B.T., *On the geometrical interpretation of nonsymmetric space-time field structures*, *Class. Quantum Grav.* **1** (1984) 105-113.
1924. McInnes B.T., *On the affine approach to Riemann-Cartan space-time geometry*, *Class. Quantum Grav.* **1** (1984) 115-123.
1925. McInnes B., *Spontaneous compactification and Ricci-flat manifolds with torsion*, *J. Math. Phys.* **27** (1986) 2029-2038.
1926. McKellar R.J., *The uniqueness of gravity as Poincaré or Lorentz gauge theory*, *J. Math. Phys.* **22** (1981) 2934-2942.
1927. McKellar R.J., *A gravitational Poincaré gauge theory and Higgs mechanism*, *J. Math. Phys.* **25** (1984) 161-166.
1928. Mecklenburg W., *Massive photons and the equivalence principle*, *Lett. Nuovo Cim.* **31** (1981) 616-618.
1929. Mehdizadeh M.R., and Ziaie A.H., *Einstein-Cartan wormhole solutions*, *Phys. Rev.* **D95** (2017) 064049 (12 pages).
1930. Mehra A.L. and Gokhroo M.K., *Static dust sphere in Einstein-Cartan theory*, *Gen. Relat. Grav.* **24** (1992) 1011-1014.
1931. Mena M. and Tahiri M., *A note on topological 4-D gravity and its metric approach*, *Mod. Phys. Lett.* **A15** (2000) 565-571.
1932. Menotti P. and Pelissetto A., *Poincaré, de Sitter and conformal gravity on the lattice*, *Phys. Rev.* **D35** (1987) 1194-1204.
1933. Mercuri S., *From the Einstein-Cartan to the Ashtekar-Barbero canonical constraints, passing through the Nieh-Yan functional*, *Phys. Rev.* **D77** (2008) 024036 (5 pages).
1934. Messios N., *Spinning particles in spacetimes with torsion*, *Int. J. Theor. Phys.* **46** (2007) 562-575.
1935. Mészáros A., *Did Eötvös torsion experiment detect Cartan's contortion?*, *Astrophys. and Space Sci.* **125** (1986) 405-410.
1936. Mészáros A., *Two notes on Cartan's contortion*, *Astrophys. and Space Sci.* **132** (1987) 415-417.
1937. Mészáros A., *Cartan's contortion as a pair of massless spin-2 fields*, *Phys. Rev.* **D35** (1987) 1176-1180.
1938. Mészáros A., *A repulsive interaction mediated by a spin-2 field?*, *Astrophys. and Space Sci.* **158** (1989) 163-167.
1939. Mészáros A., *Cartan's contorsion as a system of spin-2 and spin-0 fields*, *Gen. Relat. Grav.* **22** (1990) 417-430.
1940. Meyer H., *Moller's tetrad theory of gravitation as a special case of Poincaré theory - a coincidence?*, *Gen. Relat. Grav.* **14** (1982) 531-547.
1941. Miao R.-X., Li M., and Miao Y.-G., *Violation of the first law of black hole thermodynamics in  $f(T)$  gravity*, *JCAP* **11** (2011) 033.
1942. Mielke E.W., *Reduction of the Poincaré gauge field equations by means of duality rotations*, *J. Math. Phys.* **25** (1984) 663-668.
1943. Mielke E.W., *On pseudoparticle solutions in the Poincaré gauge theory of gravity*, *Fortschr. Phys.* **32** (1984) 639-660.

1944. Mielke E.W., *Analogue of the Witten effect in the Poincaré gauge theory of gravity*, *Phys. Lett.* **A110** (1985) 87-91.
1945. Mielke E.W., *Geometrodynamics of gauge fields - On the geometry of Yang-Mills and gravitational gauge theories* (Akademie Verlag: Berlin, 1987) 242 pp.
1946. Mielke E.W., *Algebraic resolution of the duality ansatz for Poincaré gauge fields*, *Phys. Lett.* **A128** (1988) 251-255.
1947. Mielke E.W., *Generating function for new variables in general relativity and Poincaré gauge theory*, *Phys. Lett.* **A149** (1990) 345-350; (E) *Phys. Lett.* **A151** (1990) 567.
1948. Mielke E.W., *Positive gravitational energy proof from complex variables?*, *Phys. Rev.* **D42** (1990) 3388-3394.
1949. Mielke E.W., *Ashtekar's complex variables in general relativity and its teleparallelism equivalent*, *Ann. Phys. (USA)* **219** (1992) 78-108.
1950. Mielke E.W., *Chern-Simons solution of the Ashtekar constraints for the teleparallelism equivalent of gravity*, *Acta Phys. Pol.* **29** (1998) 871-880.
1951. Mielke E.W., *Anomaly-free solution of the Ashtekar constraints for the teleparallelism equivalent of gravity*, *Phys. Lett.* **A251** (1999) 349-353.
1952. Mielke E.W., *Affine generalization of the Komar complex of general relativity*, *Phys. Rev.* **D63** (2001) 044018 (4 pages).
1953. Mielke E.W., *Beautiful gauge field equations in Cliffords*, *Int. J. Theor. Phys.* **40** (2001) 171-189.
1954. Mielke E.W., *Chern-Simons solutions of the chiral teleparallelism constraints of gravity*, *Nucl. Phys.* **B622** (2002) 457-471.
1955. Mielke E.W., *Consistent coupling to Dirac fields in teleparallelism: Comment on "Metric-affine approach to teleparallel gravity"*, *Phys. Rev.* **D69** (2004) 128501 (3 pages).
1956. Mielke E.W., *Einsteinian gravity from a topological action*, *Gen. Rel. Grav.* **40** (2008) 1311-1325.
1957. Mielke E.W., *Limitations on the topological BF scheme in Riemann-Cartan spacetime with torsion*, *Phys. Rev.* **D80** (2009) 047504 (4 pages).
1958. Mielke E.W., *Topologically modified teleparallelism, passing through the Nieh-Yan functional*, *Phys. Rev.* **D80** (2009) 067502 (4 pages).
1959. Mielke E.W., *Spontaneously broken topological  $SL(5, R)$  gauge theory with standard gravity emerging*, *Phys. Rev.* **D83** (2011) 044004 (9 pages).
1960. Mielke E.W., *Weak equivalence principle from a spontaneously broken gauge theory of gravity*, *Phys. Lett.* **B702** (2011) 187-190.
1961. Mielke E.W., *Einstein-Weyl gravity from a topological  $SL(5, R)$  gauge invariant action*, *Advances in Applied Clifford Algebras* **22** (2012) 803-817.
1962. Mielke E.W., *Symmetry breaking in topological quantum gravity*, *Int. J. Mod. Phys.* **D22** (2013) 1330009 (16 pages).
1963. Mielke E.W., *Is Einstein-Cartan theory coupled to light fermions asymptotically safe?*, *J. of Gravity* **2013** (2013) 812962 (5 pages).
1964. Mielke E.W., *Mass gap in Yang's theory of gravity*, *Int. J. Mod. Phys.* **D24** (2015) 1550070 (9 pages).
1965. Mielke E.W., *Geometrodynamics of Gauge Fields - On the Geometry of Yang-Mills and Gravitational Gauge Theories* (Springer: Berlin, 2017) 373 pp.
1966. Mielke E.W. and Baekler P., *Topological gauge model of gravity with torsion*, *Phys. Lett.* **A156** (1991) 399-403.

1967. Mielke E.W., Baekler P., Hehl F.W., Macías A., and Morales-Técotl H.A., *Yang-Mills-Clifford version of the chiral Einstein action*, in: "Gravity, Particles and Space-time", Eds. P. Pronin and G. Sardanashvily (World Scientific: Singapore, 1996) 217-254.
1968. Mielke E.W., Gronwald F., Obukhov Yu.N., Tresguerres R., and Hehl F.W., *Towards complete integrability of two-dimensional Poincaré gauge gravity*, *Phys. Rev.* **D48** (1993) 3648-3662.
1969. Mielke E.W. and Hehl F.W., *Die Entwicklung der Eichtheorien: Marginalien zu deren Wissenschaftsgeschichte*, in: "Exact sciences and their philosophical foundations – Vorträge des Internat. Hermann-Weyl-Kongresses, Kiel, 1985", Eds. W. Deppert, K. Hübner, A. Oberschelp, and V. Weidemann (P.Lang Verlag: Frankfurt a. M., 1988) 191.
1970. Mielke E.W., Hehl F.W., and McCrea J.D., *Belinfante type invariance of the Noether identities in a Riemannian and Weitzenbock space-time*, *Phys. Lett.* **A140** (1989) 368-372.
1971. Mielke E.W. and Kreimer D., *Chiral anomaly in Ashtekar's approach to canonical gravity*, *Int. J. Mod. Phys.* **D7** (1998) 535-548.
1972. Mielke E.W. and Kreimer D., *Chiral anomaly in contorted spacetimes*, *Gen. Relat. Grav.* **31** (1999) 701-712.
1973. Mielke E.W. and Macías A., *Chiral supergravity and anomalies*, *Ann. Phys. (Leipzig)* **8** (1999) 301-317.
1974. Mielke E.W., Macías A., and Morales-Técotl H.A., *Chiral fermions coupled to chiral gravity*, *Phys. Lett.* **A215** (1996) 14-20.
1975. Mielke, E.W., Macías A., and Morales-Técotl H.A., *Riemann-Cartan structure of the spatial Yang-Mills equation*, in: "Recent Developments in Gravitation and Mathematical Physics", *Proceedings of the First Mexican School on Gravitation and Mathematical Physics, Guanajuato, Mexico 12 – 16 Dec. 1994*, Eds. A. Macías, T. Matos, O. Obregón and H. Quevedo (World Scientific: Singapore, 1995) 128-135.
1976. Mielke, E.W., Macías A., and Ne'eman Y., *CP-symmetry in chiral gravity*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories" (Jerusalem, 22-27 June 1997)*, Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. B, pp. 901-903.
1977. Mielke E.M. and Maggiolo A.A.R., *Rotating black hole solution in a generalized topological 3D gravity with torsion*, *Phys. Rev.* **D68** (2003) 104026 (7 pages).
1978. Mielke E.M. and Maggiolo A.A.R., *Algebra for a BRST quantization of metric-affine gravity*, *Gen. Rel. Grav.* **35** (2003) 771-789.
1979. Mielke E.M. and Maggiolo A.A.R., *Current status of Yang's theory of gravity*, *Annales de la Fondation Louis de Broglie* **29**, n.2 (2004) 1-15.
1980. Mielke E.M. and Maggiolo A.A.R., *Duality in Yang's theory of gravity*, *Gen. Rel. Grav.* **37** (2005) 997-1007.
1981. Mielke E.M. and Maggiolo A.A.R., *S-duality in 3D gravity with torsion*, *Ann. Phys. (USA)* **322** (2007) 341-362.
1982. Mielke E.W., McCrea J.D., Ne'eman Y., and Hehl F.W., *Avoiding degenerate coframes in an affine gauge approach to quantum gravity*, *Phys. Rev.* **D48** (1993) 673-679.
1983. Mielke E.M., Obukhov Yu.N., and Hehl F.W., *Yang-Mills configurations from 3D Riemann-Cartan geometry*, *Phys. Lett.* **A192** (1994) 153-162.
1984. Mielke E.W. and Romero E.S., *Cosmological evolution of a torsion-induced quintaxion*, *Phys. Rev.* **D73** (2006) 043521 (10 pages).
1985. Mielke E.W. and Wallner R.P., *Mass and spin of double dual solutions in Poincaré gauge theory*, *Nuovo Cim.* **B101** (1988) 607-624.
1986. Mignani R. and Scipioni R., *On the solutions of the vacuum Cartan equation in metric-affine gravity*, *Gen. Relat. Grav.* **33** (2001) 683-711.

1987. Mignemi S., *Lineal gravity with dynamical torsion*, *Mod. Phys. Lett.* **A11** (1996) 1235-1245.
1988. Mignemi S., *Black hole solutions of two-dimensional Riemann-Cartan gravity with higher-derivative action*, *Ann. Phys.* **257** (1997) 1-17.
1989. Mignemi S., *Black hole solutions in four-dimensional gauge theories of gravity*, *Class. Quantum Grav.* **14** (1997) 2157-2166.
1990. Mignemi S., *Two-dimensional extended Poincaré gravity with nontrivial potential*, *Int. J. Mod. Phys.* **A12** (1997) 5767-5774.
1991. Mignemi S., *Spherically symmetric solutions in four-dimensional Poincaré gravity with non-trivial torsion*, *Class. Quantum Grav.* **15** (1998) 299-307.
1992. Mignemi S., *Two-dimensional gravity with an invariant energy scale*, *Phys. Lett.* **B551** (2003) 169-172.
1993. Mikhail F.I., Wanas M.I., and Nashed G.G.L., *Cosmological application of the new general relativity*, *Astrophys. Space Sci.* **228** (1995) 255-271.
1994. Mikhailov A.L., *Centrally-symmetric gravitational source on the background of nonmetricity*, *Moscow University Physics Bulletin*, **30**, n. 4 (1989) 83-84 (in Russian).
1995. Minkevich A.V., *Gravitational field and the local invariance principle*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 4 (1966) 117-118 (in Russian).
1996. Minkevich A.V., *Generalized cosmological Friedmann equations without gravitational singularity*, *Phys. Lett.* **A80** (1980) 232-234.
1997. Minkevich A.V., *Gravitation and limit of mass density*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 2 (1980) 87-94 (in Russian).
1998. Minkevich A.V., *Generalized cosmological Friedman equation and de Sitter solution*, *Phys. Lett.* **A95** (1983) 422-424.
1999. Minkevich A.V., *Vacuum and nonsingular inflationary universe*, *Dokl. AN BSSR* **29** (1985) 998-1001 (in Russian).
2000. Minkevich A.V., *Minimal quadratic gauge gravitational theory and non-linearity of wave fields with spin*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 5 (1986) 100-103 (in Russian).
2001. Minkevich A.V., *Gravitating systems in extremal conditions and gauge theories of gravity*, *Vesci AN Belarus, ser. fiz.-mat. nauk*, n. 4 (1995) 123-129 (in Russian).
2002. Minkevich A.V., *Problem of cosmological singularity and gauge theories of gravitation*, *Acta Phys. Pol.* **B29** (1998) 949-960.
2003. Minkevich A.V., *Regular inflationary cosmological models in general relativity theory and gauge theories of gravitation*, *Int. J. Mod. Phys.* **A17** (2002) 4441-4449.
2004. Minkevich A.V., *Problem of cosmological singularity and its possible resolution*, *Int. J. Mod. Phys.* **A20** (2005) 2436-2442.
2005. Minkevich A.V., *Gauge approach to gravitation and regular Big Bang theory*, *Grav. Cosmol.* **12** (2006) 11-20.
2006. Minkevich A.V., *Gravitation, cosmology and space-time torsion*, *Annales de la Fondation L. de Broglie* **32** (2007) 253-265.
2007. Minkevich A.V., *On gravitational repulsion effect at extreme conditions in gauge theories of gravity*, *Acta Phys. Pol.* **B38** (2007) 61-72.
2008. Minkevich A.V., *Gravitational interaction and Poincaré gauge theory of gravity*, *Acta Phys. Pol.* **B40** (2009) 229-239.
2009. Minkevich A.V., *Accelerating Universe with spacetime torsion but without dark matter and dark energy*, *Phys. Lett.* **B678** (2009) 423-426.

2010. Minkevich A.V., *De Sitter spacetime with torsion as physical spacetime in the vacuum and isotropic cosmology*, *Mod. Phys. Lett.* **A26** (2011) 259-266.
2011. Minkevich A.V., *Limiting energy density and a regular accelerating universe in Riemann-Cartan spacetime*, *JETP Lett.* **94** (2012) 831-836.
2012. Minkevich A.V., *On theory of regular accelerating universe in Riemann-Cartan spacetime*, *Mod. Phys. Lett.* **A28** (2013) 1350090 (12 pages).
2013. Minkevich A.V., *Gauge gravitation theory in Riemann-Cartan space-time and gravitational interaction*, *Gravitation and Cosmology* **22** (2016) 148-158.
2014. Minkevich A.V., *Towards the theory of regular accelerating Universe in Riemann-Cartan space-time*, *Int. J. Mod. Phys.* **A31** (2016) 1641011 (10 pages).
2015. Minkevich A.V. and Fedorov F.I., *On the Lagrangian formalism in field theory*, *Acta. Phys. Pol.* **B11** (1980) 367-377.
2016. Minkevich A.V. and Fedorov F.I., *Gauge approach in gravity and relativistic dynamics of classical systems*, in: "Fundamental Interactions", Ed. V.N.Ponomarev (Pedagogical Inst. Press: Moscow, 1984) 141-159 (in Russian).
2017. Minkevich A.V. and Fedorov F.I., *On conformal invariant gravitating systems in the Poincaré gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 5 (1985) 45-52 (in Russian).
2018. Minkevich A.V., Fedorov F.I., and Kudin V.I., *To the definition of space-time symmetries in the tetrad field theory*, *Dokl. AN BSSR* **25** (1981) 406-408 (in Russian).
2019. Minkevich A.V. and Garkun A.S., *Homogeneous isotropic models in the metric-affine gauge theory of gravity*, *Class. Quantum Grav.* **17** (2000) 3045-3054.
2020. Minkevich A.V. and Garkun A.S., *Analysis of inflationary cosmological models in gauge theories of gravitation*, *Class. Quantum Grav.* **23** (2006) 4237-4247.
2021. Minkevich A.V., Garkun A.S., and Kudin V.I. *Regular accelerating universe without dark energy in Poincaré gauge theory of gravity*, *Class. Quantum Grav.* **24** (2007) 5835-5847.
2022. Minkevich A.V., Garkun A.S., and Kudin V.I. *On some physical aspects of isotropic cosmology in Riemann-Cartan spacetime*, *JCAP* **03** (2013) 040 (15 pages).
2023. Minkevich A.V. and Karakura F.J., *On the relativistic dynamics of spinning matter in space-time with curvature torsion*, *J. Phys.* **A16** (1983) 1409-1418.
2024. Minkevich A.V. and Kudin V.I., *Gauge fields and the Lorentz group*, *Acta Phys. Pol.* **B5** (1974) 335-343.
2025. Minkevich A.V. and Kudin V.I., *The gauge A-field and the strong gravity*, *Acta Phys. Pol.* **B6** (1975) 661-666.
2026. Minkevich A.V. and Kudin V.I., *Analogue of interior Schwarzschild solution in the gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 3 (1981) 103-113 (in Russian).
2027. Minkevich A.V. and Nemenman I.M., *On the influence of the gravitating vacuum on the dynamics of homogeneous isotropic models in gauge theories of gravity*, *Class. Quantum Grav.* **12** (1995) 1259-1265.
2028. Minkevich A.V. and Nemenman I.M., *On the influence of the gravitating vacuum on the dynamics of homogeneous isotropic models in gauge theories of gravity*, *Dokl. AN Belarus* **39** (1995) 45-51 (in Russian).
2029. Minkevich A.V. and Nguen Hong Chuong, *On the equations of gauge gravitational theory for the ideal fluid ball*, *Dokl. AN BSSR* **27** (1983) 38-41 (in Russian).
2030. Minkevich A.V. and Nguyen Hong Chuong, *Qualitative analysis of isotropic models with viscosity in gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 4 (1989) 84-89 (in Russian).

2031. Minkevich A.V. and Nguyen Hong Chuong, *Gravitational vacuum and the isotropic models dynamics in gauge gravitational theory*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 5 (1989) 63-68 (in Russian).
2032. Minkevich A.V., Nguen Hong Chuong, and Fedorov F.I., *On the particles creation effect in isotropic models of Universe with torsion*, *Dokl. AN BSSR* **28** (1984) 211-214 (in Russian).
2033. Minkevich A.V., Nguen Hong Chuong, and Fedorov F.I., *On the spontaneous breakdown of gauge symmetry in isotropic cosmological models with torsion*, *Dokl. AN BSSR* **29** (1985) 697-699 (in Russian).
2034. Minkevich A.V., Nguen Hong Chuong, and Fedorov F.I., *Regular homogeneous anisotropic cosmological solution in the minimal quadratic gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 1 (1987) 98-102 (in Russian).
2035. Minkevich A.V. and Polozov M.N., *On spherically symmetric gravitational fields in the gauge theory of gravity in the Weyl-Cartan space*, *Vesci AN Belarus, ser. fiz.-mat. nauk*, n. 5 (1986) 80-83 (in Russian).
2036. Minkevich A.V. and Vereshchagin G.V., *On qualitative analysis of homogeneous isotropic models in gauge theories of gravity*, *Grav. Cosmol.* **8** (2002) 305-308.
2037. Minkowski P., *On the cosmological equations in the presence of a spatially homogeneous torsion field*, *Phys. Lett.* **B173** (1986) 247-250.
2038. Minn Hokee, *Global theory of Einstein-Cartan equations: Gödel universe with torsion*, *J. Korean Astron. Soc.* **9** (1976) 9-14.
2039. Miri M. and Rivier N., *Continuum elasticity with topological defects, including dislocations and extra-matter*, *J. Phys. A: Math. Gen.* **35** (2002) 1727.
2040. Mistura L., *Cartan connection and defects in Bravais lattices*, *Int. J. Theor. Phys.* **29** (1990) 1207-1218.
2041. Miyamoto S., Nakano T., Ohtani T., and Tamura Y., *Linear approximation for the Lorentz gauge field*, *Progr. Theor. Phys.* **66** (1981) 481-497.
2042. Miyamoto S., Nakano T., Ohtani T., and Tamura Y., *Linear approximation for massless Lorentz gauge field*, *Progr. Theor. Phys.* **69** (1983) 1236-1240.
2043. Modanese G. and Toller M., *Radial gauge in Poincaré gauge field theories*, *J. Math. Phys.* **31** (1990) 452-458.
2044. Mohanty S. and Sarkar U., *Constraints on background torsion field from K-physics*, *Phys. Lett.* **B433** (1998) 424-428.
2045. Mohseni M., *Torsion-induced spin precession*, *Eur. Phys. J.* **C56** (2008) 607-612.
2046. Mol I., *The non-metricity formulation of general relativity*, *Advances in Applied Clifford Algebras* **27** (2017) 1-32.
2047. Monteiro W. S.Jr. and Tomimura N.A., *Existence and causality of cylindrically symmetric cosmological models with rotating spin fluids*, *Class. Quantum Grav.* **8** (1991) 977-984.
2048. Monteiro W. S.Jr. and Tomimura N.A., *Existence and causality of cosmological models with spinning fluids and torsion*, *Phys. Lett.* **A197** (1995) 267-274.
2049. Montesinos M., *Connection with torsion from Ashtekar-Barbero connection*, *J. Math. Phys.* **40** (1999) 4084-4088.
2050. Moon T., Lee J., and Oh P., *Conformal invariance in Einstein-Cartan-Weyl space*, *Mod. Phys. Lett.* **A25** (2010) 3129-3143.
2051. Moraes F., *Geodesics around a dislocation*, *Phys. Lett.* **A214** (1996) 189-192.
2052. Moraes F., *Condensed matter physics as a laboratory for gravitation and cosmology*, *Brazilian J. Phys.* **30** (2000) 304-308.

2053. Morales-Técotl H.A. and Esposito G., *Self-dual action for fermionic fields and gravitation*, *Nuovo Cim.* **B109** (1994) 973-982.
2054. Morales-Técotl H.A., Macías A., and Mielke E.W., *Geometrical phases and translations*, in: "Recent Developments in Gravitation and Mathematical Physics", *Proceedings of the First Mexican School on Gravitation and Mathematical Physics*, Guanajuato, Mexico 12 – 16 Dec. 1994, Eds. A. Macías, T. Matos, O. Obregón and H. Quevedo (World Scientific: Singapore, 1995) 242-247.
2055. Morand K. and Solodukhin S.N., *Dual massive gravity*, *Phys. Lett.* **B715** (2012) 260-266.
2056. Morganstern R.E., *Riemann curvature tensor and closed geodesic paths*, *Am. J. Phys.* **45** (1977) 1173-1175.
2057. Morita K., *Quaternionic variational formalism for Poincaré gauge theory and supergravity*, *Progr. Theor. Phys.* **73** (1985) 999-1015.
2058. Morita K., *Quaternionic variational formalism for general relativity in Riemann and Riemann-Cartan space-times*, *Progr. Theor. Phys.* **128** (2012) 1107-1139.
2059. Moritsch O. and Schweda M., *On the algebraic structure of gravity with torsion including Weyl symmetry*, *Helv. Phys. Acta* **67** (1994) 289-344.
2060. Moritsch O., Schweda M., and Sorella S.P., *Algebraic structure of gravity with torsion*, *Class. Quantum Grav.* **11** (1994) 1225-1241.
2061. Moritsch O., Schweda M., and Sommer T., *Yang-Mills gauge anomalies in the presence of gravity with torsion*, *Class. Quantum Grav.* **12** (1995) 2059-2069.
2062. Mosna R.A. and Saa A., *Volume elements and torsion*, *J. Math. Phys.* **46** (2005) 112502.
2063. Mozhen W., Changgi S., and Changbai H., *Analytic solution of gauge field equation for Lorentz gravity*, *Int. J. Theor. Phys.* **28** (1989) 1437-1441.
2064. Muench U., Gronwald F., Hehl F.W., *A brief guide to variations in teleparallel gauge theories of gravity and the Kaniel-Itin model*, *Gen. Relat. Grav.* **30** (1998) 933-961.
2065. Mukhopadhyaya B. and Sengugta S., *A geometrical interpretation of parity violation in gravity with torsion*, *Phys. Lett.* **B458** (1999) 8-12.
2066. Mukhopadhyaya B., Sengugta S., and Sur S., *Space-time torsion and parity violation: A gauge-invariant formulation*, *Mod. Phys. Lett.* **A17** (2002) 43.
2067. Mukku C., *Transmogrifying torsion*, *Annales de la Fondation L. de Broglie* **32** (2007) 311-322.
2068. Mukku C. and Sayed W.A., *Torsion without torsion*, *Phys. Lett.* **B82** (1979) 382-386.
2069. Mukunda N., *An elementary introduction to the gauge approach to gravity*, In: "Gravitation, gauge theories and the early universe", Eds. B.R. Iyer, C.V. Vishveshwara, and N. Mukunda. (Dordrecht: Kluwer, 1989) 467-479.
2070. Müller-Hoissen F., *Friedman cosmology with torsion*, *Phys. Lett.* **A92** (1982) 433-434.
2071. Müller-Hoissen F., *A gauge theoretical approach to space-time structures*, *Ann. Inst. H. Poincaré* **A40** (1984) 21-34.
2072. Müller-Hoissen F. and Nitsch J., *Teleparallelism - a viable theory of gravity?*, *Phys. Rev.* **D28** (1983) 718-728.
2073. Müller-Hoissen F. and Nitsch J., *On the tetrad theory of gravity*, *Gen. Relat. Grav.* **17** (1985) 747-760.
2074. Mullick L. and Bandyopadhyay P., *Gravity without metric, torsion, and topological gravity*, *J. Math. Phys.* **36** (1995) 370-382.
2075. Mullick L. and Bandyopadhyay P., *Nonperturbative quantum geometry and torsion*, *Int. J. Mod. Phys.* **A11** (1996) 2575-2589.

2076. Mullick L. and Bandyopadhyay P., *Gravity without the metric, torsion and the cosmological-constant problem*, *Nuovo Cim.* **B112** (1997) 1289-1297.
2077. Mullick L. and Bandyopadhyay P., *Quantum gravity, torsion and space-time singularity*, *Nuovo Cim.* **B113** (1998) 1083-1089.
2078. Muschik W. and von Borzeszkowski H.-H., *Entropy identity and equilibrium conditions in relativistic thermodynamics*, *Gen. Rel. Grav.* **41** (2009) 1285-1304.
2079. Muslih S.I., *Hamilton-Jacobi quantization of two-dimensional gravity with torsion*, *Mod. Phys. Lett.* **A19** (2004) 151.
2080. Muzinich I.J., *Some remarks on torsion in Kaluza-Klein unification*, *J. Math. Phys.* **26** (1985) 1942-1947.

### $\mathcal{N}$

2081. Naik P.C., *Analog of Aharonov-Bohm effect in spin gauge theory*, *Pramana* **27** (1986) 629-635.
2082. Naik P.C., *Diffraction Aharonov-Bohm effect in spin gauge theory*, *Indian J. Phys.* **A62** (1988) 533-538.
2083. Naik P.C. and Pradhan T., *Long range interaction between spins*, *J. Phys.* **A14** (1981) 2795-2805.
2084. Nair V.P., Randjbar-Daemi S., and Rubakov V., *Massive spin-2 fields of geometric origin in curved spacetimes*, *Phys. Rev.* **D80** (2009) 104031 (8 pages).
2085. Nakamichi A., Sugamoto A., and Oda I., *Topological four-dimensional self-dual gravity*, *Phys. Rev.* **D44** (1991) 3835-3842.
2086. Nakamura A., Kikukawa H., and Kikukawa J., *Modified Maurer-Cartan condition and fiber bundle structure of the gauge theory of gravity*, *Progr. Theor. Phys.* **91** (1994) 611-624.
2087. Nakariki S., *A spinor approach to Poincaré gauge theory*, *Progr. Theor. Phys.* **81** (1989) 523-540.
2088. Nakariki S., *An exact solution in Poincaré gauge theory: monopole solution*, *J. Math. Phys.* **32** (1991) 1612-1618.
2089. Nakariki S., *Abelian solutions in Poincaré gauge gravity*, *Nuovo Cim.* **B106** (1991) 315-321.
2090. Nakariki S., *A monopole solution in Poincaré gauge gravity*, *Nuovo Cim.* **B106** (1991) 945-955.
2091. Nakariki S., Masaki T., Fukuma K., Fukui T., Mizouchi M., Ohtani T., and Tashiro T., *Interaction of massless Dirac field with a Poincaré gauge field*, *Progr. Theor. Phys.* **100** (1998) 179-190.
2092. Narlikar J.V. and Kembhavi A.K., *Non-standard cosmologies*, *Fundamentals of Cosmic Physics* **6**, no. 1-2 (1980) 1-186.
2093. Nastina L.V., *About one problem of new nonlinear electrodynamics*, *Vestnik Mosk. Univ., Ser. Astr. Fiz.* **n 6** (1978) 94-96 (in Russian).
2094. Nashed G.G.L., *General spherically symmetric nonsingular black hole solutions in a teleparallel theory of gravitation*, *Phys. Rev.* **D66** (2002) 064015 (8 pages).
2095. Nashed G.G.L., *Charged axially symmetric solution and energy in teleparallel theory equivalent to general relativity*, *Eur. Phys. J.* **C49** (2007) 851-857.
2096. Nashed G.G.L., *General regular charged space-times in teleparallel equivalent of general relativity*, *Eur. Phys. J.* **C51** (2007) 377-383.

2097. Nashed G.G.L., *Charged dilaton, energy, momentum and angular-momentum in teleparallel theory equivalent to general relativity*, *Eur. Phys. J.* **C54** (2008) 291-302.
2098. Nashed G.G.L., *Self-dual Lorentzian wormholes and energy in teleparallel theory of gravity*, *Grav. Cosmol.* **15** (2009) 256-262.
2099. Nashed G.G.L., *Cylindrically symmetric solution in teleparallel theory*, *Chinese Phys. Lett* **27** (2010) 100401.
2100. Nashed G.G.L., *Brane world black holes in teleparallel theory equivalent to general relativity and their Killing vectors, energy, momentum and angular momentum*, *Chinese Phys.* **B19** (2010) 020401.
2101. Nashed G.G.L., *Teleparallel equivalent theory of (1+1)-dimensional gravity*, *Chinese Phys.* **B19** (2010) 110401.
2102. Nashed G.G.L., *Gravitational radiation fields in teleparallel equivalent of general relativity and their energies*, *Chinese Phys.* **B19** (2010) 110402.
2103. Nashed G.G.L., *Stationary axisymmetric solutions and their energy contents in teleparallel equivalent of Einstein theory*, *Astropys. and Space Sci.* **330** (2010) 173-181.
2104. Nashed G.G.L., *Axially symmetric solution in teleparallel equivalent of general relativity, energy, momentum and angular momentum*, *Astropys. and Space Sci.* **330** (2010) 367-372.
2105. Nashed G.G.L., *Nonlinear electrodynamics coupled to teleparallel theory of gravity*, *Chinese Phys.* **B20** (2011) 020402.
2106. Nashed G.G.L., *Energy and momentum of rotating frames in tetrad gravity*, *Chinese Phys.* **B20** (2011) 100402.
2107. Nashed G.G.L., *Energy and spatial momentum of charged rotating frames in tetrad gravity*, *Chinese Phys.* **B20** (2011) 110401.
2108. Nashed G.G.L., *Five-dimensional teleparallel theory equivalent to general relativity, the axially symmetric solution, energy and spatial momentum*, *Chinese Phys.* **B20** (2011) 110402.
2109. Nashed G.G.L., *Energy and momentum of a spherically symmetric dilaton frame as regularized by teleparallel gravity*, *Ann. d. Phys. (Berlin)* **523** (2011) 450-458.
2110. Nashed G.G.L., *Killing vectors of Schwarzschild space-times in teleparallel equivalent of general relativity*, *Astropys. and Space Sci.* **333** (2011) 317-325.
2111. Nashed G.G.L., *Gravitational energy, momentum and angular momentum of Schwarzschild Anti-de Sitter space-times in the teleparallel equivalent of general relativity*, *Astropys. and Space Sci.* **335** (2011) 593-598.
2112. Nashed G.G.L., *Spherically symmetric solutions on a non-trivial frame in  $f(T)$  theories of gravity*, *Chinese Phys. Lett* **29** (2012) 050402.
2113. Nashed G.G.L., *Energy and momentum of general spherically symmetric frames on the regularizing teleparallelism*, *Chinese Phys.* **B21** (2012) 010401.
2114. Nashed G.G.L., *Energy, momentum and angular momentum in the dyadosphere of a charged spacetime in teleparallel equivalent of general relativity*, *Chinese Phys.* **B21** (2012) 030401.
2115. Nashed G.G.L., *Thermal properties of Lense-Thirring spacetime in tetrad theory of gravity*, *Chinese Phys.* **B21** (2012) 080401.
2116. Nashed G.G.L., *Cosmological application on five-dimensional teleparallel theory equivalent to general relativity*, *Chinese Phys.* **B21** (2012) 100401.
2117. Nashed G.G.L., *On the regularization of Kerr-NUT spacetime. I*, *Progr. Theor. Phys.* **127** (2012) 561-576.

2118. Nashed G.G.L., *Regularization of Kerr-NUT spacetimes and their thermodynamical quantities*, *Progr. Theor. Phys.* **127** (2012) 1077-1098.
2119. Nashed G.G.L., *Total conserved charges of Lense-Thirring spacetime in gravity theory with diffeomorphism and local lorentz symmetry*, *Progr. Theor. Phys.* **128** (2012) 767-783.
2120. Nashed G.G.L., *Spherically symmetric solution in higher-dimensional teleparallel equivalent of general relativity*, *Chinese Phys.* **B22** (2013) 020401.
2121. Nashed G.G.L., *Local Lorentz transformation and exact spherically symmetric vacuum solutions in  $f(T)$  gravity theories*, *Eur. Phys. J.* **C73** (2013) 2394.
2122. Nashed G.G.L., *Conserved charges of Kerr-Ads spacetime using Poincaré gauge version*, *Chinese Phys.* **B22** (2013) 120401.
2123. Nashed G.G.L., *Total conserved charges of Kerr-Newman spacetimes in gravity theory using a Poincaré gauge version of the teleparallel equivalent of general relativity*, *Adv. High En. Phys.* **2013** (2013) 692505 (16 pages).
2124. Nashed G.G.L., *A special exact spherically symmetric solution in  $f(T)$  gravity theories*, *Gen. Rel. Grav.* **45** (2013) 1887-1899.
2125. Nashed G.G.L., *Spherically symmetric charged-dS solution in  $f(T)$  gravity theories*, *Phys. Rev.* **D88** (2013) 104034 (6 pages).
2126. Nashed G.G.L., *Total conserved charges of charged rotating Ads solution using a Poincaré gauge version*, *Int. J. Theor. Phys.* **52** (2013) 4298-4307.
2127. Nashed G.G.L., *Thermal properties of charged dilaton black hole, energy and momentum in teleparallel equivalent of general relativity*, *Astropys. and Space Sci.* **344** (2013) 87-97.
2128. Nashed G.G.L., *Local Lorentz transformation and exact solution in  $f(T)$  gravity theories*, *Astropys. and Space Sci.* **348** (2013) 591-598.
2129. Nashed G.G.L., *Schwarzschild-NUT solution in modified teleparallel gravity theory*, *Astropys. and Space Sci.* **350** (2014) 791-797.
2130. Nashed G.G.L., *Relocalization of total conserved charge for Schwarzschild-AdS spacetime*, *Int. J. Theor. Phys.* **53** (2014) 1580-1589.
2131. Nashed G.G.L., *Energy and momentum of Robinson-Trautman space-times*, *Int. J. Theor. Phys.* **53** (2014) 1654-1665.
2132. Nashed G.G.L., *Schwarzschild solution in extended teleparallel gravity*, *Eur. Phys. Lett.* **105** (2014) 10001.
2133. Nashed G.G.L., *Total conserved charges of Kerr spacetime with one rotation parameter in 5-dimensions using Poincaré gauge theory*, *Int. J. Theor. Phys.* **54** (2015) 3490-3499.
2134. Nashed G.G.L., *FRW in quadratic form of  $f(T)$  gravitational theories*, *Gen. Rel. Gravit.* **47** (2015) 75.
2135. Nashed G.G.L., *(1 + 4)-dimensional spherically symmetric black holes in  $f(T)$* , *Gravit. Cosmol.* **23** (2017) 63-69.
2136. Nazarovski E.A. and Obukhov Yu.N., *On renormalizability of gauge theory of gravity*, *Sov. J. Phys.* **26** (1983) 1042-1047 [*Izvestia VUZov, Fizika*, n. 12 (1984) 43-49 (in Russian)].
2137. Nduka A., *Charged static fluid spheres in Einstein-Cartan theory*, *Gen. Relat. Grav.* **8** (1977) 371-377.
2138. Ne'eman Y., *Gravitational interaction of hadrons: band-spinor representations of  $GL(n, R)$* , *Proc. Nat. Acad. Sci. USA* **74** (1977) 4157-4159.
2139. Ne'eman Y., *Spinor-type fields with linear, affine and general coordinate transformations*, *Ann. Inst. H. Poincaré* **A28** (1978) 369-378.

2140. Ne'eman Y., *Gravity as the gauge theory of the parallel-transport modification of the Poincaré group*, in: "Differential Geometry Methods in Mathematical Physics", Eds. K. Bleuler, H.K. Petry and A. Reetz (Springer: Berlin, 1979) *Lect. Notes Math* **676** (1979) 189-216.
2141. Ne'eman Y., *Gravity, groups and gauges*, in: "General Relativity and Gravitation: One Hundred Years after the Birth of Albert Einstein", Ed. A. Held (Plenum: New York, 1980) vol. **1**, 309.
2142. Ne'eman Y., *Gauged and affine quantum gravity*, in: "To Fulfill a vision", Jerusalem Einstein Centennial Symposium on Gauge theories and Unification of Physical Forces, Ed. Y. Ne'eman (Addison-Wesley: Reading, 1981) 99-114.
2143. Ne'eman Y., *Superalgebras, supergroups, and geometric gauging*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 177-226.
2144. Ne'eman Y., *World spinors and the hadrons*, in: "Proc. of the 4th Marcel Grossmann Meeting on General relativity", Ed. R. Ruffini (North Holland: Amsterdam, 1986) 259-274.
2145. Ne'eman Y., *The theory of world spinors*, in: "Spinors in physics and geometry", Eds. A. Trautman and G. Furlan (World Scientific: Singapore, 1987) 313-346.
2146. Ne'eman Y., *Particle-field algebraic interplay and  $GL(4, R)$* , *Nucl. Phys. B., Proc. Suppl.* **B6** (1989) 96-101.
2147. Ne'eman Y., *A superconnection for Riemannian gravity as a broken  $SL(4, R)$  gauge theory*, *Phys. Lett.* **B427** (1998) 19-25.
2148. Ne'eman Y., *Geometric and gravity-inspired methods in QCD*, *Hadronic J.* **21** (1998) 255-278.
2149. Ne'eman Y., *Gauge theories of gravity*, *Acta Phys. Pol.* **29** (1998) 827-843.
2150. Ne'eman Y., *World spinors in metric and nonmetric backgrounds*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?", *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. de Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 134-172.
2151. Ne'eman Y., *Quantizing gravity and spacetime: where do we stand?*, *Ann. d. Phys. (Leipzig)* **8** (1999) 3-17.
2152. Ne'eman Y., *Broken symmetry: Applying the method of the superconnection for Riemannian gravity*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997), Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 40-46.
2153. Ne'eman Y. and Hehl F.W., *Test matter in a spacetime with nonmetricity*, *Class. Quantum Grav.* **14** (1997) A251-A259.
2154. Ne'eman Y., Hehl F.W., and Mielke E.W., *The generalized Erlangen program and setting a geometry for four-dimensional conformal fields*, "Math. Physics Towards the 21st Century", Eds. R.N. Sen and A. Gersten (Ben-Gurion Univ. of the Negev Press: Beer-Sheva, 1994) 59-73.
2155. Ne'eman Y. and Regge T., *Gravity and supergravity as gauge theories on a group manifold*, *Phys. Lett.* **B74** (1978) 54-56.
2156. Ne'eman Y. and Regge T., *Gauge theory of gravity and supergravity on a group manifold*, *Riv. Nuovo Cim.* **1**, no **5**, ser. 3 (1978) 1-42.
2157. Ne'eman Y. and Šijački Dj., *Unified affine gauge theory of gravity and strong interactions with finite and infinite  $GL(4, R)$  spinor fields*, *Ann. Phys. (USA)* **120** (1979) 292-315.
2158. Ne'eman Y. and Šijački Dj., *Minimal and centered graded spin-extensions of the  $SL(3, R)$  algebra*, *J. Math. Phys.* **21** (1980) 1312-1317.
2159. Ne'eman Y. and Šijački Dj.,  *$SL(4, R)$  world spinors and gravity*, *Phys. Lett.* **B157** (1985) 275-279.

2160. Ne'eman Y. and Šijački Dj., *Hadrons in an  $\overline{SL}(4, R)$  classification*, *Phys. Rev.* **D37** (1988) 3267-3283.
2161. Ne'eman Y. and Šijački Dj., *Gravity from symmetry breakdown of a gauge affine theory*, *Phys. Lett.* **B200** (1988) 489-494.
2162. Ne'eman Y. and Šijački Dj., *World spinors – construction and some applications*, *Found. Phys.* **27** (1997) 1105-1122.
2163. Nelson P.C., *Gravity with propagating pseudoscalar torsion*, *Phys. Lett.* **A79** (1980) 285-287.
2164. Nesseris S., Basilakos S., Saridakis E.N., and Perivolaropoulos L., *Viable  $f(T)$  models are practically indistinguishable from  $\Lambda$ CDM*, *Phys. Rev.* **D88** (2013) 103010 (16 pages).
2165. Nester J., *Effective equivalence of the Einstein-Cartan and Einstein theories of gravity*, *Phys. Rev.* **D16** (1977) 2395-2401.
2166. Nester J.M., *Gravity, torsion and gauge theory*, in: *An introduction to Kaluza-Klein theories*, Ed. H.C.Lee (World Scientific: Singapore, 1984) 83-115.
2167. Nester J.M., *Is there really a problem with the teleparallel theory?*, *Class. Quantum Grav.* **5** (1988) 1003-1010.
2168. Nester J.M., *A covariant Hamiltonian for gravity theories*, *Mod. Phys. Lett.* **A06** (1991) 2655.
2169. Nester J.M., *Special orthonormal frames*, *J. Math. Phys.* **33** (1992) 910-913.
2170. Nester J.M., *Some progress in classical canonical gravity*, in: *Directions in General Relativity: Proc. of Misner-Brill 1993 Int. Symp.*, Maryland, Eds. B.L. Hu, M.P. Ryan, Jr., and C.V. Vishveshwara (Cambridge: University Press, 1993) vol. 1, 245-260.
2171. Nester J.M., *General pseudotensors and quasilocal quantities*, *Class. Quantum Grav.* **21** (2004) S261.
2172. Nester J.M. and Chen C.-M., *Gravity: A gauge theory perspective*, *Int. J. Mod. Phys.* **D25** (2016) 1645002 [11 pages].
2173. Nester J.M. and Isenberg J., *Torsion singularities*, *Phys. Rev.* **D15** (1977) 2078-2087.
2174. Nester J.M., Tung R.-S., and Zhytnikov V.V., *Some spinor-curvature identities*, *Class. Quantum Grav.* **11** (1994) 983-987.
2175. Nester J.M. and Wang C., *Can torsion be treated as just another tensor field?*, *Int. J. Mod. Phys. Conf. Ser.* **07** (2012) 158.
2176. Nester J.M. and Yo H.J., *Symmetric teleparallel general relativity*, *Chinese J. Phys.* **37** (1999) 113-117.
2177. Neville D.E., *Gravity Lagrangian with ghost-free curvature-squared terms*, *Phys. Rev.* **D18** (1978) 3535-3543.
2178. Neville D.E., *“No hair” theorem for black holes in the Mansouri-Chang theory of gravitation*, *Phys. Rev.* **D19** (1979) 1033-1035.
2179. Neville D.E., *Gravity theories with propagating torsion*, *Phys. Rev.* **D21** (1980) 867-873.
2180. Neville D.E., *Experimental bounds of the coupling strength of torsion potentials*, *Phys. Rev.* **D21** (1982) 2075-2080.
2181. Neville D.E., *Spin-2 propagating torsion*, *Phys. Rev.* **D23** (1981) 1244-1249.
2182. Neville D.E., *Experimental bounds of the coupling of massless spin-1 torsion*, *Phys. Rev.* **D25** (1982) 573-576.
2183. Neville D.E., *Conformal divergences and space-time foam in an  $R + R^2$  theory*, *Phys. Rev.* **D25** (1982) 2638-2644.

2184. Neville D.E., *Torsion and chiral fermions in Kaluza-Klein theories*, *Phys. Rev.* **D33** (1986) 363-369.
2185. Nguyen Hong Chuong, *Some analytical solutions for isotropic viscous models in the Poincaré gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 5, 70-75 (in Russian).
2186. Nguyen Hong Chuong, *Cosmological constant and regular Bianchi type-I models with radiation in the minimum quadratic gauge theory of gravity*, *Phys. Lett.* **A142** (1989) 326-330.
2187. Nguyen Hong Chuong, *Cosmological constant and the dynamics of the Bianchi type I models with spin and torsion: A qualitative investigation*, *Gen. Relat. Grav.* **22** (1990) 1229-1238.
2188. Nguyen Hong Chuong and Nguen Van Hoang, *On the new solutions for cosmological equations of the Poincaré gauge theory of gravity*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 3 (1986) 94-97 (in Russian).
2189. Nguyen Hong Chuong and Nguen Van Hoang, *Cosmological constant and gravitational repulsion effect. 1. Homogeneous models with radiation*, *Int. J. Mod. Phys.* **D2** (1993) 443-450.
2190. Ni W.-T., *Hoiman-Rosenbaum-Ryan-Shepley torsion theory and Eotvos-Dicke-Braginsky experiments*, *Phys. Rev.* **D19** (1979) 2260-2263.
2191. Ni W.-T., *Conditions for an affine manifold with torsion to have a Riemann-Cartan structure*, *Math. Proc. Camb. Phil. Soc.* **90** (1981) 517-527.
2192. Ni W.-T., *Searches for the role of spin and polarization in gravity*, *Rep. Prog. Phys.* **73** (2010) 056901.
2193. Nolić I., *Dirac Hamiltonian structure of  $R + R^2 + T^2$  Poincaré gauge theory of gravity without gauge fixing*, *Phys. Rev.* **D30** (1984) 2508-2520.
2194. Nolić I.A., *Canonical structure of Poincaré gauge-invariant theory of gravity*, *Fizika (SFRJ)* **18** (1986) 135-139.
2195. Nolić I.A., *Schwinger's energy-energy brackets in the  $U_4$ -theory of gravity*, *Class. Quantum Grav.* **7** (1990) 1895-1903.
2196. Nolić I.A., *Constraint algebra from local Poincaré symmetry*, *Gen. Relat. Grav.* **24** (1992) 159-170.
2197. Niederle J., *Gauge formulation of gravitation theories*, *Czech. J. Phys.* **B32** (1982) 672-673.
2198. Niederle J., *On gauge formulations of gravitation theories*, in: "Gauge Theories of Fundamental Interactions. Proc. XXXII Semester in Banach Int. Math. Center, Warsaw, 1988", Eds. M.Pawlowski and R.Raczka (World Scientific: Singapore, 1990) 329-340.
2199. Nieh H.T., *Gauss-Bonnet and Bianchi identities in Riemann-Cartan type gravitational theories*, *J. Math. Phys.* **21** (1980) 1432-1441.
2200. Nieh H.T., *A spontaneously broken conformal gauge theory of gravitation*, *Phys. Lett.* **A88** (1982) 388-390.
2201. Nieh H.T., *Local Poincaré symmetry: quantum effects*, in: "Proc. of the 4th Marcel Grossmann Meeting on General relativity", Ed. R. Ruffini (North Holland: Amsterdam, 1986) 675-683.
2202. Nieh H.T. and Rauch R., *Riemann-Cartan type gravitational theories satisfying Birkhoff's theorem*, *Phys. Lett.* **A81** (1981) 113-115.
2203. Nieh H.T. and Yan M.L., *Quantized Dirac field in curved Riemann-Cartan background. I. Symmetry properties, Green's functions*, *Ann. Phys. (USA)* **138** (1982) 237-259.
2204. Nieh H.T. and Yan M.L., *An identity in Riemann-Cartan geometry*, *J. Math. Phys.* **23** (1982) 373-374.
2205. Nikiforova V., Randjbar-Daemi S., and Rubakov V., *Infrared modified gravity with dynamical torsion*, *Phys. Rev.* **D80** (2009) 124050 (10 pages).

2206. Nikiforova V., Randjbar-Daemi S., and Rubakov V., *Self-accelerating universe in modified gravity with dynamical torsion*, *Phys. Rev.* **D95** (2017) 024013 (6 pages).
2207. Nikolaenko V.M., *Gauge-invariant gravitational field equations in a theory with quadratic Lagrangians*, *Sov. Phys. Doklady* **19** (1975) 684-685 [*Doklady AN SSSR* **218**, n. 11 (1974) 1068-1070 (in Russian)].
2208. Nikolaenko V.M., *Second order Lagrangians of curvature in the theory of gravity*, *Acta Phys. Pol.* **B7** (1976) 681-692.
2209. Nikolaenko V.M., *Back reaction of a quantized field in the gauge treatment of gravity*, *Acta Phys. Pol.* **B8** (1977) 911-918.
2210. Ning B. and Li F.-L., *Relative entropy and torsion coupling*, *Phys. Rev. D* **94** (2016) 126007 (16 pages).
2211. Nikolic I.A., *Dirac Hamiltonian formulation and algebra of the constraints in the Einstein-Cartan theory*, *Class. Quantum Grav.* **12** (1995) 3103-3114.
2212. Nishida K., *A generalized gauge theory of gravity*, *Prog. Theor. Phys.* **123** (2010) 227-235.
2213. Nishino H., *On supergravity with propagating Lorentz connection*, *Progr. Theor. Phys.* **66** (1981) 287-302.
2214. Nishino H., *Local tensor calculus in supergravity with propagating Lorentz connection*, *Progr. Theor. Phys.* **68** (1982) 975-988.
2215. Nishino H., *Consistency of couplings in supergravity theory with propagating Lorentz connection*, *Progr. Theor. Phys.* **68** (1982) 1765-1775.
2216. Nishino H., and Rajpoot S., *Teleparallelism for a massive spin-2 field*, *Class. Quantum Grav.* **28** (2011) 125019 (11 pages).
2217. Nishioka M., *A remark on coefficients of connection with two gauge fields*, *Gen. Relat. Grav.* **15** (1983) 591-593.
2218. Nishioka M., *Local transformation and symmetric connection in space-time*, *Lett. Nuovo Cim.* **36** (1983) 266-268.
2219. Nitsch J., *The macroscopic limit of the Poincaré gauge field theory of gravitation*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 63-91.
2220. Nitsch J. and Hehl F.W., *Translational gauge theory of gravity: post-Newtonian approximation and spin precession*, *Phys. Lett.* **B90** (1980) 98-102.
2221. Nomura K., Shirafuji T., and Hayashi K., *Spinning test particles in spacetime with torsion*, *Progr. Theor. Phys.* **86** (1991) 1239-1258.
2222. Nomura K., Shirafuji T., and Hayashi K., *Semiclassical particles with arbitrary spin in the Riemann-Cartan space-time*, *Progr. Theor. Phys.* **87** (1992) 1275-1291.
2223. Norris L.K., *Einstein-Maxwell dynamics as a  $P(4)$  affine theory*, *Phys. Rev.* **D31** (1985) 3090-3098.
2224. Norris L.K. and Davis W.R., *Gauge formulations of gravitation and their fundamental relationship to Riemannian holonomy structure*, *Lett. Nuovo Cim.* **27** (1980) 122-128.
2225. Norris L.K., Fulp R.O., and Davis W.R., *Underlying fiber bundle structure of  $A(4)$  gauge theories*, *Phys. Lett.* **A79** (1980) 278-282.
2226. Novello M., *Scalar and massless vector fields in Cartan space*, *Phys. Lett.* **A59** (1976) 105-106.
2227. Nurgaliev I.S., *Spherically symmetric static solutions in Einstein-Cartan theory*, *Sov. J. Phys.* **25** (1982) 820-824 [*Izvestia VUZov, Fizika*, n. 9 (1982) 60-64 (in Russian)].

2228. Nurgaliev I.S., *Cosmological nucleosynthesis and torsion of space-time*, *Izvestia VUZov, Fizika*, n. 5 (1984) 115-117 (in Russian).
2229. Nurgaliev I.S., *Spinor-like spinning nonpolarised liquid in Einstein-Cartan theory*, in: "On Relativity Theory. Proc. of Sir A.Eddington Centenary Symp.", Eds. Y. Choquet-Bruhat and T.M. Karade (World Scientific: Singapore, 1985) vol. 2, 174-177.
2230. Nurgaliev I.S. and Piskareva O.B., *Structural stability of cosmological models in Einstein-Cartan theory*, *Vestnik Mosk. Univ., ser. Fiz. Astr.* **29**, n. 2 (1988) 16-22 (in Russian).
2231. Nurgaliev I.S. and Ponomariev V.N., *Cosmological constant and torsion of space-time*, *Sov. J. Phys.* **25** (1982) 796-798 [*Izvestia VUZov, Fizika*, n. 9 (1982) 32-35 (in Russian)].
2232. Nurgaliev I.S. and Ponomariev V.N., *Torsion of space-time and cosmology*, *Sov. J. Phys.* **25** (1982) 929-931 [*Izvestia VUZov, Fizika*, n. 10 (1982) 63-65 (in Russian)].
2233. Nurgaliev I.S. and Ponomariev V.N., *The earliest evolutionary stages of the Universe and space-time torsion*, *Phys. Lett.* **B130** (1983) 378-379.
2234. Nurgaliev I.S. and Ponomariev V.N., *Pre-Friedmann stage of the evolution and space-time torsion*, in: "Probl. Theor. Grav. and Element. Particles", Ed. K.P. Stanyukovich (Energoatomizdat: Moscow, 1985) **16**, 68-76 (in Russian).

## O

2235. Oancea S., *The effects of torsion for fields of axial symmetry*, *Acta Phys. Hungar.* **71** (1992) 193-200.
2236. Oancea S., *The spin of the Kerr field*, *Bull. Inst. Politeh. Iasi I* **41** N 1-2 (1995) 107-111.
2237. Obata T., *Torsion coupling constant of new general relativity*, *Progr. Theor. Phys.* **70** (1983) 622-634.
2238. Obata T. and Oshima H., *Schouten manifold with a spinor structure*, *Progr. Theor. Phys.* **69** (1983) 314-322.
2239. Obata T. and Oshima H., *Dirac equation on the Schouten spacetime*, *Progr. Theor. Phys.* **71** (1984) 183-200.
2240. O'Brien G.M. and Tchraikian D.H., *Spin connection generalized Yang-Mills fields on double dual generalized Einstein-Cartan backgrounds*, *J. Math. Phys.* **29** (1988) 1212-1219.
2241. Obukhov Yu.N., *Conformal invariance and space-time torsion*, *Phys. Lett.* **A90** (1982) 13-16.
2242. Obukhov Yu.N., *Spectral geometry of the Riemann-Cartan space-time and the axial anomaly*, *Phys. Lett.* **B108** (1982) 308-310.
2243. Obukhov Yu.N., *Spectral geometry of the Riemann-Cartan space-time*, *Nucl. Phys.* **B212** (1983) 237-254.
2244. Obukhov Yu.N., *Arbitrary spin field equations and anomalies in the Riemann-Cartan space-time*, *J. Phys.* **A16** (1983) 3795-3804.
2245. Obukhov Yu.N., *The Palatini principle for manifold with boundary*, *Class. Quantum Grav.* **4** (1987) 1085-1091.
2246. Obukhov Yu.N., *Spin-driven inflation*, *Phys. Lett.* **A182** (1993) 214-216.
2247. Obukhov Yu.N., *Two-dimensional Poincaré gauge gravity with matter*, *Phys. Rev.* **D50** (1994) 5072-5086.
2248. Obukhov Yu.N., *On a model of an unconstrained hyperfluid*, *Phys. Lett.* **A210** (1996) 163-167.

2249. Obukhov Yu.N., *Connection forms in gravity models*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 269-297.
2250. Obukhov Yu.N., *On the gravitational moments of a Dirac particle*, *Acta Phys. Polon.* **B29** (1998) 1131-1138.
2251. Obukhov Yu.N., *Analogue of black string in Yang-Mills gauge theory*, *Int. J. Theor. Phys.* **37** (1998) 1455-1467.
2252. Obukhov Yu.N., *On physical foundations and observational effects of cosmic rotation*, accepted in "Colloquium on Cosmic Rotation" (Berlin, Feb. 1998), Eds. M. Scherfner, T. Chrobok, and M. Shefaat (Wissenschaft und Technik Verlag: Berlin, 2000) 23-96.
2253. Obukhov Yu.N., *New solutions in 3D gravity*, *Phys. Rev.* **D68** (2004) 124015 (8 pages).
2254. Obukhov Yu.N., *Two-dimensional metric-affine gravity*, *Phys. Rev.* **D69** (2004) 064009 (6 pages).
2255. Obukhov Yu.N., *Plane waves in metric-affine gravity*, *Phys. Rev.* **D73** (2006) 024025 (6 pages).
2256. Obukhov Yu.N., *Equations of motion in the gauge gravity models*, *Acta Phys. Pol. B Proc. Suppl.* **1** (2008) 161-166.
2257. Obukhov Yu.N., *Gravitational waves in Poincaré gauge gravity theory*, *Phys. Rev.* **D95** (2017) 084028 (12 pages).
2258. Obukhov Yu.N. and Hehl F.W., *On the relation between quadratic and linear curvature Lagrangians in Poincaré gauge gravity*, *Acta Phys. Polon.* **B27** (1996) 2685-2693.
2259. Obukhov Yu.N. and Hehl F.W., *Black holes in two dimensions*, in: "Black holes: Theory and observations", *Proc. of 179th Heraeus Seminar, Bad Honnef, Germany, 18-22 August 1997*, Eds. F.W. Hehl, C. Kiefer and R.J.K. Metzler (Springer: Berlin) *Lect. Notes Phys.* **514** (1998) 289-316.
2260. Obukhov Yu.N. and Hehl F.W., *Extended Einstein-Cartan theory á la Diakonov: The field equations*, *Phys. Lett.* **B713** (2012) 321-325.
2261. Obukhov Yu.N. and Hehl F.W., *Quantum gravity model with fundamental spinor fields*, *Phys. Part. Nucl.* **45** (2014) 223-225.
2262. Obukhov Yu.N. and Korotky V.A., *The Weyssenhoff fluid in Einstein-Cartan theory*, *Class. Quantum Grav.* **4** (1987) 1633-1657.
2263. Obukhov Yu.N., Krechet V.G., and Ponomariev V.N., *Torsion, segmentary curvature and the structure of space-time*, in: "Gravitation and General Relativity", Ed. V.R. Kaigorodov (Univ. Press: Kazan, 1978) vol. **14**, 121-127 (in Russian).
2264. Obukhov Yu.N., Mielke E.W., Budczies J., and Hehl F.W., *On the chiral anomaly in non-Riemannian spacetimes*, *Found. Phys.* **27** (1997) 1221-1236.
2265. Obukhov Yu.N. and Nazarowski E.A., *On quantum gravity with dynamical torsion*, in: "Relativistic Astrophysics and Cosmology. Proc. of the Sir A.Eddington Centenary Symp.", Eds. V. DeSabbata and T.M. Karade (World Scientific: Singapore, 1984) vol. **1**, 137-169.
2266. Obukhov Yu.N. and Pereira J.G., *Teleparallel origin of the Fierz picture for spin-2 particle*, *Phys. Rev.* **D67** (2003) 044008 (5 pages).
2267. Obukhov Yu.N. and Pereira J.G., *Metric-affine approach to teleparallel gravity*, *Phys. Rev.* **D67** (2003) 044016 (17 pages).
2268. Obukhov Yu.N. and Pereira J.G., *Lessons of spin and torsion: Reply to "Consistent coupling to Dirac fields in teleparallelism"*, *Phys. Rev.* **D69** (2004) 128502 (3 pages).
2269. Obukhov Yu.N. and Piskareva O.B., *Spinning fluid in general relativity*, *Class. Quantum Grav.* **6** (1989) 15-19.

2270. Obukhov Yu.N., Ponomarev V.N., and Zhytnikov V.V., *Quadratic Poincaré gauge theory of gravity: a comparison with the general relativity theory*, *Gen. Relat. Grav.* **21** (1989) 1107-1142.
2271. Obukhov Yu.N., Portales-Oliva F., Puetzfeld D., and Rubilar G.F., *Invariant conserved currents in generalized gravity*, *Phys. Rev.* **D92** (2015) 104010 (16 pages).
2272. Obukhov Yu.N. and Pronin P.I., *Quantum field theory in Riemann-Cartan space*, in: "Problems of Gravity", Ed. D.V. Gal'tsov (Moscow Univ.Publ. House: Moscow, 1986) 130-150 (in Russian).
2273. Obukhov Yu.N. and Pronin P.I., *Renormalization of gauge field theories in Riemann-Cartan space-time. I. Abelian models*, *Acta Phys. Polon.* **B14** (1988) 541-560.
2274. Obukhov Yu.N. and Pronin P.I., *Problems and perspectives of quantum gauge theory of gravity*, in: "Problems of Gravity. Proc. VII Soviet Grav. Conf." (Yerevan Univ.: Yerevan, 1989) 121-136 (in Russian).
2275. Obukhov Yu.N. and Pronin P.I., *Physical effects in the theory of gravity with torsion*, in: "Itogi Nauki i Techniki. Ser. Class. Field Theory and Gravity", Ed. V.N. Melnikov (VINITI: Moscow, 1991) vol. **2**, 112-170 (in Russian).
2276. Obukhov Yu.N. and Puetzfeld D., *Conservation laws in gravitational theories with general nonminimal coupling*, *Phys. Rev.* **D87** (2013) 081502(R) (6 pages).
2277. Obukhov Yu.N. and Puetzfeld D., *Multipolar test body equations of motion in generalized gravity theories*, in: "Equations of Motion in Relativistic Gravity", Eds. D. Puetzfeld, C. Lämmerzahl, and B. Schutz, *Fundamental Theories of Physics* **179** (Springer: Switzerland, 2015) 67-119.
2278. Obukhov Yu.N. and Rubilar G.F., *Covariance properties and regularization of conserved currents in tetrad gravity*, *Phys. Rev.* **D73** (2006) 124017 (14 pages).
2279. Obukhov Yu.N. and Rubilar G.F., *Invariant conserved currents in gravity theories with local Lorentz and diffeomorphism symmetry*, *Phys. Rev.* **D74** (2006) 064002 (19 pages).
2280. Obukhov Yu.N. and Rubilar G.F., *Invariant conserved currents in gravity theories: Diffeomorphisms and local gauge symmetries*, *Phys. Rev.* **D76** (2007) 124030 (13 pages).
2281. Obukhov Yu.N. and Rubilar G.F., *Invariant conserved currents for gravity*, *Phys. Lett.* **B660** (2008) 240-246.
2282. Obukhov Yu.N., Rubilar G.F., and Pereira J.G. *Conserved currents in gravitational models with quasi-invariant Lagrangians: Application to teleparallel gravity*, *Phys. Rev.* **D74** (2006) 104007 (10 pages).
2283. Obukhov Yu.N., Silenko A.J., and Teryaev O.V., *Spin-torsion coupling and gravitational moments of Dirac fermions: Theory and experimental bounds*, *Phys. Rev.* **D90** (2014) 124068 (13 pages).
2284. Obukhov Yu.N., Silenko A.J., and Teryaev O.V., *Spin-gravity interactions and equivalence principle*, *Int. J. Mod. Phys.: Conf. Ser.* **40** (2016) 1660081 (6 pages).
2285. Obukhov Yu.N. and Solodukhin S.N., *Dynamical gravity and conformal and Lorentz anomalies in two dimensions*, *Class. Quantum Grav.* **7** (1990) 2045-2054.
2286. Obukhov Yu.N., Solodukhin S.N., and Mielke E.W., *Coupling of lineal Poincaré gauge gravity to scalar fields*, *Class. Quantum Grav.* **11** (1994) 3069-3079.
2287. Obukhov Yu.N. and Tresguerres R., *Hyperfluid – a model of classical matter with hypermomentum*, *Phys. Lett.* **A184** (1993) 17-22.
2288. Obukhov Yu.N. and Vargas T., *Gödel type solution in teleparallel gravity*, *Phys. Lett.* **A327** (2004) 365-373.
2289. Obukhov Yu.N. and Vlachynsky E.J., *Einstein-Proca model: spherically symmetric solutions*, *Ann. Phys. (USA)* **8** (1999) 497-509.

2290. Obukhov Yu.N., Vlachynsky E.J., Esser W., and Hehl F.W., *Effective Einstein theory from metric-affine gravity models via irreducible decompositions*, *Phys. Rev.* **D56** (1997) 7769-7778.
2291. Obukhov Yu.N., Vlachynsky E.J., Esser W., Tresguerres R., and Hehl F.W., *An exact solution of the metric-affine gauge theory with dilation, shear, and spin charges*, *Phys. Lett.* **A220** (1996) 1-9.
2292. Obukhov Yu.N. and Yakushin I.V., *On the experimental estimates of the axial torsion mass and coupling constants*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 197-215.
2293. Obukhov Yu.N. and Yakushin I.V., *Experimental bounds on parameters of spin-spin interaction in gauge theory of gravitation*, *Theor. Math. Phys.* **90** (1992) 209-213.
2294. Obukhov Yu.N. and Yakushin I.V., *Experimental restrictions on spin-spin interactions in gauge gravity*, *Int. J. Theor. Phys.* **31** (1992) 1993-2001.
2295. O'Connell R.F., *Contact interactions in the Einstein and Einstein-Cartan-Sciama-Kibble (ECSK) theories of gravitation*, *Phys. Rev. Lett.* **37** (1976) 1653-1655; **38**, 1977 (E).
2296. O'Connell R.F., *Attractive spin-spin contact interactions in the Einstein-Cartan-Sciama-Kibble torsion theory of gravitation*, *Phys. Rev.* **D16** (1977) 1247.
2297. O'Raifeartaigh L. and Straumann N., *Gauge theory: Historical origins and some modern developments*, *Rev. Mod. Phys.* **72** (2000) 1-23.
2298. Oda I., *Topological four-dimensional gravity*, *Progr. Theor. Phys., Suppl.* **110** (1992) 41-56.
2299. Odintsov S.D., *Some remarks on the effective action in curved space-time with torsion*, *Acta Phys. Pol.* **B21** (1990) 863-870.
2300. Odintsov S.D., *Renormalization group, effective action and grand unification theories in curved space-time*, *Fortschr. Phys.* **39** (1991) 621-641.
2301. Odintsov S.D., *Nonlocal effective action in two-dimensional curved space with torsion, and cosmology*, *Russ. Phys. J.* **34** (1991) 851-854 [*Izvestia VUZov, Fizika*, n. 10 (1991) 5-8 (in Russian)].
2302. Ogino S., *Roles of torsion particles in the very early Universe*, *Progr. Theor. Phys.* **73** (1985) 84-99.
2303. Ogino S., *Stability of the Friedmann universe in the Poincaré gauge theory*, *Progr. Theor. Phys.* **82** (1989) 723-736.
2304. Ogino S., *Stability of the Friedmann universe in the Poincaré gauge theory II*, *Progr. Theor. Phys.* **89** (1993) 371-388.
2305. Oh P., *Introduction to Poincaré gravity*, *J. Korean Phys. Soc. Suppl.* **28** (1995) S543-S548.
2306. Oh C. H. and Singh K., *Some comments on torsion in the Kaluza-Klein scheme*, *Class. Quantum Grav.* **6** (1989) 1053-1055.
2307. Okolów A., *ADM-like Hamiltonian formulation of gravity in the teleparallel geometry*, *Gen. Rel. Grav.* **45** (2013) 2569-2610.
2308. Okolów A., *ADM-like Hamiltonian formulation of gravity in the teleparallel geometry: derivation of constraint algebra*, *Gen. Rel. Grav.* **46** (2014) 1636 (32 pages).
2309. Okolów A. and Świeżewski J., *Hamiltonian formulation of a simple theory of the teleparallel geometry*, *Class. Quantum Grav.* **29** (2012) 045008.
2310. Öktem F., *On affine field laws*, *Proc. Roy. Irish Soc.* **A64** (1966) 107-113.
2311. Okubo S., *A BRST-like operator for space with zero curvature but non-zero torsion tensor*, *Gen. Relat. Grav.* **23** (1991) 599-605.

2312. Okubo S., *Nienhuis-Bianchi identity and BRST-like operators*, *J. Math. Phys.* **33** (1992) 895-900.
2313. Oliveira C.G. and Amaral M., *Torsion generated field in a four-dimensional Weitzenböck space*, *Rev. bras. fis.* **13** (1983) 201-219.
2314. Oliveira J.D. de, Teixeira A.F.F., and Tiomno J., *Homogeneous cosmos of Weyssenhoff fluid in Einstein-Cartan space*, *Phys. Rev.* **D34** (1986) 3661-3665.
2315. Oliveira H.P. de, *Spinning fluids in the Einstein-Cartan theory: a variational formulation*, *Gen. Relat. Grav.* **25** (1993) 473-481.
2316. Oliveira H.P. de, and Salim J.M., *Spinning fluids in general relativity: a variational formulation*, *Class. Quant. Grav.* **8** (1991) 1199-1204.
2317. Oliveira H.P. de, and Salim J.M., *On the influence of spin on entropy production in curved spacetimes*, *Phys. Lett.* **A170** (1992) 370-372.
2318. Oliveira H.P. de, and Salim J.M., *A variational approach to charged spinning fluids in general relativity*, *Phys. Lett.* **A201** (1995) 381-386.
2319. Oliveira H.P. de, Salim J.M., and Sautu S.L., *Non-singular inflationary cosmologies in Weyl integrable spacetime*, *Class. Quantum Grav.* **14** (1997) 2833-2843.
2320. Olmo G.J. and Rubiera-Garcia D., *Importance of torsion and invariant volumes in Palatini theories of gravity*, *Phys. Rev.* **D88** (2013) 084030 (11 pages).
2321. Ong Y.C., Izumi K., Nester J.M., and Chen P., *Problems with propagation and time evolution in  $f(T)$  gravity*, *Phys. Rev.* **D88** (2013) 024019 (15 pages).
2322. Orsaesli C.A., *Torsion dynamics in multidimensional unified theories*, *Phys. Lett.* **B140** (1984) 29-43.
2323. Ortín T., *Gravity and Strings* (Cambridge University Press: Cambridge. 2004).
2324. Otalora G., *Scaling attractors in interacting teleparallel dark energy*, *JCAP* **07** (2013) 044.
2325. Otalora G., *A novel teleparallel dark energy model*, *Int. J. Mod. Phys.* **D25** (2016) 1650025 (15 pages).
2326. Otalora G. and Saridakis E.N., *Modified teleparallel gravity with higher-derivative torsion terms*, *Phys. Rev.* **D94** (2016) 084021 (13 pages).
2327. Ozakin A. and Yavari A., *A geometric theory of thermal stresses*, *J. Math. Phys.* **51** (2010) 032902 (32 pages).

## P

2328. Pagani C. and Percacci R., *Quantization and fixed points of non-integrable Weyl theory*, *Class. Quantum Grav.* **31** (2014) 115005.
2329. Pagani C. and Percacci R., *Quantum gravity with torsion and non-metricity*, *Class. Quantum Grav.* **32** (2015) 195019 (17 pages).
2330. Page C.H., *Electromagnetism in non-Riemannian space*, *J. Res. Nat. Bur. Stand.* **B81** (1977) 1-3.
2331. Pagels H., *Spin and gravitation*, *Ann. Phys. (USA)* **31** (1965) 64-87.
2332. Pagels H.R., *Gauge theory of gravity*, *Phys. Rev.* **D27** (1983) 2299-2304.
2333. Pagels H.R., *Gravitational gauge fields and the cosmological constant*, *Phys. Rev.* **D29** (1984) 1690-1698.

2334. Pak D.G., Kim Y., and Tsukioka T., *Lorentz gauge theory as a model of emergent gravity*, *Phys. Rev.* **D85** (2012) 084006 (11 pages).
2335. Palatini A., *Deduzione invariante delle equazioni gravitazionali del principio di Hamilton*, *Rend. Circ. Mat. Palermo* **43** (1919) 203-212; English translation: *Invariant deduction of the gravitational equations from the principle of Hamilton*, in: "Proc. 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 477-488.
2336. Palese M. and Winterroth E., *Higgs fields on spinor gauge-natural bundles*, *J. Phys.: Conf. Ser.* **411** (2013) 012025 (5 pages).
2337. Palle D., *On certain relationships between cosmological observables in the Einstein-Cartan gravity*, *Nuovo Cim.* **B111** (1996) 671-675.
2338. Palle D., *On primordial cosmological density fluctuations in the Einstein-Cartan gravity and COBE data*, *Nuovo Cim.* **B114** (1999) 853-860.
2339. Palle D., *On the Einstein-Cartan cosmology vs. Planck data*, *J. Exp. Theor. Phys.* **118** (2014) 587-592.
2340. Pan Y.Y. and Ren J., *Spin-rotation coupling in the teleparallelism description in high speed rotation system*, *Int. J. Theor. Phys.* **50** (2011) 724-728.
2341. Pandey S.N., Pandey H.S., and Tiwari R., *Some perfect fluid spheres with non-zero spin density*, *Indian J. Pure and Appl. Math.* **13** (1982) 88-95.
2342. Park C.J. and Yoon Y., *Conformal couplings in induced gravity*, *Gen. Rel. Grav.* **29** (1997) 765-772.
2343. Pascual-Sánchez J.-F., *Macroscopic limit of the Poincaré gauge theory of gravity*, *Lett. Nuovo Cim.* **44** (1985) 326-328.
2344. Pascual-Sánchez J.-F., *On the limits of Poincaré gauge theory*, *Phys. Lett.* **A108** (1985) 387-390.
2345. Pasic V. and Barakovic E., *PP-waves with torsion: a metric-affine model for the massless neutrino*, *Gen. Rel. Grav.* **46** (2014) 1787.
2346. Pasic V., Barakovic E., and Okcic N., *A new representation of the field equations of quadratic metric-affine gravity*, *Advances in Mathematics* **3**, n. 1 (2014) 33-46.
2347. Pasic V. and Vassiliev D., *pp-waves with torsion and metric-affine gravity*, *Class. Quantum Grav.* **22** (2005) 3961.
2348. Pasini A., *Precession of opposite chirality for the spin vector in a Riemann-Cartan framework*, *Phys. Lett.* **A151** (1990) 459-463.
2349. Pasini A., Pelino V., and Potesta S., *Torsion and attractors in the Kolmogorov hydrodynamical system*, *Phys. Lett.* **A241** (1998) 77-83.
2350. Pasmatsiou K., Tsagas C.G., and Barrow J.D., *Kinematics of Einstein-Cartan universes*, *Phys. Rev.* **D95** (2017) 104007 (15 pages).
2351. Pavelle R., *Unphysical characteristics of Yang's pure-space equations*, *Phys. Rev. Lett.* **37** (1976) 961-964.
2352. Pavelle R., *Mansouri-Chang gravitation theory*, *Phys. Rev. Lett.* **40** (1978) 267-270.
2353. Pavelkin V.N. and Panov V.F., *A nonstationary cosmological model with rotation in the Einstein-Cartan theory*, *Izvestiya VUZov, Fizika*, n. **8** (1993) 90-94 (in Russian).
2354. Penrose R., *Spin and torsion in general relativity*, *Found. Phys.* **13** (1983) 325-339.
2355. Percacci R., *General relativity as a soldered nonlinear sigma model*, *Gen. Relat. Grav.* **14** (1982) 1043-1049.

2356. Percacci R., *Gauge group of gravity, spinors, and anomalies*, *Int. J. Theor. Phys.* **25** (1986) 493-507.
2357. Percacci R., *Geometry of nonlinear field theories* (World Scientific: Singapore, 1986) 255 p.
2358. Percacci R., *The Higgs phenomenon in quantum gravity*, *Nucl. Phys.* **B353** (1991) 271-290.
2359. Pereira J.G., *Lorentz connections and gravitation*, *AIP Conf. Proc.* **1483**, n. 1 (2012) 239-259.
2360. Pereira J.G., *Teleparallelism: A new insight into gravity*, in: "Springer Handbook of Space-time" Eds. Ashtekar A., and Petkov V. (Springer: Berlin, 2014) 197-212.
2361. Pereira J.G. and Vargas T., *Regge calculus in teleparallel gravity*, *Class. Quantum Grav.* **19** (2002) 4807.
2362. Pereira J.G., Vargas T., and Zhang C.M., *Axial-vector torsion and the teleparallel Kerr spacetime*, *Class. Quantum Grav.* **18** (2001) 833-841.
2363. Pereira J.G. and Zhang C.M., *Some remarks on the neutrino oscillation phase in a gravitational field*, *Gen. Relat. Grav.* **32** (2000) 1633-1637.
2364. Peres A., *Spinor fields in generally covariant theories*, *Suppl. Nuovo Cim.* **24** (1962) 389-452.
2365. Petti R.J., *Some aspects of the geometry of first-quantized theories*, *Gen. Relat. Grav.* **7** (1976) 869-883.
2366. Petti R.J., *Some aspects of the geometry of first-quantized theories II*, *Gen. Relat. Grav.* **8** (1977) 887-903.
2367. Petti R.J., *On the local geometry of rotating matter*, *Gen. Relat. Grav.* **18** (1986) 441-460.
2368. Petti R.J., *Affine defects and gravitation*, *Gen. Relat. Grav.* **33** (2001) 163-172.
2369. Petti R.J., *Translational spacetime symmetries in gravitational theories*, *Class. Quantum Grav.* **23** (2006) 737.
2370. Pietropaolo F. and Toller M., *The motion of a Dirac wave packet in gravitational field*, *Nuovo Cim.* **B77** (1983) 129-142.
2371. Pilch K.A., *Geometrical meaning of the Poincaré group gauge theory*, *Lett. Math. Phys.* **4** (1980) 49-51.
2372. Pinheiro M.J. *The flyby anomaly and the effect of a topological torsion current*, *Phys. Lett.* **A378** (2014) 3007-3011.
2373. Pirogov Yu.F., *Gravity as the affine goldstone phenomenon and beyond*, *Phys. Atomic Nucl.* **68** (2005) 1904-1915.
2374. Pitts J.B. and Schieve W.C., *On parametrized general relativity*, *Found. Phys.* **28** (1998) 1417-1424.
2375. Platania G. and Rosania R., *An anisotropic cosmological model with spin-1/2 particles*, *Nuovo Cim.* **B111** (1996) 1195-1209.
2376. Platania G. and Rosania R., *A universe with torsion*, *Europhys. Lett.* **38** (1997) 7-11.
2377. Platania G. and Rosania R., *A universe with torsion and cosmological constant*, *Nuovo Cim.* **B112** (1997) 1337-1349.
2378. Poberii E.A., *Nonmetricity driven inflation*, *Helv. Phys. Acta* **67** (1994) 745-752.
2379. Poberii E.A., *Metric-affine scale-covariant gravity*, *Gen. Rel. Grav.* **26** (1994) 1011-1054.
2380. Poberii E.A., *Spinor-type objects in metric-affine space-time*, *J. High Ener. Phys.* **(7)** (1998) U327-U343.

2381. Polishchuk R.F., *Energy problem in the Einstein-Cartan theory*, *Gravitation and Cosmology* **21** (2015) 220-235.
2382. Polishchuk R.F., *Tetrad currents in the Einstein-Cartan theory*, *Int. J. Mod. Phys.: Conf. Ser.* **41** (2016) 1660118 (5 pages).
2383. Ponomarev V.N., *Observable effects of torsion in space-time*, *Bull. Acad. Polon. Sci., ser. mat., astr., phys.*, **19** (1971) 545-550.
2384. Ponomarev V.N., *Observable effects of space-time torsion*, *Vestnik Mosk. Univ., ser. fiz., astr.*, n. 5 (1974) 139-143 (in Russian).
2385. Ponomarev V.N., *Cosmological model of space-time with torsion*, *Vestnik Mosk. Univ., ser. fiz., astr.*, n. 12 (1974) 730-735 (in Russian).
2386. Ponomarev V.N., *Singularities and torsion*, in: "Relativity Theory and Gravitation", Ed. V.I.Rodichev (Nauka: Moscow, 1976) 167-176.
2387. Ponomarev V.N., Barvinski A.O., and Obukhov Yu.N., *Geometrodynamical methods and the gauge approach to the theory of gravitational interactions* (Ergoatomizdat: Moscow, 1985) 168 pp. (in Russian).
2388. Ponomarev V.N., Krechet V.G., and Pronin P.I., *Problems of classical and quantum collapse IX, Torsion, wave fields, cosmology*, *Sov. J. Phys.* **21** (1978) 279-284 [*Izvestiya VUZov, Fizika*, n. 3 (1978) 18-24 (in Russian)].
2389. Ponomarev V.N. and Obukhov Yu.N., *The generalized Einstein-Maxwell theory of gravitation*, *Gen. Relat. Grav.* **14** (1982) 309-330.
2390. Ponomarev V.N. and Pronin P.I., *Problems of classical and quantum collapse X, Massless scalarparticle production through the torsion field*, *Sov. J. Phys.* **21** (1978) 1197-1200 [*Izvestiya VUZov, Fizika*, n. 9 (1978) 105-109 (in Russian)].
2391. Ponomarev V.N. and Pronin P.I., *Creation of massless scalar particles by the torsion field*, *Theor. Math. Phys.* **39** (1979) 559-561 [*Teor. Mat. Fiz.* **39** (1979) 425-428 (in Russian)].
2392. Ponomarev V.N. and Pronin P.I., *Spontaneous symmetry breaking by the torsion field and nonlinear spinor theorems*, *Sov. J. Phys.* **22** (1979) 1324-1325 [*Izvestiya VUZov, Fizika*, n. 12 (1979) 81-82 (in Russian)].
2393. Ponomarev V.N. and Smetanin E.V., *New variant of nonlinear electrodynamics. I*, *Sov. J. Phys.* **21** (1978) 1177-1181 [*Izvestiya VUZov, Fizika*, n. 9 (1978) 81-87 (in Russian)].
2394. Ponomarev V.N. and Smetanin E.V., *New variant of nonlinear electrodynamics. II*, *Sov. J. Phys.* **21** (1978) 1288-1293 [*Izvestiya VUZov, Fizika*, n. 10 (1978) 46-53 (in Russian)].
2395. Ponomarev V.N. and Smetanin E.V., *New version of nonlinear electrodynamics. I*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 5 (1978) 29-37 (in Russian).
2396. Ponomarev V.N. and Smetanin E.V., *New version of nonlinear electrodynamics. II*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 6 (1978) 3-8 (in Russian).
2397. Ponomarev V.N. and Tseytlin A.A., *On the correct usage of the Palatini principle in the gravitational theories*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **19**, n. 6 (1978) 57-59 (in Russian).
2398. Ponomarev V.N. and Tseytlin A.A., *Microscopic gravity as a Yang-Mills field*, *Phys. Lett.* **A70** (1979) 164-166.
2399. Ponomarev V.N. and Tseytlin A.A., *On variational principle in general relativity*, *Vestnik Mosk. Univ., ser. fiz., astr.*, **20**, n. 5 (1979) 68-80 (in Russian).
2400. Ponomarev V.N. and Tseytlin A.A., *Gauge approach in gravity*, in: "Gravitation and General Relativity", Ed. V.R. Kaigorodov (Kazan Univ. Publ. House: Kazan, 1983) **19**, 124-133 (in Russian).
2401. Poplawski N., *Interacting dark energy in  $f(R)$  gravity*, *Phys. Rev.* **D74** (2006) 084032 (8 pages).

2402. Popławski N., *Propagating torsion in the Einstein frame*, *J. Math. Phys.* **47** (2006) 112504.
2403. Popławski N., *Acceleration of the universe in the Einstein frame of a metric-affine  $f(R)$  gravity*, *Class. Quantum Grav.* **23** (2006) 2011-2020.
2404. Popławski N., *The present universe in the Einstein frame, metric-affine  $R + 1/R$  gravity*, *Class. Quantum Grav.* **23** (2006) 4819-4828.
2405. Popławski N., *On the nonsymmetric purely affine gravity*, *Mod. Phys. Lett.* **A22** (2007) 2701-2020.
2406. Popławski N., *Massive vectors from projective invariance breaking*, *Annales de la Fondation L. de Broglie* **32** (2007) 335-353.
2407. Popławski N., *A variational formulation of relativistic hydrodynamics*, *Phys. Lett.* **A373** (2009) 2620-2621.
2408. Popławski N., *Gravitation, electromagnetism and cosmological constant in purely affine gravity*, *Found. Phys.* **39** (2009) 307-330.
2409. Popławski N., *Nonsingular Dirac particles in spacetime with torsion*, *Phys. Lett.* **B690** (2010) 73-77.
2410. Popławski N., *Cosmology with torsion: An alternative to cosmic inflation*, *Phys. Lett.* **B694** (2010) 181-185.
2411. Popławski N., *Torsion as electromagnetism and spin*, *Int. J. Theor. Phys.* **49** (2010) 1481-1488.
2412. Popławski N., *Cosmological constant from quarks and torsion*, *Ann. d. Phys. (Berlin)* **523** (2011) 291-295.
2413. Popławski N., *Matter-antimatter asymmetry and dark matter from torsion*, *Phys. Rev.* **D83** (2011) 084033 (5 pages).
2414. Popławski N., *Nonsingular, big-bounce cosmology from spinor-torsion coupling*, *Phys. Rev.* **D85** (2012) 107502 (5 pages).
2415. Popławski N., *Four-fermion interaction from torsion as dark energy*, *Gen. Rel. Grav.* **44** (2012) 491-499.
2416. Popławski N., *Big bounce from spin and torsion*, *Gen. Rel. Grav.* **44** (2012) 1007-1014.
2417. Popławski N., *Cosmological consequences of gravity with spin and torsion*, *Astronomical Review*, **8** (2013) 108-115.
2418. Popławski N., *Affine theory of gravitation*, *Gen. Rel. Grav.* **46** (2014) 1625.
2419. Popławski N., *The energy and momentum of the Universe*, *Class. Quantum Grav.* **31** (2014) 065005.
2420. Popławski N., *Schwinger's variational principle in Einstein-Cartan gravity*, *Phys. Rev.* **D89** (2014) 027501 (3 pages).
2421. Popławski N., *Universe in a black hole in Einstein-Cartan gravity*, *Astrophys. J.* **832** (2016) 96 [8 pages].
2422. Popov D.A., *Theory of Yang-Mills fields*, *Theor. Math. Phys.* **24** (1975) 879-885 [*Teor. Mat. Fiz.* **24** (1975) 347-356 (in Russian)].
2423. Popov D.A. and Daikhin L.I., *Einstein spaces and Yang-Mills fields*, *Sov. Phys. Dokl.* **20** (1976) 818-820 [*Dokl. AN SSSR* **225** (1975) 790-793 (in Russian)].
2424. Popović D.S. and Sazdović B., *The geometrical form for the string space-time action*, *Eur. Phys. J.* **C50** (2007) 683-689.
2425. Poznanin P.L., *Quantization of torsion in a space of zero Riemannian curvature*, *Sov. J. Phys.* **18** (1975) 95-100 [*Izvestia VUZov, Fizika*, n. 1 (1975) 116-121 (in Russian)].

2426. Poznanin P.L., *Long-range interaction between spinors and a torsion field by means of photon exchange*, *Sov. J. Phys.* **18** (1975) 566-567 [*Izvestia VUZov, Fizika*, n. 4 (1975) 134-135 (in Russian)].
2427. Poznanin P.L., *Quantization of torsion in the weak field approximation*, in: "Probl. Theor. Grav. and Elem. Particles", Ed. K.P. Stanyukovich (Atomizdat: Moscow) **7** (1976) 94-109 (in Russian).
2428. Poznanin P.L., *Torsionic interactions in quantum theory*, in: "Advances in the relativity and gravity theory" (Nauka: Moscow, 1977) 86-90 (in Russian).
2429. Pradhan T., Malik R.P., and Naik P.C., *The fifth interaction: universal long range force between spins*, *Pramana J. Phys.* **24** (1985) 77-94.
2430. Prasanna A.R., *Cylindrically symmetric matter distribution with a magnetic field in Einstein-Cartan theory*, *Pramana* **5** (1975) 289-293.
2431. Prasanna A.R., *Maxwell's equations in Riemann-Cartan space  $U_4$* , *Phys. Lett.* **A54** (1975) 17-18.
2432. Prasanna A.R., *Static fluid spheres in Einstein-Cartan theory*, *Phys. Rev.* **D11** (1975) 2076-2082.
2433. Prasanna A.R., *Static cylinder of perfect fluid with nonzero spin density*, *Phys. Rev.* **D11** (1975) 2083-2087.
2434. Prasanna A.R., *Differential forms and Einstein-Cartan theory*, In: "Gravitation, gauge theories and the early universe", Eds. B.R. Iyer, C.V. Vishveshwara, and N. Mukunda. (Dordrecht: Kluwer, 1989), 119-132.
2435. Prasanna A.R. and Mohanty S., *Electromagnetic wave propagation in general spacetimes with curvature and/or torsion ( $U_4$ )*, in: "The Universe: Visions and Perspectives", Eds. N. Dadhich and A. Kembhavi (Springer: Amsterdam, 2000) 277-284.
2436. Prasanna A.R. and Mohanty S., *Photon propagation in torsion background*, *Gen. Rel. Grav.* **41** (2009) 1905-1908.
2437. Preuss O., Haugan M.P., Solanki S.K., and Jordan S., *An astronomical search for evidence of new physics: Limits on gravity-induced birefringence from the magnetic white dwarf RE J0317-853*, *Phys. Rev.* **D70** (2004) 067101 (4 pages).
2438. Preuss O., Solanki S.K., Haugan M.P., and Jordan S., *Gravity-induced birefringence within the framework of Poincaré gauge theory*, *Phys. Rev.* **D72** (2005) 042001 (6 pages).
2439. Prokof'ev A. V., *N cosmic strings in Riemann-Cartan space* *Vestnik Moskov. Univ. Ser. III Fiz. Astronom.* **36**, n. 3 (1995) 8-14 (in Russian).
2440. Pronin P.I., *The induced torsional long range interaction in the Einstein-Cartan theory*, in: "Relativistic Astrophysics and Cosmology. Proc. of the Sir A.Eddington Centenary Symp.", Eds. V. DeSabbata and T.M. Karade (World Scientific: Singapore, 1984) vol. **1**, 137-169.
2441. Pronin P.I., *Renormalization of nonabelian gauge fields in the Riemann-Cartan space-time*, *Hadronic J.* **11** (1988) 291-298.
2442. Pronin P.I., *How we can measure torsion?*, in: "Experimental tests of gravity", Eds. V.B. Braginsky and V.I. Denisov (Moscow Univ. Publ. House: Moscow, 1989) 146-152 (in Russian).
2443. Pronin P.I., *Renormalization of field theories in Riemann-Cartan space-time*, in: *Proc. NATO Adv. Res. Workshop "Quantum Mechanics in Curved Space-time" (Erice, Italy, 1989)*, Eds. J. Audretsch and V. De Sabbata (Plenum Press: New York, 1990) 517-550.
2444. Pronin P.I., *Physical effects and measurability of spin-torsion interaction*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?", *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. de Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 173-198.

2445. Puetzfeld D., *A plane-fronted wave solution in metric-affine gravity*, in: "Exact Solutions and Scalar Fields in Gravity: Recent Developments", Eds. Macias A., Cervantes-Cota J., and Lämmerzahl C. (Springer: NY, 2001) 141-151.
2446. Puetzfeld D., *A cosmological model in Weyl-Cartan spacetime: I. Field equations and solutions*, *Class. Quantum Grav.* **19** (2002) 3263.
2447. Puetzfeld D., *A cosmological model in Weyl-Cartan spacetime: II. Magnitude-redshift relation*, *Class. Quantum Grav.* **19** (2002) 4463.
2448. Puetzfeld D., *A non-standard cosmological model*, *Int. J. Mod. Phys.* **A17** (2002) 2772.
2449. Puetzfeld D., *Prospects of Non-Riemannian Cosmology*, in: "22nd Texas Symposium on Relativistic Astrophysics at Stanford University" (Dec. 13-17, 2004) 1221 (5 pages).
2450. Puetzfeld D., *Status of non-Riemannian cosmology*, *New Astronomy Reviews* **49** (2005) 59-64.
2451. Puetzfeld D., *The motion of test bodies with microstructure in gauge gravity models*, *Acta Phys. Pol. B Proc. Suppl.* **1** (2008) 167-172.
2452. Puetzfeld D. and Chen X., *Testing non-standard cosmological models with supernovae*, *Class. Quantum Grav.* **21** (2004) 270.
2453. Puetzfeld D. and Obukhov Yu.N., *Propagation equations for deformable test bodies with microstructure in extended theories of gravity*, *Phys. Rev.* **D76** (2007) 084025 (20 pages).
2454. Puetzfeld D. and Obukhov Yu.N., *Probing non-Riemannian spacetime geometry*, *Phys. Lett.* **A372** (2008) 6711-6716.
2455. Puetzfeld D. and Obukhov Yu.N., *Equations of motion in gravity theories with nonminimal coupling: A loophole to detect torsion macroscopically?* *Phys. Rev.* **D88** (2013) 064025 (9 pages).
2456. Puetzfeld D. and Obukhov Yu.N., *Prospects of detecting spacetime torsion*, *Int. J. Mod. Phys.* **D23** (2014) 1442004 (6 pages).
2457. Puetzfeld D. and Obukhov Yu.N., *Equations of motion in metric-affine gravity: A covariant unified framework*, *Phys. Rev.* **D90** (2014) 084034 (13 pages).
2458. Puetzfeld D., Pohl M., and Zhu Z.-H., *Complementary constraints from Fanaroff-Riley type IIb radio galaxies and X-ray gas mass fractions in clusters on nonstandard cosmological models*, *Astrophys. J.* **619** (2005) 657.
2459. Puetzfeld D. and Tresguerres R., *A cosmological model in Weyl-Cartan spacetime*, *Class. Quantum Grav.* **18** (2001) 677-693.
2460. Pullin J., *The parallel transport of a vector: its physical meaning in three geometrical unified field theories*, *Gen. Relat. Grav.* **18** (1986) 1087-1091.
2461. Pullin J., *Massive solutions to Einstein-Cartan-Dirac theory*, *Ann. d. Phys.* **46** (1989) 559-560.
2462. Puntigam R.A., *Spinning cosmic strings as spacetime distortions*, in: "Gravity, Theoretical Physics and Computers", *Proc. of 6th Conf. on Theor. Phys. - Gen. Rel. Grav., Bistrica, Romania, 1996*, Eds. D.N. Vulcanov and I.I. Cotaescu (Mirton Publ. House: Timisoara, 1997) 24-34.
2463. Puntigam R.A., Lämmerzahl C., and Hehl F.W., *Maxwell's theory on a post-Riemannian spacetime and the equivalence principle*, *Class. Quantum Grav.* **14** (1997) 1347-1356.
2464. Puntigam R.A. and Soleng H.H., *Volterra distortions, spinning strings, and cosmic defects*, *Class. Quantum Grav.* **14** (1997) 1129-1149.
2465. Purcell A.J., *Pseudoscalar action in a Cartan spacetime*, *Phys. Rev.* **D18** (1978) 2730-2732.

## Q

2466. Qiu X.M., Yan J., and Peng D.Y., *String theory and an integrable model in two-dimensional gravity with dynamical torsion*, *Gen. Rel. Grav.* **29** (1997) 1195-1207.
2467. Quin Si-Ming and Wu Yon-Lin,  *$T_4$  gauge theory of gravity and new general relativity*, *Gen. Relat. Grav.* **22** (1990) 511-524.

## R

2468. Radchenkov O.N., *Gauge model of gravity*, *Pis'ma ZhETF* **53** (1991) 514-517 (in Russian).
2469. Radovanovic V. and Šijački Dj., *Spacetime geometry of three-dimensional Yang-Mills theory*, *Class. Quantum Grav.* **12** (1995) 1791-1800.
2470. Ragusa S. and Bailyn M., *Traceless pole particles with intrinsic spin in space with torsion*, *Phys. Rev.* **D36** (1987) 1676-1678.
2471. Ramaswamy S. and Yasskin P., *Birkhoff theorem for an  $R + R^2$  theory of gravity with torsion*, *Phys. Rev.* **D19** (1979) 2264-2267.
2472. Randono A., *Canonical Lagrangian dynamics and general relativity*, *Class. Quantum Grav.* **25** (2008) 205017.
2473. Randono A., *A new perspective on covariant canonical gravity*, *Class. Quantum Grav.* **25** (2008) 235017.
2474. Randono A., *de Sitter spaces: topological ramifications of gravity as a gauge theory*, *Class. Quantum Grav.* **27** (2010) 105008.
2475. Randono A., *A mesoscopic quantum gravity effect*, *Gen. Rel. Grav.* **42** (2010) 1909-1917.
2476. Randono A., *Do spinors give rise to a frame-dragging effect?* *Phys. Rev.* **D81** (2010) 024027 (11 pages).
2477. Randono A. and Hughes T.L., *Torsional monopoles and torqued geometries in gravity and condensed matter*, *Phys. Rev. Lett.* **106** (2011) 161102 (4 pages).
2478. Randono A. and Sloan D., *Internal spin angular momentum of an asymptotically flat spacetime*, *Phys. Rev.* **D80** (2009) 044006 (16 pages).
2479. Rapoport D.L., *Stochastic processes in conformal Riemann-Cartan-Weyl gravitation*, *Int. J. Theor. Phys.* **30** (1991) 1497-1515.
2480. Rapoport D.L., *Riemann-Cartan-Weyl quantum geometry, I: Laplacians and supersymmetric systems*, *Int. J. Theor. Phys.* **35** (1996) 287-309.
2481. Rapoport D.L., *Riemann-Cartan-Weyl quantum geometry, II: Cartan stochastic copying method, Fokker-Planck operator and Maxwell-De Rham equations*, *Int. J. Theor. Phys.* **36** (1997) 2115-2152.
2482. Rapoport D.L., *Riemann-Cartan-Weyl geometries, quantum diffusions and the equivalence of the free Maxwell and Dirac-Hestenes equations*, *Adv. Appl. Clifford Alg.* **8** (1998) 129-146.
2483. Rapoport D.L., *Torsion, thermodynamical, quantum and hydrodynamical fluctuations*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997). Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 469-471.
2484. Rapoport D.L., *On the unification of geometric and random structures through torsion fields: Brownian motions, viscous and magneto-fluid-dynamics*, *Found. Phys.* **35** (2005) 1205-1244.

2485. Rapoport D.L., *Cartan-Weyl Dirac and Laplacian operators, Brownian motions: The quantum potential and scalar curvature, Maxwell's and Dirac-Hestenes equations, and supersymmetric systems*, *Found. Phys.* **35** (2005) 1383-1431.
2486. Rapoport D.L., *Torsion fields, Cartan-Weyl space-time and state-space quantum geometries, their Brownian motions, and the time variables*, *Found. Phys.* **37** (2007) 813-854.
2487. Rapoport D. and Sternberg S., *Classical mechanics without Lagrangians and Hamiltonians. II. The motion of a massive spinning particles in the presence of a flat metric and vector torsion*, *Nuovo Cim.* **A80** (1984) 371-383.
2488. Rapoport D. and Sternberg S., *On the interaction of spin and torsion*, *Ann. Phys. (USA)* **158** (1984) 447-475.
2489. Rapoport D., Rodrigues W., de Souza Q., and Vaz J., *The Einstein-Cartan effective geometry associated to Dirac-Hestenes spinor field*, *Algebras, Groups and Geometries* **14** (1994) 23-35.
2490. Rauch R.T., *Asymptotic flatness, reflection symmetry, and Birkhoff's theorem for  $R + R^2$  actions containing quadratic torsion*, *Phys. Rev.* **D25** (1982) 577-580.
2491. Rauch R.T., *Equivalence of an  $R + R^2$  theory of gravity to Einstein-Cartan-Sciama-Kibble theory in the presence of matter*, *Phys. Rev.* **D26** (1982) 931-933.
2492. Rauch R.T., *Time dependent gravitational coupling and torsion*, *Phys. Rev. Lett.* **52** (1984) 1842-1844.
2493. Rauch R. and Nieh H.T., *Birkhoff's theorem for general Riemann-Cartan  $R + R^2$  theories of gravity*, *Phys. Rev.* **D24** (1981) 2029-2048.
2494. Rauch R.T., Shaw S.J., and Nieh H.T., *Birkhoff's theorem for ghost-free tachyon-free  $R + R^2 + Q^2$  theories with torsion*, *Gen. Relat. Grav.* **14** (1982) 331-354.
2495. Ray J.R. and Smalley L.L., *Spinning fluids in general relativity*, *Phys. Rev.* **D26** (1982) 2619-2622.
2496. Ray J.R. and Smalley L.L., *Perfect fluids in the Einstein-Cartan theory*, *Phys. Rev.* **D26** (1982) 2615-2618.
2497. Ray J.R. and Smalley L.L., *Improved perfect fluid energy-momentum tensor with spin in Einstein-Cartan space-time*, *Phys. Rev. Lett.* **49** (1982) 1059-1061.
2498. Ray J.R. and Smalley L.L., *Spinning fluids in the Einstein-Cartan theory*, *Phys. Rev.* **D27** (1983) 1383-1385.
2499. Ray J.R. and Smalley L.L., *Consistency relations for spinning matter in gravitational theories*, *Phys. Rev.* **D34** (1986) 3268-3269.
2500. Ray J.R. and Smalley L.L., *Equation of state spinning fluids in the Einstein-Cartan theory*, *Phys. Rev.* **D35** (1987) 1185-1188.
2501. Ray J.R., Smalley L.L., and Krisch J., *Spinning fluids in general relativity. II, Self-consistent formulation*, *Phys. Rev.* **D35** (1987) 3261-3262.
2502. Ray S., *Static spherically symmetric electromagnetic mass models with charged dust sources in Einstein-Cartan theory: Lane-Emden models*, *Astropys. and Space Sci.* **280** (2002) 345-355.
2503. Raychaudhuri A.K., *Einstein-Cartan cosmologies with a magnetic field*, *Phys. Rev.* **D12** (1975) 952-955.
2504. Raychaudhuri A.K., *Theoretical cosmology* (Clarendon Press: Oxford, 1979), chapter 10, 182-198.
2505. Raychaudhuri A.K. and Banerji S., *Einstein-Cartan spheres*, *Phys. Rev.* **D16** (1977) 281-285.

2506. Reid J.A. and Wang C.H.T., *Conformal holonomy in MacDowell-Mansouri gravity*, *J. Math. Phys.* **55** (2014) 032501.
2507. Reifler F. and Morris R., *Hestenes' tetrad and spin connections*, *Int. J. Theor. Phys.* **44** (2005) 1307-1324.
2508. Richter O. and Rudolph G., *Torsion and spontaneous compactification*, *Int. J. Mod. Phys.* **A8** (1993) 1903-1921.
2509. Richter O. and Rudolph G., *On dimensional reduction of gravity with torsion*, *Class. Quant. Grav. Suppl.* **10** (1993) S127-S134.
2510. Ringermacher H.I., *An electrodynamic connection*, *Class. Quantum Grav.* **11** (1994) 2383-2394.
2511. Rizaoglu E., *Two-dimensional nonlinear fermionic models in curved space with torsion*, *Class. Quantum Grav.* **14** (1997) 1771-1779.
2512. Rocha-Neto J.F. da, and Maluf J.W., *The angular momentum of plane-fronted gravitational waves in the teleparallel equivalent of general relativity*, *Gen. Rel. Grav.* **46** (2014) 1667.
2513. Rocha R. da, Fabbri L., da Silva J.M.H., Cavalcanti R.T., and Silva-Neto J.A., *Flag-dipole spinor fields in ESK gravities*, *J. Math. Phys.* **54** (2013) 102505.
2514. Rocha R. da, and Rodrigues W.A. Jr., *Rigorous formulation of duality in gravitational theories*, *J. Phys. A: Math. Theor.* **43** (2010) 205206.
2515. Rochev V.E., *Massive vector field as a source of space-time torsion*, *Teor. Mat. Fiz.* **18** (1974) 226-232 (in Russian) [*Theor. Math. Phys.* **18** (1974) 160-164].
2516. Rodichev V.I., *Space with torsion and nonlinear field equations*, *ZhETF* **40** (1961) 1469-1472 (in Russian) [*Sov. Phys. JETP* **13** (1961) 1029-1031].
2517. Rodichev V.I., *Theory of gravity in orthogonal repère* (Nauka: Moscow, 1974) 184 p. (in Russian).
2518. Rodrigues M.E., Hamani Daouda M., Houndjo M.J.S., Myrzakulov R., and Sharif M., *Inhomogeneous universe in  $f(T)$  theory*, *Grav. Cosmol.* **20** (2014) 80-89.
2519. Rodrigues M.E., Houndjo M.J.S., Sáez-Gómez D., and Rahaman F., *Anisotropic universe models in  $f(T)$  gravity*, *Phys. Rev.* **D86** (2012) 104059 (13 pages).
2520. Rodrigues M.E., Houndjo M.J.S., Tossa J., Momeni D., and Myrzakulov R., *Charged black holes in generalized teleparallel gravity*, *JCAP* **11** (2013) 024.
2521. Rodrigues W.A., Jr., *Differential forms on Riemannian (Lorentzian) and Riemann-Cartan structures and some applications to physics*, *Annales de la Fondation L. de Broglie* **32** (2007) 425-478.
2522. Rodrigues W.A., Jr., and de Oliveira E.C., *The many faces of Maxwell, Dirac and Einstein equations: A Clifford bundle approach* (Springer: Berlin, 2007).
2523. Rodrigues W.A., Jr., de Souza Q.A.G., and Vaz J., Jr., *Lagrangian formulation in the Clifford bundle of Dirac-Hestenes equation on a Riemann-Cartan spacetime*, in: *Gravitation: the spacetime structure* (Águas de Lindóia, 1993) (World Scientific: River Edge, NJ, 1994), 534-543.
2524. Rodrigues W.A., Jr., de Souza Q.A.G., Vaz J., Jr., and Lounesto P., *Dirac-Hestenes spinor fields on Riemann-Cartan manifolds*, *Int. J. Theor. Phys.* **35** (1996) 1849-1900.
2525. Rohm R., *Quantized torsion*, In: "Unified string theories". *Proc. workshop, Santa Barbara, USA, July 29 - August 16, 1985*, Eds. M. Green, and D. Gross (Singapore: World Scientific, 1986), 719-726.
2526. Ronchetti L. and De Sabbata V., *Quantum gravity theory with torsion and spin fluctuations*, *Grav. Cosmol.* **8** (2002) 175-180.

2527. Rose B., *Field equations for gravity quadratic in curvature*, *Int. J. Theor. Phys.* **31** (1992) 1283-1293.
2528. Rosen N., *Weyl's geometry and physics*, *Found. Phys.* **12** (1982) 213-248.
2529. Rosenbaum M. and Luehr C.P., *Poincaré group gauge theory of gravitation in the fibre-bundle picture*, *Hadronic J.* **4** (1981) 153-163.
2530. Rosenbaum M., Ryan M., and Shepley L.C., *Spacetime homogeneous models with torsion*, *J. Math. Phys.* **20** (1979) 744-751.
2531. Rosenbaum M., Ryan M.P., Jr., and Urrutia L.F., *Dynamical torsion in a gravitational theory coupled to first order twist-tensor matter field*, *Phys. Rev.* **D26** (1982) 761-769.
2532. Ross D.K., *Gauge groups in a fiber bundle approach to superspace and the gravitational fields*, *J. Math. Phys.* **22** (1981) 557-561.
2533. Ross D.K., *Plank's constant, torsion and space-time defects*, *Int. J. Theor. Phys.* **28** (1989) 1333-1340.
2534. Ross D.K., *Torsion in fiber bundle models of gauge theories*, *Nuovo Cim.* **A109** (1996) 185-192.
2535. Ross D.K., *Normalization of quantized area using torsion and spin*, *Gen. Rel. Grav.* **33** (2001) 29-36.
2536. Rovelli C., *Ghosts in gravity theories with a scalar field*, *Nuovo Cim.* **B78** (1983) 167-177.
2537. Roy A. and Bandyopadhyay P., *Gravitational anomaly and the world topology*, *Mod. Phys. Lett.* **A7** (1992) 33-42.
2538. Rubilar G.F., *On the universality of Einstein(-Cartan) field equations in the presence of matter fields*, *Class. Quantum Grav.* **15** (1998) 239-244.
2539. Rubilar G.F., Obukhov Yu.N., and Hehl F.W., *Torsion nonminimally coupled to the electromagnetic field and birefringence*, *Class. Quantum Grav.* **20** (2003) L185-L191.
2540. Rudenko A.S. and Khriplovich I.B., *Gravitational four-fermion interaction in the early Universe*, *Phys. Usp.* **57** (2014) 167-170.
2541. Ruggiero M.L., *Light bending in  $f(T)$  gravity*, *Int. J. Mod. Phys.* **D25** (2016) 1650073 (12 pages).
2542. Ruggiero M.L. and Radicella N., *Weak-field spherically symmetric solutions in  $f(T)$  gravity*, *Phys. Rev.* **D91** (2015) 104014 (6 pages).
2543. Rumpf H., *On the translational part of the Lagrangian in the Poincaré gauge theory of gravitation*, *Z. Naturforsch.* **33a** (1978) 1224-1225.
2544. Rumpf H., *Creation of Dirac particles in general relativity with torsion and electromagnetism. I. The general formalism*, *Gen. Relat. Grav.* **10** (1979) 509-523.
2545. Rumpf H., *Creation of Dirac particles in general relativity with torsion and electromagnetism. II. The constant electric field - a pedagogical example*, *Gen. Relat. Grav.* **10** (1979) 525-533.
2546. Rumpf H., *Creation of Dirac particles in general relativity with torsion and electromagnetism. III. Matter production in a modes of torsion*, *Gen. Relat. Grav.* **10** (1979) 647-658.
2547. Rumpf H., *Quasiclassical limit of the Dirac equation and the equivalence principle in the Riemann-Cartan geometry*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 93-104.
2548. Rumpf H., *Supersymmetric Dirac particles in Riemann-Cartan space-time*, *Gen. Relat. Grav.* **14** (1982) 873-892.

2549. Russel N., *Lorentz violation and torsion*, in: "CPT and Lorentz Symmetry", Proc. 4th Meeting Indiana University, Bloomington, USA, 8-11 August 2007 (World Scientific: Singapore, 2007) pp. 282-286.
2550. Ryder L.H., *Introduction to general relativity* (Cambridge University Press: Cambridge, 2009) 441 p.
2551. Ryder L.H. and Shapiro I.L., *On the interaction of massive spinor particles with external electromagnetic and torsion fields*, Phys. Lett. **A247** (1998) 21-26.

## S

2552. Saa A., *On minimal coupling in Riemann-Cartan space-times*, Mod. Phys. Lett. **A8** (1993) 2565-2568.
2553. Saa A., *Gauge fields on Riemann-Cartan space-times*, Mod. Phys. Lett. **A9** (1994) 971-981.
2554. Saa A., *Volume-forms and minimal action principles in affine manifolds*, J. Geom. Phys. **15** (1995) 102-108.
2555. Saa A., *Propagating torsion from first principles*, Gen. Rel. Grav. **29** (1997) 205-220.
2556. Sabbir G., Khan A., Qureshi M.A., and Kara A.H., *A note on classification of teleparallel conformal symmetries in non-static plane symmetric space-times in the teleparallel theory of gravitation using diagonal tetrads*, Int. J. Geom. Methods Mod. Phys. **13** (2016) 1650046 (8 pages).
2557. Safko J.L. and Elston F., *Lagrange multipliers and gravitational theory*, J. Math. Phys. **17** (1976) 1531-1537.
2558. Safko J.L., Tsamparlis M., and Elston F., *Variational methods with torsion in general relativity*, Phys. Lett. **A60** (1977) 1-2.
2559. Safko J.L., Tsamparlis M., and Elson F., *Variational methods with torsion in general relativity (Errata)*, Phys. Lett. **A61** (1977) 493.
2560. Saitoh H., *Gravity as a gauge theory with spontaneously broken internal super-Poincaré symmetry*, Progr. Theor. Phys. **83** (1990) 1003-1024.
2561. Sakane E. and Kawai T., *Energy-momentum and angular momentum carried by gravitational waves in extended New General Relativity* Prog. Theor. Phys. **108** (2002) 615-639.
2562. Sakharov A.D., *The topological structure of elementary charges and CPT symmetry*, in: "Problems of Theoretical Physics", Volume dedicated to the memory of I.E. Tamm (Nauka: Moscow, 1972) 243-247 (in Russian); in: A.D. Sakharov: *Collected Scientific Works*, D. ter Haar et al., eds. (M. Dekker, New York 1982) 199-204.
2563. Salako I.G., Rodrigues M.E., Kpadonou A.V., Houndjo M.J.S., and Tossa J., *A  $\Lambda$ CDM model in  $f(T)$  gravity: reconstruction, thermodynamics and stability*, JCAP **11** (2013) 060 (22 pages).
2564. Salgado P., Cataldo M., and del Campo S., *Supergravity and the Poincaré group*, Phys. Rev. **D65** (2002) 084032 (7 pages).
2565. Salgado P., Cataldo M., and del Campo S., *Higher dimensional gravity invariant under the Poincaré group*, Phys. Rev. **D66** (2002) 024013 (6 pages).
2566. Salgado P., Rubilar G., Crisóstomo J., and del Campo S., *A note about teleparallel super-gravity*, Eur. Phys. J. **C44** (2005) 587-590.
2567. Samanta S., *Diffeomorphism symmetry in the Lagrangian formulation of gravity*, Int. J. Theor. Phys. **48** (2009) 1436-1448.
2568. Samokhvalov S.E. and Vanyashin V.S., *Group theory approach to unification of gravity with internal gauge interactions*, Class. Quantum Grav. **8** (1991) 2277-2282.

2569. Sandberg V.D., *Are torsion theories of gravitation equivalent to metric theories?*, *Phys. Rev.* **D12** (1975) 3013-3018.
2570. Saniga M., *Conformally symmetrical Abelian Higgs sunspot and nonmetricity of the sun's spacetime*, *Astrophys. and Space Sci.* **193** (1992) 155-159.
2571. Sardanashvily G.A., *Compensation, nonlinearity, and geometry*, *Sov. J. Phys.* **17** (1974) 1741-1745 [*Izvestia VUZov, Fizika*, n. 12 (1974) 128-134 (in Russian)].
2572. Sardanashvily G.A., *Compensation and nonlinear theory*, *Sov. J. Phys.* **18** (1975) 1645-1650 [*Izvestia VUZov, Fizika*, n. 12 (1974) 7-13 (in Russian)].
2573. Sardanashvily G.A., *Systems with interactions and fibre bundles*, *Sov. J. Phys.* **21** (1978) 4-7 [*Izvestia VUZov, Fizika*, n. 1 (1978) 10-14 (in Russian)].
2574. Sardanashvily G.A., *The gravity as goldstone field in the Lorentz gauge theory*, *Phys. Lett.* **A75** (1980) 257-258.
2575. Sardanashvily G., *What are the Poincaré gauge fields?*, *Czech. J. Phys.* **B33** (1983) 610-615.
2576. Sardanashvily G., *On the definition of gauge transformations in gauge theory*, *Ann. der Phys.* **41** (1984) 23-28.
2577. Sardanashvily G.A., *The gauge model of the fifth force*, *Acta Phys. Pol.* **B21** (1990) 583-587.
2578. Sardanashvily G.A., *Spontaneous symmetry breaking in the gauge gravitational theory*, *Preprint IC/90/73* (ICTP: Trieste, 1990) 1-22.
2579. Sardanashvily G.A., *Gauge theory of gravity*, in: "Modern Problems of Theoretical Physics. Festschrift for Professor D.Ivanenko", Eds. P.I. Pronin and Yu.N. Obukhov (World Scientific: Singapore, 1991) 75-99.
2580. Sardanashvily G.A., *Spontaneous symmetry breaking in gauge gravity*, in: "Perspectives of the unified theory", Eds. D.V. Gal'tsov, L.S.Kuz'menkov and P.I.Pronin (Moscow Univ. Publ. House: Moscow, 1991) 182-203.
2581. Sardanashvily G., *Gauge gravitation theory*, *Int. J. Theor. Phys.* **30** (1991) 721-735.
2582. Sardanashvily G., *Gauge Theory in Jet Manifolds*, (Hadronic Press: Palm Harbor, 1993).
2583. Sardanashvily G., *Gravity as Higgs field*, in: "New Frontiers in Gravitation", Ed. G.A. Sardanashvily (Hadronic Press: Palm Harbor, 1996) 299-336.
2584. Sardanashvily G., *Stress-energy-momentum tensors in constraint field theories*, *J. Math. Phys.* **38** (1997) 847-866.
2585. Sardanashvily G., *Stress-energy-momentum conservation law in gauge gravitation theory*, *Class. Quantum Grav.* **14** (1997) 1357-1370.
2586. Sardanashvily G., *Universal spin structure*, *Int. J. Theor. Phys.* **37** (1998) 1265-1287.
2587. Sardanashvily G., *Covariant spin structure*, *J. Math. Phys.* **39** (1998) 4874-4890.
2588. Sardanashvily G., *Classical gauge gravitation theory*, *Int. J. Geom. Methods Mod. Phys.* **08** (2011) 1869-1895.
2589. Sardanashvily G., *Classical Higgs fields*, *Theor. Math. Phys.* **181** (2014) 1599-1611.
2590. Sardanashvily G., *Gauge gravitation theory: Gravity as a Higgs field*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650086 [35 pages].
2591. Sardanashvily G. and Gogberashvily M., *The dislocation treatment of gauge fields of space-time translations*, *Mod. Phys. Lett.* **A2** (1987) 609-616.
2592. Sardanashvily G.A. and Ikhlov B.L., *Higgs vacuum in the gravity theory*, *Moscow University Physics Bulletin*, **27**, n. 2 (1986) 17-19 (in Russian).

2593. Sardanashvily G.A. and Timoshenko E.G., *Gauge model of the fifth force*, Moscow University Physics Bulletin, **31**, n. 4 (1990) 70-72 (in Russian).
2594. Sardanashvily G.A. and Zakharov O.A., *Gauge transformations in gravitation theory*, Acta Phys. Pol. **B20** (1989) 651-653.
2595. Sardanashvily G. and Zakharov O., *Gauge Gravitation Theory*, (World Scientific: Singapore, 1992).
2596. Sarkar S., *String theory backgrounds with torsion in the presence of fermions and implications for leptogenesis*, J. Phys.: Conf. Ser. **631** (2015) 012089 (9 pages).
2597. Satarov A.G. and Syromyatnikov A.G., *Some features of two approaches to the affine metric theory of gravitation*, Teor. Mat. Fiz. **92** (1992) 150-153 (in Russian); English translation: Theor. Math. Phys. **92** (1992) 799-801.
2598. Schaller P. and Strobl T., *Canonical quantization of two-dimensional gravity with torsion and the problem of time*, Class. Quantum Grav. **11** (1994) 331-346.
2599. Scherk J. and Schwarz J.H., *Dual models and the geometry of space-time*, Phys. Lett. **B52** (1974) 347-350.
2600. Schouten J.A., *Tensor analysis for physicists* (Clarendon Press: Oxford, 1951).
2601. Schouten J.A., *Ricci-calculus. An introduction to tensor analysis and its geometrical applications*, 2nd edition (Springer: Berlin, 1954), 516 p.
2602. Schouten J.A. and Struik D.J., *Einführung in die neuen Methoden der Differentialgeometrie. Bd. I. (Schouten) Algebra und Übertragungslehre* (Noordhoff: Groningen-Batavia, 1935).
2603. Schouten J.A. and Struik D.J., *Einführung in die neuen Methoden der Differentialgeometrie. Bd. II. (Struik) Geometrie* (Noordhoff: Groningen-Batavia, 1938).
2604. Schrödinger E., *The general unitary theory of the physical fields*, Proc. Roy. Irish Acad. **A49** (1943) 43-58.
2605. Schrödinger E., *The Earth's and the Sun's permanent magnetic fields in the unitary field theory*, Proc. Roy. Irish Acad. **A49** (1943) 135-148.
2606. Schrödinger E., *Unitary field theory: Conservation identities and relation to Weyl and Eddington*, Proc. Roy. Irish Acad. **A49** (1944) 237-244.
2607. Schrödinger E., *The union of the three fundamental fields (gravitation, meson, electromagnetism)*, Proc. Roy. Irish Acad. **A49** (1944) 275-287.
2608. Schrödinger E., *The affine connexion in physical field theories*, Nature **153** (1944) 572-575.
2609. Schrödinger E., *On distant affine connection*, Proc. Roy. Irish Acad. **A50** (1945) 143-154.
2610. Schrödinger E., *The general affine field laws*, Proc. Roy. Irish Acad. **A51** (1946) 41-50.
2611. Schrödinger E., *The relation between metric and affinity*, Proc. Roy. Irish Acad. **A51** (1947) 147-150.
2612. Schrödinger E., *The final affine field laws. I*, Proc. Roy. Irish Acad. **A51** (1947) 163-171.
2613. Schrödinger E., *The final affine field laws. II*, Proc. Roy. Irish Acad. **A51** (1948) 205-216.
2614. Schrödinger E., *The final affine field laws. III*, Proc. Roy. Irish Acad. **A52** (1948) 1-9.
2615. Schrödinger E., *On the differential identities of an affinity*, Proc. Roy. Irish Acad. **A54** (1951) 79-85.
2616. Schrödinger E., *Space-time structure* (Cambridge Univ. Press: Cambridge, 1950).

2617. Schrüfer E., Hehl F.W., and McCrea J.D., *Exterior calculus on a computer: the REDUCE-package EXCALC applied to general relativity and to Poincaré gauge theory*, *Gen. Relat. Grav.* **19** (1987) 197-218.
2618. Schücker T. and Tilquin A., *Torsion, an alternative to the cosmological constant?*, *Int. J. Mod. Phys.* **D21** (2012) 1250089 (9 pages).
2619. Schücker T. and ZouZou S.R., *On a weak Gauss law in general relativity and torsion*, *Class. Quantum Grav.* **29** (2012) 245009.
2620. Schucking E.L. and Surowitz E.J., *Einstein's Apple: Homogeneous Einstein Fields* (World Scientific: Singapore, 2015) 316 pages.
2621. Schütze D., *On unifying electromagnetism and gravitation without curvature*, *J. Math. Phys.* **26** (1985) 2596-2600.
2622. Schweizer M.A., *The gauge symmetries of gravitation*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity" (Erice, 1979), Eds. P.G.Bergmann and V.De Sabbata (Plenum: New York, 1980) 117-124.
2623. Schweizer M. and Straumann N., *Poincaré gauge theory of gravitation and the binary pulsar 1913+16*, *Phys. Lett.* **A71** (1979) 493-495.
2624. Schweizer M., Straumann N., and Wipf A., *Post-Newtonian generation of gravitational waves in a theory of gravity with torsion*, *Gen. Relat. Grav.* **12** (1980) 951-961.
2625. Sciamia D.W., *On a non-symmetric theory of pure gravitational field*, *Proc. Cambr. Philos. Soc.* **54** (1958) 72-80.
2626. Sciamia D.W., *On the interpretation of the Einstein-Schrödinger unified field theory*, *J. Math. Phys.* **2** (1961) 472-477.
2627. Sciamia D.W., *On the analogy between charge and spin in general relativity*, in: "Recent Developments in General Relativity" (Pergamon: Oxford, 1962) 415-439.
2628. Sciamia D.W., *The physical structure of general relativity*, *Rev. Mod. Phys.* **36** (1964) 463-469, 1103(E).
2629. Scipioni R., *Hidden symmetries in non-Riemannian gravity*, *Helv. Phys. Acta* **72** (1999) 211-219.
2630. Scipioni R., *Isomorphism between non-Riemannian gravity and Einstein-Proca-Weyl theories extended to a class of scalar gravity theories*, *Class. Quantum Grav.* **16** (1999) 2471-2478.
2631. Scipioni R., *Effective Levi-Civita dilaton theory from metric affine dilaton gravity*, *Phys. Lett.* **A259** (1999) 104-107.
2632. Scipioni R., *On the triviality of certain non-Riemannian models of gravitation*, *Phys. Lett.* **A269** (2000) 263-268.
2633. Scipioni R., *On the extension of the Obukhov theorem in non-Riemannian gravity. I*, *J. Math. Phys.* **41** (2000) 3028-3034.
2634. Seeger A., *Recent advances in the theory of defects in crystals*, *Physica status solidi (b)* **1** (1961) 669-698.
2635. Seitz M., *The gravitational moment densities of classical matter fields with spin*, *Ann. der Phys.* **41** (1984) 280-290.
2636. Seitz M., *On solutions of the Einstein-Cartan-Dirac theory*, *Class. Quantum Grav.* **2** (1985) 919-925.
2637. Seitz M., *Proca field in space-time with curvature and torsion*, *Class. Quantum Grav.* **3** (1986) 1265-1273.
2638. Seitz M., *A quadratic Lagrangian for the Poincaré gauge field theory of gravity*, *Class. Quantum Grav.* **3** (1986) 175-182.

2639. Sen A., *Gravity as a spin system*, *Phys. Lett.* **B119** (1982) 89-91.
2640. Sen D.K., *Fields and/or particles* (Ryerson Press / Academic Press: Toronto, 1968) 140 p.
2641. Sen D.K., and Vanstone J.R., *On Weyl and Lyra manifolds*, *J. Math. Phys.* **13** (1972) 990-993.
2642. Sengupta S., *Parity violation in a gravitational theory with torsion: A geometrical interpretation*, *Pramana* **53** (1999) 1115-1119.
2643. Sengupta S., *Dark energy from the gravity vacuum*, *Class. Quantum Grav.* **32** (2015) 195005.
2644. Sengupta S. and Sinha A., *Fermion helicity flip by parity violating torsion*, *Phys. Lett.* **B514** (2001) 109-113.
2645. Sert Ö. and Adak M., *Spinor coupling to the weak Poincaré gauge theory of gravity in three dimensions*, *Phys. Rev.* **D86** (2012) 124033 (6 pages).
2646. Setare M.R. and Mohammadipour N., *Can  $f(T)$  gravity theories mimic  $\Lambda$ CDM cosmic history*, *JCAP* **01** (2013) 015.
2647. Sezgin E., *A class of ghost-free gravity Lagrangians with massive or massless propagating torsion*, *Phys. Rev.* **D24** (1981) 1677-1680.
2648. Sezgin E. and van Nieuwenhuizen P., *New ghost-free gravity Lagrangians with propagating torsion*, *Phys. Rev.* **D21** (1980) 3269-3280.
2649. Shabanov S.V., *Constrained systems and analytical mechanics in spaces with torsion*, *J. Phys. A: Math. Gen.* **A31** (1998) 5177-5190.
2650. Shankar K.H., Balaraman A., and Wali K.C., *Metric theory of gravity with torsion in an extra dimension*, *Phys. Rev.* **D86** (2012) 024007 (13 pages).
2651. Shao C.-G. and Xu B.-Q., *ISO(3, 1) gauge theory of gravity*, *Int. J. Theor. Phys.* **4** (1986) 347-353.
2652. Shapiro I.L., *The interaction of quantized matter fields with torsion and metric: NJL model*, *Mod. Phys. Lett.* **A9** (1994) 729-733.
2653. Shapiro I.L., *Physical aspects of the space-time torsion*, *Phys. Repts.* **357** (2002) 113-213.
2654. Shapiro I.L., *Effective QFT and what it tells us about dynamical torsion*, in: "CPT and Lorentz Symmetry", *Proc. 5th Meeting Indiana University, Bloomington, USA, 28 June-2 July 2010*, Ed. Kostelecký V.A. (World Scientific: Singapore, 2010) p. 65-69.
2655. Shapiro I.L. and Teixeira P.M., *Quantum Einstein-Cartan theory with the Holst term*, *Class. Quantum Grav.* **31** (2014) 185002.
2656. Sharif M., *Comments on torsion, massive electrodynamics and gravitation induced by scalar fields*, *Astrophys. Space Sci.* **238** (1996) 309-310.
2657. Sharin Yu.A., *Physical interpretation of curved space with torsion*, *Sov. Phys. J.* **33** (1990) 861-864 [*Izvestia VUZov, Fizika*, n. 10 (1990) 59-62 (in Russian)].
2658. Sharin Yu.A., *Physical interpretation of a curvilinear space with torsion*, *Russ. Phys. J.* **52** (2009) 1113-1115.
2659. Shen J.-Q., *A gravitational constant and a cosmological constant in a spin-connection gravitational gauge field theory*, *J. Phys. A: Math. Theor.* **42** (2009) 155401.
2660. Shen J.-Q., *Gravitational gauge theory developed based on the Stephenson-Kilmister-Yang equation*, *Int. J. Theor. Phys.* **48** (2009) 1566-1582.
2661. Shestopal V.O., *Feasibility of describing the gravitational field as a space with absolute parallelism*, *Sov. J. Phys.* **16** (1973) 1692-1694 [*Izvestia VUZov, Fizika* n. 12 (1973) 83-85 (in Russian)].

2662. Shie K.-F., Nester J.M., and Yo H.-J., *Torsion cosmology and the accelerating universe*, *Phys. Rev.* **D78** (2008) 023522 (16 pages).
2663. Shipov G.I., *Tetrad field equations in the space of absolute parallelism*, *Sov. J. Phys.* **19** (1976) 798-800 [*Izvestia VUZov, Fizika* n. 6 (1976) 132-133 (in Russian)].
2664. Shipov G.I., *Nonlinear spinor equations in general relativity*, *Sov. J. Phys.* **20** (1977) 378-382 [*Izvestia VUZov, Fizika* n. 3 (1976) 121-125 (in Russian)].
2665. Shipov G.I., *Theory of gravitation in the space of absolute parallelism*, *Sov. J. Phys.* **20** (1977) 813-817 [*Izvestia VUZov, Fizika* n. 6 (1976) 142-147 (in Russian)].
2666. Shipov G.I., *Geometry of absolute parallelism in the theory of physical vacuum*, *Russ. Phys. J.* **37** (1994) 853-879 [*Izvestia VUZov, Fizika* n. 9 (1994) 64-88 (in Russian)].
2667. Shirafuji T., *Lorentz invariant theory of gravitation. Gravitational interaction of spin-1/2 particles*, *Progr. Theor. Phys.* **62** (1979) 802-822.
2668. Shirafuji T. and Nashed G.G.L., *Energy and momentum in the tetrad theory of gravitation*, *Progr. Theor. Phys.* **98** (1997) 1355-1370.
2669. Shirafuji T., Nashed G.G.L., and Hayashi K., *Energy of general spherically symmetric solution in tetrad theory of gravitation*, *Progr. Theor. Phys.* **95** (1996) 665-678.
2670. Shirafuji T., Nashed G.G.L., and Kobayashi Y., *Equivalence principle in the new general relativity*, *Progr. Theor. Phys.* **96** (1996) 933-947.
2671. Shirafuji T. and Suzuki M., *Gauge theory of gravitation. A unified formulation of Poincaré and (anti-)de Sitter gauge theories*, *Progr. Theor. Phys.* **80** (1988) 711-730.
2672. Shtanov Yu.V. and Yushchenko S.A., *Conformally invariant cosmology based on Riemann-Cartan spacetime*, *Class. Quantum Grav.* **11** (1994) 2455-2482.
2673. Šijački Dj., *The unitary irreducible representations of  $\overline{SL}(3, R)$* , *J. Math. Phys.* **16** (1975) 298-311.
2674. Šijački Dj., *Quark confinement and the short-range component of general affine gauge gravity*, *Phys. Lett.* **B109** (1982) 435-438.
2675. Šijački Dj., *Affine gauge theories of gravity and particle physics*, in: "Proc. of the 4th Marcel Grossmann Meeting on General relativity" / Ed. R. Ruffini (North Holland: Amsterdam, 1986) 693-703.
2676. Šijački Dj.,  *$\overline{SL}(n, R)$  spinors for particles, gravity and superstrings*, in: "Spinors in Physics and Geometry", Eds. A. Trautman and G. Furlan (World scientific: Singapore, 1988) 191-206.
2677. Šijački Dj., *All  $SL(3, R)$  ladder representations*, *J. Math. Phys.* **31** (1990) 1872-1876.
2678. Šijački Dj., *World spinors revisited*, *Acta Phys. Pol.* **B29** (1998) 1089-1097.
2679. Šijački Dj., *Towards hypergravity*, in: *Proc. of the 8th Marcel Grossmann meeting "On recent developments in theoretical and experimental general relativity, gravitation, and relativistic field theories"* (Jerusalem, 22-27 June 1997). Eds. T. Piran and R. Ruffini (World Scientific: Singapore, 1999) pt. A, pp. 482-484.
2680. Šijački Dj.,  *$\overline{SL}(4, R)$  embedding for a 3D world spinor equation*, *Class. Quantum Grav.* **21** (2004) 4575.
2681. Šijački Dj. and Ne'eman Y., *Algebra and physics of the unitary multiplicity-free representations of  $SL(4, R)$* , *J. Math. Phys.* **26** (1985) 2457-2464.
2682. Šijački Dj. and Ne'eman Y., *Hadrons in an  $\overline{SL}(4, R)$  classification. 2. Mesons and C-assignments, P-assignments*, *Phys. Rev.* **D47** (1993) 4133-4141.
2683. Singh K.D., *Subspaces of a space with torsion*, *Tensor* **6** (1956) 6-14.

2684. Singh P., *On axial vector torsion in vacuum quadratic Poincaré gauge field theory*, *Phys. Lett.* **A145** (1990) 7-10.
2685. Singh P., *On null tratorial torsion in vacuum quadratic Poincaré gauge field theory*, *Class. Quantum Grav.* **7** (1990) 2125-2130.
2686. Singh P. and Griffiths J.B., *On neutrino fields in Einstein-Cartan theory*, *Phys. Lett.* **A132** (1988) 88-90.
2687. Singh P. and Griffiths J.B., *The application of spin coefficient techniques in the vacuum quadratic Poincaré gauge field theory*, *Gen. Relat. Grav.* **22** (1990) 269-287.
2688. Singh P. and Griffiths J.B., *A new class of exact solutions of the vacuum quadratic Poincaré gauge field theory*, *Gen. Relat. Grav.* **22** (1990) 947-956.
2689. Singh P. and Ryder L.H., *Einstein-Cartan-Dirac theory in the low-energy limit*, *Class. Quantum Grav.* **14** (1997) 3513-3525.
2690. Singh P., *On metrical Finsler connections with torsion of generalized Finsler spaces*, *Indian J. Pure Appl. Math.* **25** (1994) 399-404.
2691. Singh T., Srivastava R.S., and Helmi A.M., *Cylindrically-symmetric matter distribution with a magnetic field in Einstein-Cartan theory*, *Astropys. and Space Sci.* **178** (1991) 277-285.
2692. Singh T. and Yadaw R.B.S., *Some exact solutions of charged fluid spheres in Einstein-Cartan theory*, *Acta Phys. Pol.* **B9** (1978) 831-836.
2693. Singh T. and Yadaw R.B.S., *Static fluid spheres in Einstein-Cartan theory*, *Acta Phys. Pol.* **B9** (1978) 837-847.
2694. Sippel R. and Goenner H., *Symmetry classes of pp-waves*, *Gen. Rel. Grav.* **18** (1986) 1229-1243.
2695. Sivaram C. and DeSabbata V., *Maximum acceleration and magnetic field in the early universe*, *Astropys. and Space Sci.* **176** (1991) 145-148.
2696. Sivaram C. and Garcia De Andrade L.C., *Astrophysical limits on gauge invariance breaking in electrodynamics with torsion*, *Astropys. and Space Sci.* **201** (1993) 121-123.
2697. Sivaram C. and Garcia De Andrade L.C., *Gravitating systems with torsion self-interaction*, *Astropys. and Space Sci.* **201** (1993) 131-133.
2698. Sivaram C. and Garcia De Andrade L.C., *Torsion and massive electrodynamics in the early universe*, *Gen. Relat. Grav.* **26** (1994) 615-618.
2699. Sivaram C. and Garcia De Andrade L.C., *Torsion modified plasma screening in astrophysics*, *Mod. Phys. Lett.* **A15** (2000) 1275-1277.
2700. Skinner R. and Gregorash D., *Generalized Einstein-Cartan field equations*, *Phys. Rev.* **D14** (1976) 3314-3321.
2701. Skinner R. and Webb I., *Junction conditions for the Einstein-Cartan theory*, *Acta Phys. Pol.* **B8** (1977) 81-91.
2702. Skugoreva M.A., Saridakis E.N., and Toporensky A.V., *Dynamical features of scalar-torsion theories*, *Phys. Rev.* **D91** (2015) 044023 (11 pages).
2703. Slawianowski J.J., *GL(n, R) as a candidate for fundamental symmetry in field theory*, *Nuovo Cim.* **B106** (1991) 645-668.
2704. Slawianowski J.J., *Elimination of scale from the theory of mutually interacting gravitational and spinor fields*, *Rept. Math. Phys.* **33** (1993) 191-202.
2705. Slawianowski J.J., *Spinors, gravity and recalibration invariance. Microphysical motivation for the Weyl geometry*, *Rept. Math. Phys.* **35** (1995) 1-31.

2706. Slawianowski J.J., *New approach to the  $U(2, 2)$ -symmetry in spinor and gravitation theory*, *Fortschr. Phys.* **44** (1996) 105-141.
2707. Smalley L.L., *Variational principle for general relativity with torsion and non-metricity*, *Phys. Lett.* **A61** (1977) 436-438.
2708. Smalley L.L., *Comment on scalar-metric-torsion gravitational theories*, *Phys. Rev.* **D18** (1978) 3896-3897.
2709. Smalley L.L., *Volume preserving and conformal transformations in the metric affine gravitational theory*, *Lett. Nuovo Cim.* **24** (1979) 406-410.
2710. Smalley L.L., *Post-Newtonian approximation of the Poincaré gauge theory of gravity*, *Phys. Rev.* **D21** (1980) 328-331.
2711. Smalley L.L., *Riemann curvature tensor in non-holonomic coordinates and non-Riemann space-times*, *Int. J. Theor. Phys.* **23** (1984) 1001-1008.
2712. Smalley L.L., *Gödel cosmology in Riemann-Cartan space-time with spin density*, *Phys. Rev.* **D32** (1985) 3124-3127.
2713. Smalley L.L., *Brans-Dicke type models with nonmetricity*, *Phys. Rev.* **D33** (1986) 3590-3593.
2714. Smalley L.L., *Self-consistent Gödel cosmology with spin-density in Riemann-Cartan space-time*, *Phys. Lett.* **A113** (1986) 463-466.
2715. Smalley L.L., *On the extension of geometric optics from Riemannian to Riemann-Cartan spacetime*, *Phys. Lett.* **A117** (1986) 267-269.
2716. Smalley L.L., *Energy conditions in non-Riemannian spacetimes*, *Phys. Lett.* **A196** (1994) 147-153.
2717. Smalley L.L., *On the implications of perfect fluids in metric-affine spacetime*, *Class. Quantum Grav.* **17** (2000) 1447-1453.
2718. Smalley L.L. and Krisch J.P., *Energy-momentum tensor for minimally coupled charged perfect fluids with spin*, *Class. Quantum Grav.* **8** (1991) 1889-1894.
2719. Smalley L.L. and Krisch J.P., *Minimal coupling of electromagnetic fields in Riemann-Cartan space-times for perfect fluids with spin density*, *J. Math. Phys.* **33** (1992) 1073-1081.
2720. Smalley L.L. and Krisch J.P., *Minimal coupling of electromagnetic field in Riemann-Cartan spacetime for perfect fluid*, *Int. J. Theor. Phys.* **31** (1992) 1253-1265.
2721. Smalley L.L. and Krisch J.P., *Spinning fluid cosmologies in Einstein-Cartan theory*, *Class. Quantum Grav.* **11** (1994) 2375-2382.
2722. Smalley L.L. and Krisch J.P., *On spherically symmetric cosmology in Riemann-Cartan spacetime with spin-density*, *Class. Quantum Grav.* **11** (1994) 2517-2522.
2723. Smalley L.L. and Krisch J.P., *Energy conditions in non-Riemannian spacetimes*, *Phys. Lett.* **A196** (1994) 147-153.
2724. Smalley L.L. and Krisch J.P., *Fluids with spin and twist*, *J. Math. Phys.* **36** (1995) 778-795.
2725. Smalley L.L. and Krisch J.P., *String fluid dynamics*, *Class. Quantum Grav.* **13** (1996) L19-L22.
2726. Smalley L.L. and Krisch J.P., *Comparison of string fluid dynamical models in general relativity and Einstein-Cartan theory*, *Gen. Relat. Grav.* **31** (1999) 799-814.
2727. Smalley L.L. and Krisch J.P., *String fluid dynamical models in the Einstein-Cartan theory*, *Class. Quantum Grav.* **16** (1999) 1011-1019.
2728. Smalley L.L. and Ray J.R., *Geometrization of spin and the Weyssenhoff fluid conjecture*, *Gen. Relat. Grav.* **18** (1986) 549-553.

2729. Smalley L.L. and Ray J.R., *On the Lagrangian formulation of spinning fluids in Riemann-Cartan space-time*, *Phys. Lett.* **A134** (1988) 87-92.
2730. Smalley L.L. and Ray J.R., *Weysenhoff spinning fluid in Riemann-Cartan spacetime: direct variation with respect to the spin density of the fluid*, *Class. Quantum Grav.* **7** (1990) 1445-1461.
2731. Smetanin E.V., *Maxwell and Proca equations in space-time with torsion*, *Sov. J. Phys.* **25** (1982) 26-30 [*Izvestia VUZov, Fizika*, n. 1 (1982) 28-32 (in Russian)].
2732. Smetanin E.V., *Electromagnetic field in space-time with torsion. New solutions*, *Sov. J. Phys.* **25** (1982) 107-110 [*Izvestia VUZov, Fizika*, n. 2 (1982) 30-33 (in Russian)].
2733. Smetanin E.V., *Torsion and conservation laws*, *Sov. J. Phys.* **29** (1986) 123-125 [*Izvestia VUZov, Fizika*, n. 2 (1986) 50-52 (in Russian)].
2734. Smetanin E.V., *Magnetic monopole and dyon in space with torsion*, *Izvestia VUZov, Fizika*, n. 10 (1989) 121-123 (in Russian).
2735. Smrz P.K., *De Sitter gauge transformations of Dirac equations*, *Progr. Theor. Phys.* **57** (1977) 1771-1780.
2736. Smrz P.K., *Geometry of gauge translations*, *Lett. Nuovo Cim.* **38** (1983) 141-144.
2737. Smrz P.K., *Construction of space-time by gauge translations*, *J. Math. Phys.* **28** (1987) 2824-2828.
2738. Smrz P.K., *De Sitter gauge structure of space-time*, In: "Recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories". *Proc. of 5th Marcel Grossmann meeting, Perth, Australia, August 8-13, 1988*, Eds. D.G. Blair, and M.J. Buckingham (Singapore: World Scientific, 1989) Pt. **A**, 685-688.
2739. Smrz P.K., *Generalized linear connections on real manifolds*, *Gen. Relat. Grav.* **24** (1992) 857-865.
2740. Smrz P.K., *Einstein-Ötsuki vacuum equations*, *Gen. Relat. Grav.* **25** (1993) 33-40.
2741. Smrz P.K., *Geometrical models of elementary particles*, *Austral. J. Phys.* **48** (1995) 1045-1054.
2742. Smrz P.K., *Construction of real spacetime from complex linear metric connections*, *Austral. J. Phys.* **50** (1995) 793-802.
2743. So L.-L., *Gravitational energy from a combination of a tetrad expression and Einstein's pseudotensor*, *Class. Quantum Grav.* **25** (2008) 175012.
2744. Sobreiro R.F., Tomaz A.A., and Vasquez Otoya V.J., *de Sitter gauge theories and induced gravities*, *Eur. Phys. J.* **C72** (2012) 1991.
2745. Sobreiro R.F., Tomaz A.A., and Vasquez Otoya V.J., *Induced gravity from gauge theories*, *J. Phys.: Conf. Ser.* **453** (2013) 012014 (17 pages).
2746. Sobreiro R.F. and Vasquez Otoya V.J., *On the topological reduction from the affine to the orthogonal gauge theory of gravity*, *J. Geom. Phys.* **61** (2011) 137-150.
2747. Sobreiro R.F. and Vasquez Otoya V.J., *Affine gauge theory of gravity and its reduction to the Riemann-Cartan geometry*, *J. Phys.: Conf. Ser.* **283** (2011) 012032.
2748. Socolovsky M., *Locally inertial coordinates with totally antisymmetric torsion*, *Annales de la Fondation L. de Broglie* **37** (2012) 73-81.
2749. Socorro, J., Lämmerzahl C., Macías A., and Mielke E.W., *Multipole-like solutions in metric-affine gravity*, *Phys. Lett.*, **A244** (1998) 317-323.
2750. Socorro, J., Macías A., and Hehl F.W., *Computer algebra in gravity: Reduce-Excalc programs for (Non-)Riemannian space-times. I*, *Comp. Phys. Commun.* **115** (1998) 264-283.

2751. Soffel M., Müller B., and Greiner W., *Vacuum with spin in the ECKS-theory of gravitation*, *Phys. Lett.* **A70** (1979) 167-170.
2752. Solanki S.K., Preuss O., Haugan M.P., Gandorfer A., Povel H.P., Steiner P., Stucki K., Bernasconi P.N., and Soltan D., *Solar constraints on new couplings between electromagnetism and gravity*, *Phys. Rev.* **D69** (2004) 062001 (11 pages).
2753. Soleng H.H., *Scalar-tensor theory of gravitation with spin-scalar-torsion coupling*, *Class. Quantum Grav.* **5** (1988) 1489-1500.
2754. Soleng H.H., *Motions of photons in scalar-tensor theory of gravitation with spin-torsion coupling*, *Class. Quantum Grav.* **5** (1988) 1501-1511.
2755. Soleng H.H., *Generalized affine geometry and algebraically extended relativity: a new approach to spin-torsion coupling*, *Class. Quantum Grav.* **6** (1989) 785-795.
2756. Soleng H.H., *Gödel type cosmologies with spin density*, *Phys. Lett.* **A137** (1989) 826-828.
2757. Soleng H.H., *Spin-polarized cylinder in Einstein-Cartan theory*, *Class. Quantum Grav.* **7** (1990) 999-1007.
2758. Soleng H.H., *Torsion vector and variable G*, *Gen. Relat. Grav.* **23** (1991) 1089-1112.
2759. Soleng H.H., *A spinning string*, *Gen. Relat. Grav.* **24** (1992) 111-117.
2760. Soleng H.H., *Negative energy densities in extended sources generating closed timelike curves in general relativity with and without torsion*, *Phys. Rev.* **D49** (1994) 1124-1125.
2761. Soleng H.H. and Eeg J.O., *Skewons and gravitons*, *Acta Phys. Polon.* **B23** (1992) 87-114.
2762. Solodukhin S.N., *Black-hole solution in 2D gravity with torsion*, *Pis'ma ZhETF* **57** (1993) 317-322; English translation: *JETP Lett.* **57** (1993) 329-334.
2763. Solodukhin S.N., *Topological 2D Riemann-Cartan-Weyl gravity*, *Class. Quantum Grav.* **10** (1993) 1011-1021.
2764. Solodukhin S.N., *Two-dimensional black hole with torsion*, *Phys. Lett.* **B319** (1993) 87-95.
2765. Solodukhin S.N., *Cosmological solutions in 2D Poincaré gravity*, *Int. J. Mod. Phys.* **D3** (1994) 269-272.
2766. Solodukhin S.N., *On exact integrability of 2-D gravity*, *Mod. Phys. Lett.* **A9** (1994) 2817-2823.
2767. Solodukhin S.N., *Exact solution of two-dimensional Poincaré gravity coupled to fermion matter*, *Phys. Rev.* **D51** (1995) 603-608.
2768. Som M.M. and Bedran M., *Static dust sphere in Einstein-Cartan theory*, *Phys. Rev.* **D24** (1981) 2561-2561.
2769. Som M.M., Bedran M.L., and Amaral C.M., *Static perfect fluid in the Einstein-Cartan theory*, *Progr. Theor. Phys.* **67** (1982) 683-688.
2770. Som M.M., Bedran M.L., and Vaidya E.P.V., *Inhomogeneous cosmology with spinning fluid*, *Nuovo Cim.* **B102** (1988) 573-581.
2771. Soo C., *Adler-Bell-Jackiw anomaly, the Nieh-Yan term, and vacuum polarization*, *Phys. Rev.* **D59** (1999) 045006 (7 pages).
2772. Sotiriou T.P.,  *$f(R)$  gravity, torsion and non-metricity*, *Class. Quantum Grav.* **26** (2009) 152001.
2773. Sotiriou T.P. and Faraoni V.,  *$f(R)$  theories of gravity*, *Rev. Mod. Phys.* **82** (2010) 451-497.
2774. Sotiriou T.P., Li B., and Barrow J.D., *Generalizations of teleparallel gravity and local Lorentz symmetry*, *Phys. Rev.* **D83** (2011) 104030 (6 pages).

2775. Sotiriou T.P. and Liberati S., *The metric-affine formalism of  $f(R)$  gravity*, *J. Phys.: Conf. Ser.* **68** (2007) 012022.
2776. Sotiriou T.P. and Liberati S., *Metric-affine  $f(R)$  theories of gravity*, *Ann. Phys. (USA)* **322** (2007) .
2777. Sousa A.A. and Maluf J.W., *Canonical formulation of gravitational teleparallelism in  $2+1$  dimensions in Schwinger's time gauge*, *Progr. Theor. Phys.* **104** (2000) 531-543.
2778. Sousa A.A., Pereira R.B., and da Rocha-Neto J.F., *Angular momentum of the BTZ black hole in the teleparallel geometry*, *Progr. Theor. Phys.* **114** (2007) 1179-1190.
2779. Sousa A.A., Pereira R.B., and Silva A.C., *Energy and angular momentum densities in a Gödel-type universe in teleparallel geometry*, *Grav. Cosmol.* **16** (2010) 25-33.
2780. Spaniol E.P., Belo L.R.A., de Andrade V.C., and de Deus J.A., *The role of observers in the measurement of teleparallel gravitoelectromagnetic fields in Schwarzschild spacetime*, *Grav. Cosmol.* **19** (2013) 85-91.
2781. Spinoso R., *Proca test field in a space-time with torsion*, *Class. Quantum Grav.* **4** (1987) 473-484.
2782. Spinoso R., *On minimal coupling of classical spinning matter fields in space-time with torsion*, *Phys. Lett.* **A125** (1987) 227-230.
2783. Spinoso R., *Spin- $\frac{3}{2}$  test field in a space-time with torsion*, *Class. Quantum Grav.* **4** (1987) 1799-1808.
2784. Spinoso R., *Über Spin  $\frac{3}{2}$  Felder in Riemann Raum-Zeiten mit Torsion*, *Ann. der Phys. bf* **46** (1989) 179-188.
2785. Stachel J., *On the interpretation of the Einstein-Cartan formalism*, in: "On Einstein's path", *Essays in honor of E.Schucking*, Ed. A. Harvey (Springer: New York, 1998) 475-485.
2786. Stachowiak T. and Szydłowski M., *Exact solutions in bouncing cosmology*, *Phys. Lett.* **B646** (2007) 209-214.
2787. Stavrinou P.C. and Kouretsis A.P., *FRW-metric and Friedmann equations in a generalized cosmological model*, *J. Phys.: Conf. Ser.* **68** (2007) 012023.
2788. Stedile E., *Geometrical gauge approach for electromagnetism and gravitation*, *Phys. Lett.* **A118** (1986) 439-442.
2789. Stedile E., *Riemannian approach for gravitational Yang-Mills equations in the macroscopic case*, *Rev. Bras. Fis.* **69** (1989) 584-590.
2790. Stedile E. and Duarte R., *Yang-Mills gauge theory and gravitation*, *Int. J. Theor. Phys.* **34** (1995) 945-950.
2791. Stelle K.S. and West P.C., *De Sitter gauge invariance and the geometry of the Einstein-Cartan theory*, *J. Phys.* **A12** (1979) L205-L210.
2792. Stelle K.S. and West P.C., *Spontaneously broken de Sitter symmetry and the gravitational holonomy group*, *Phys. Rev.* **D21** (1980) 1466-1481.
2793. Stepanov V.E., *Lie differentiation for the Cartan spinors in a space-time of the affine connection with the Weyl type of nonmetricity*, *Russ. Phys. J.* **37** (1994) 1098-1101 [*Izvestia VUZov, Fizika* n. 11 (1994) 102-106 (in Russian)].
2794. Stephenson G., *Variational principles and gauge theories of gravitation*, *J. Phys.* **A10** (1977) 181-184.
2795. Sternberg S., *The interaction spin and torsion. II. The principle of general covariance*, *Ann. Phys. (USA)* **162** (1985) 85-99.
2796. Stoeger W.R., *The physics of detecting torsion and placing limits on its effects*, *Gen. Relat. Grav.* **17** (1985) 981-988.

2797. Straumann N., *General relativity* (Springer: Dordrecht, 2013) 735 p.
2798. Strobl T., *All symmetries of non-Einsteinian gravity in  $d = 2$* , *Int. J. Mod. Phys.* **8** (1993) 1383-1397.
2799. Strobl T., *Comment on "Gravity and the Poincaré group"*, *Phys. Rev.* **D48** (1993) 5029-5031.
2800. Stueckelberg E.C.G., *A possible new type of spin-spin interactions*, *Phys. Rev.* **73** (1948) 808.
2801. Sundermeyer K., *Symmetries in fundamental physics* (Springer: Berlin, 2014).
2802. Suè M., *Involutive systems of differential equations: Einstein's strength versus Cartan's degre d'arbitraire*, *J. Math. Phys.* **32** (1991) 392-399.
2803. Suè M. and Mielke E., *The strength of the Poincaré gauge field equations in first order formalism*, *Phys. Lett.* **A139** (1989) 21-26.
2804. Suh Y.B., *Remarks on the static spherically symmetric solutions in Einstein-Cartan theory*, *Progr. Theor. Phys.* **59** (1978) 1852-1859.
2805. Suh Y.B., *The path of a sound signal in Einstein and Einstein-Cartan theory*, *Phys. Lett.* **A84** (1981) 454-458.
2806. Sukhov A.M., *Energy-momentum tensor and transformation properties of the gauge potentials in gravitation theory*, *J. Math. Phys.* **32** (1991) 1035-1036.
2807. Sundermeyer K., *Extra "gauge" symmetries in  $R + T^2$  gravitational theories*, *Phys. Lett.* **B134** (1984) 415-418.
2808. Sur S. and Bhatia A.S., *Constraining torsion in maximally symmetric (sub)spaces*, *Class. Quantum Grav.* **31** (2014) 025020.
2809. Syromyatnikov A.G., *A problem of gauge invariance in the field theory with dynamical torsion*, *Teor. Mat. Fiz.* **87** (1991) 157-160 (in Russian); English translation: *Theor. Math. Phys.* **87** (1991) 444-446.
2810. Syromyatnikov A.G., *The  $g - 2$  muon anomaly in di-muon production with the torsion in LHC*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650093 (27 pages).
2811. Szczyrba W., *Field equations and contracted Bianchi identities in the generalized Einstein-Cartan theory*, *Lett. Math. Phys.* **2** (1978) 265-274.
2812. Szczyrba W., *The canonical variables, the symplectic structure and the initial value formulation of the generalized Einstein-Cartan theory of gravity*, *Commun. Math. Phys.* **60** (1978) 215-232.
2813. Szczyrba W., *Contracted Bianchi identities and conservation laws in Poincaré gauge theories of gravity*, in: "Proc. of the 6th Course of Internat. School on Cosmology and Gravitation: Spin, Torsion, Rotation and Supergravity" (Erice, 1979), Eds. P.G. Bergmann and V. DeSabbata (Plenum: New York, 1980) 105-116.
2814. Szczyrba W., *Hamiltonian dynamics of gauge theories of gravity*, *Phys. Rev.* **D25** (1982) 2548-2568.
2815. Szczyrba W., *The dynamical structure of the Einstein-Cartan-Sciama-Kibble theory of gravity*, *Ann. Phys. (USA)* **158** (1984) 320-373.
2816. Szczyrba W., *Stephenson-Kilmister-Yang theory of gravity and its dynamics*, *Phys. Rev.* **D36** (1987) 351-374.
2817. Szczyrba W., *Dynamics of quadratic Lagrangians in gravity. Fairchild's theory*, *Contemp. Math.* **71** (1988) 167-180.
2818. Szczyrba W. and Antonowicz M., *The dynamical structure of gravitational theories with  $GL(4, R)$  connections*, *J. Math. Phys.* **26** (1985) 1711-1727.
2819. Szydłowski M. and Krawiec A., *Cosmological model with macroscopic spin fluid*, *Phys. Rev.* **D70** (2004) 043510 (12 pages).

*T*

2820. Tafel J., *Non singular homogeneous universe with torsion*, *Phys. Lett.* **A45** (1973) 341-343.
2821. Tafel J., *A class of cosmological models with torsion and spin*, *Acta Phys. Pol.* **B6** (1975) 537-554.
2822. Tafel J., *Cosmological models with a spinor field*, *Bull. Acad. Pol. Sci. sér. sci. math., astr. phys.*, **25** (1977) 593-596
2823. Talebaoui W., *Theory of self-interacting scalar fields and gravitation*, *Mod. Phys. Lett.* **A8** (1993) 2053-2060.
2824. Tamanini N. and Böhmer C.G., *Good and bad tetrads in  $f(T)$  gravity*, *Phys. Rev.* **D86** (2012) 044009 (12 pages).
2825. Tapia V. and Ujevic M., *Universal field equations for metric-affine theories of gravity*, *Class. Quantum Grav.* **15** (1998) 3719-3729.
2826. Tartaglia A., *Space-time defects as a source of curvature and torsion*, *Int. J. Mod. Phys.* **A20** (2005) 2336.
2827. Tasson J.D., *Lorentz violation, gravitomagnetism, and intrinsic spin*, *Phys. Rev.* **D86** (2012) 124021 (5 pages).
2828. Tasson J.D., *Antimatter, the SME, and gravity*, *Hyperfine Interactions* **213** (2012) 137-146.
2829. Tasson J.D., *What do we know about Lorentz invariance?* *Rep. Prog. Phys.* **77** (2014) 062901.
2830. Tchraikian D.H., *On the dimensional reduction of gravity with torsion*, *Class. Quantum Grav.* **4** (1987) L217-L224.
2831. Teixeira A.F., da F., *Homogeneous, nonsingular, closed Einstein-Cartan cosmological model*, *Phys. Rev.* **D31** (1985) 2132-2134.
2832. Teyssander P., *Linearized  $R+R^2$  gravity: a new gauge exact solution*, *Class. Quantum Grav.* **6** (1989) 219-229.
2833. Teyssander P. and Tucker R.W., *Gravity, gauges and clocks*, *Class. Quantum Grav.* **13** (1996) 145-152.
2834. Teyssander P., Tucker R.W., and Wang C., *On an interpretation of non-Riemannian gravitation*, *Acta Phys. Pol.* **29** (1998) 987-994.
2835. Thienel H.-P., *BRST approach to translational symmetry and the geometry of flat manifolds with torsion*, *Gen. Relat. Grav.* **25** (1993) 483-490.
2836. Thirring W., *Gauge theories of gravitation*, in: "Facts and prospects of gauge theories", Ed. P. Urban (Springer: Wien, 1978) *Acta Phys. Austr. Suppl.* **29** (1978) 439-462.
2837. Thirring W., *Gauge theories of gravitation*, in: *Lect. Notes Phys.* **116** (1980) 272-275.
2838. Thirring W., *Classical field theory, a course of mathematical physics 2*, 2nd ed. (Springer: New York, 1986).
2839. Tiemblo A. and Tresguerres R., *Conformal symmetry and role of torsion in a Maxwell-Einstein system*, *Z. Phys.* **C51** (1991) 571-580.
2840. Tiemblo A. and Tresguerres R., *Invariant foliation of dynamical spacetimes*, *Gen. Relat. Grav.* **30** (1998) 239-262.
2841. Tiemblo A. and Tresguerres R., *Towards a general solution of the Hamiltonian constraints of General Relativity*, *Gen. Rel. Grav.* **39** (2007) 1839-1859.
2842. Tilquin A. and Schücker T., *Torsion, an alternative to dark matter?*, *Gen. Relat. Grav.* **43** (2011) 2965-2978.

2843. Timan H., *Variational principle with torsion*, *Ingenier* **5** (1970) 82.
2844. Timofeev V., *Dirac equation and optical scalars in the Einstein-Cartan theory*, *Int. J. Mod. Phys.: Conf. Ser.* **41** (2016) 1660123 (8 pages).
2845. Tiwari R.N. and Ray S., *Static spherical charged dust electromagnetic mass models in Einstein-Cartan theory*, *Gen. Rel. Grav.* **29** (1997) 683-690
2846. Tod K.P., *Conical singularities and torsion*, *Class. Quantum Grav.* **11** (1994) 1331-1339.
2847. Toller M., *Extended test particles in geometric fields*, *J. Math. Phys.* **24** (1983) 613-619.
2848. Toller M. and Vaia R., *A complete multipole expansion for a test particle in geometric fields*, *J. Math. Phys.* **25** (1984) 1039-1044.
2849. Toma N., *A Kerr metric solution in new general relativity*, *Progr. Theor. Phys.* **86** (1991) 659-670.
2850. Tonnelat M.A., *Einstein's unified field theory* (Gordon and Breach: NY, 1966) 186 pp.
2851. Tonnelat M.A., *The principles of electromagnetic theory and of relativity* (Reidel: Dordrecht, 1966) 476 pp.
2852. Toussaint M., *A gauge theoretical view of the charge concept in Einstein gravity*, *Gen. Rel. Grav.* **32** (2000) 885-896.
2853. Toussaint M., *A numeric solution for Einstein's gravitational theory with Proca matter and metric-affine gravity*, *Gen. Rel. Grav.* **32** (2000) 1689-1709.
2854. Townsend P.K., *Small-scale structure of spacetime as the origin of the gravitational constant*, *Phys. Rev.* **D15** (1977) 2795-2801.
2855. Trautman A., *Fiber bundles associated with space-time*, *Repts Math. Phys.* **1** (1970) 29-62.
2856. Trautman A., *On the Einstein-Cartan equations I*, *Bull. Acad. Polon. Sci., sér. sci. math. astr. phys.* **20** (1972) 185-190.
2857. Trautman A., *On the Einstein-Cartan equations II*, *Bull. Acad. Polon. Sci., sér. sci. math. astr. phys.* **20** (1972) 503-506.
2858. Trautman A., *On the Einstein-Cartan equations III*, *Bull. Acad. Polon. Sci., sér. sci. math. astr. phys.* **20** (1972) 895-896.
2859. Trautman A., *On the Einstein-Cartan equations IV*, *Bull. Acad. Polon. Sci., sér. sci. math. astr. phys.* **21** (1973) 345-346.
2860. Trautman A., *On the structure of the Einstein-Cartan equations*, *Symposia Mathematica* **12** (1973) 139-160.
2861. Trautman A., *Spin and torsion may avert gravitational singularity*, *Nature Phys. Sci.* **242** (1973) 7-8.
2862. Trautman A., *Theory of gravitation*, in: "The Physicists's Conception of Nature", Ed. J. Mehra (D. Reidel Publ.: Dordrecht, 1973) pp. 179-198.
2863. Trautman A., *Recent advances in the Einstein-Cartan theory of gravity*, *Ann. N.Y. Acad. Sci.* **262** (1975) 241-245.
2864. Trautman A., *A classification of space-time structures*, *Repts Math. Phys.* **10** (1976) 297-310.
2865. Trautman A., *The geometry of gauge fields*, *Czech. J. Phys.* **B29** (1979) 107-116.
2866. Trautman A., *Fiber bundles, gauge fields and gravitation*, in: "General Relativity and Gravitation: One Hundred Years after the Birth of Albert Einstein", Ed. A. Held (Plenum: New York, 1980) vol. **1**, 287-308.
2867. Trautman A., *On groups of gauge transformations*, *Lect. Notes Phys.* **129** (1980) 114-120.

2868. Trautman A., *Differential geometry for physicists* (Bibliopolis: Naples, 1984).
2869. Trautman A., *Gauge and optical aspects of gravitation*, *Class. Quantum Grav.* **16** (1999) A157-A175.
2870. Trautman A., *Einstein-Cartan theory*, in: *Encyclopedia of Math. Physics* J.-P. Francoise et al., eds. (Elsevier, Oxford, 2006) pp. 189-195.
2871. Treder H.-J., and von Borzeszkowski H.-H., *Covariance and quantum principles-censors of the space-time structure*, *Found. Phys.* **36** (2006) 757-763.
2872. Trefilov V.V. and Nelipa N.F., *Topological invariants in generalized theories of gravitation*, *Vestnik Mosk. Univ., ser. fiz., astr.* **24**, n 5 (1983) 71-74 (in Russian).
2873. Tresguerres R., *An exact solution of (2+1)-dimensional topological gravity in metric-affine spacetime*, *Phys. Lett.* **A168** (1992) 174-178.
2874. Tresguerres R., *Topological theory in a three-dimensional metric-affine space-time*, *J. Math. Phys.* **33** (1992) 4231-4238.
2875. Tresguerres R., *Weyl-Cartan model for cosmology before mass generation*, in: "Relativity in general" *Proc. of ERE-93 Spanish Relativity Meeting* (Salas, Asturias, Spaine, Sept. 7-10, 1993). Eds. J. Diaz Alonso, and M. Lorente Paramo. (Gif-sur-Yvette: Ed. Frontieres, 1994) 407-413.
2876. Tresguerres R., *Exact vacuum solutions of 4-dimensional metric-affine gauge theories of gravitation*, *Z. Phys.* **C65** (1995) 347-354.
2877. Tresguerres R., *Exact static vacuum solution of four-dimensional metric-affine gravity with nontrivial torsion*, *Phys. Lett.* **A200** (1995) 405-410.
2878. Tresguerres R., *Unified description of interactions in terms of composite fiber bundles*, *Phys. Rev.* **D66** (2002) 064025 (15 pages).
2879. Tresguerres R., *Translations and dynamics*, *Int. J. Geom. Methods Mod. Phys.* **05** (2008) 905-946.
2880. Tresguerres R., *Dynamically broken anti-de Sitter action for gravity*, *Int. J. Geom. Methods Mod. Phys.* **05** (2008) 171-183.
2881. Tresguerres R., *Motion in gauge theories of gravity*, *Int. J. Geom. Methods Mod. Phys.* **10** (2013) 1250085 (46 pages).
2882. Tresguerres R., *Thermodynamics in dynamical spacetimes*, *Phys. Rev.* **D89** (2014) 064032 (14 pages).
2883. Tresguerres R. and Mielke E.W., *Gravitational Goldstone fields from affine gauge theory*, *Phys. Rev.* **D62** (2000) 044004 (7 pages).
2884. Troncoso R. and Zanelli J., *Higher-dimensional gravity, propagating torsion and AdS gauge invariance*, *Class. Quantum Grav.* **17** (2000) 4451-4466.
2885. Trzęsowski A., *Geometrical and physical gauging in the theory of dislocations*, *Rep. Math. Phys.* **32** (1993) 71-98.
2886. Trzęsowski A., *Dislocations and internal length measurement in continuized crystals. 2. Closed teleparallelism*, *Int. J. Theor. Phys.* **33** (1994) 951-966.
2887. Tsamparlis M., *On the Palatini method of variation*, *J. Math. Phys.* **19** (1978) 555-557.
2888. Tsamparlis M., *Cosmological principle and torsion*, *Phys. Lett.* **A75** (1979) 27-28.
2889. Tsamparlis M., *Methods for deriving solutions in generalized theories of gravitation*, *Phys. Rev.* **D24** (1981) 1451-1457.
2890. Tse C.H., *Parallel torsions on Lie groups and spontaneous compactification in Kaluza-Klein theories*, *Phys. Lett.* **B128** (1983) 160-164.

2891. Tseng H.-H., Lee C.-C., and Geng C.-Q., *Scalar-torsion mode in a cosmological model of the Poincaré gauge theory of gravity*, *JCAP* **11** (2012) 013.
2892. Tseytlin A.A., *Poincaré and de Sitter gauge theories of gravity with propagating torsion*, *Phys. Rev.* **D26** (1982) 3327-3341.
2893. Tseytlin A.A., *On the first-order formalism in quantum gravity*, *J. Phys.* **A15** (1982) L105-L110.
2894. Tsoubelis D., *Bianchi  $VI_0, VII_0$  cosmological models with spin and torsion*, *Phys. Rev.* **D20** (1979) 3004-3008.
2895. Tsoubelis D., *Einstein-Cartan-Sciama-Kibble cosmological models with spinning matter and magnetic field*, *Phys. Rev.* **D23** (1981) 823-828.
2896. Tsoubelis D., *Gravitational field of a spin-polarized cylinder in the Einstein-Cartan theory of gravitation*, *Phys. Rev. Lett.* **51** (1983) 2235-2237.
2897. Tsoubelis D., *Static spin-polarized cylinder in the Einstein-Cartan theory of gravitation*, *Phys. Rev.* **D30** (1984) 1632-1636.
2898. Tsuda M., Shirafuji T., and Xie H., *General considerations of matter coupling with the self-dual connection*, *Class. Quantum Grav.* **12** (1995) 3067-3076.
2899. Tucker R.W., *A Clifford calculus for physical field theories*, in: "Clifford algebras and their applications in mathematical physics. Proc. of NATO and SERC workshop", Canterbury, UK, September 15-27, 1985, Eds. J.S.R. Chisholm, and A.K. Common (Reidel: Dordrecht, 1986) 177-199.
2900. Tucker R.W. and Wang C., *Black holes with Weyl charge and non-Riemannian waves*, *Class. Quantum Grav.* **12** (1995) 2587-2605.
2901. Tucker R.W. and Wang C., *An Einstein-Proca-fluid model for dark matter gravitational interactions*, *Nucl. Phys. B (Proc. Suppl.)* **57** (1997) 259-262.
2902. Tucker R.W. and Wang C., *Non-Riemannian gravitational interactions*, in: "Mathematics of gravitation", Part II (Warsaw, 1996), Banach Center Publications **41** (Polish Acad. Sci.: Warsaw, 1997) 263-271.
2903. Tucker R.W. and Wang C., *Dark matter gravitational interactions*, *Class. Quantum Grav.* **15** (1998) 933-954.
2904. Tung R.-S., *Gravitation as a supersymmetric gauge theory*, *Phys. Lett.* **A264** (2000) 341-345.
2905. Tung R.-S., Chang C.-H., Chern D.-C., and Nester J.M., *Asymptotic anti-de Sitter conditions for Poincaré gauge theory*, *Progr. Theor. Phys.* **88** (1992) 291-305.
2906. Tung R.-S. and Nester J.M., *The quadratic spinor Lagrangian is equivalent to the teleparallel theory*, *Phys. Rev.* **D60** (1999) 021501(R) (5 pages).
2907. Tunyak V.N., *Central-symmetric static field of a charged mass in an affine space with the Riemannian metric and torsion*, *Izvestiya AN BSSR, ser. fiz.-mat. nauk*, n. 2 (1972) 114-117 (in Russian).
2908. Tunyak V.N., *A theory of gravitation with variable gravitational constant in an affinely connected space with Riemannian metric and torsion*, *Izvestiya AN BSSR, ser. fiz.-mat. nauk*, n. 2 (1973) 84-87, 137 (in Russian).
2909. Tunyak V.N., *A tetrad theory of gravitation in a space with absolute parallelism*, *Izvestiya AN BSSR, ser. fiz.-mat. nauk*, n. 5 (1973) 98-104, 139 (in Russian).
2910. Tunyak V.N., *Axially symmetric static electrovacuum solutions in an affinely connected space with Riemannian metric and torsion*, *Izvestiya AN BSSR, ser. fiz.-mat. nauk*, n. 5 (1974) 97-101, 143 (in Russian).
2911. Tunyak V.N., *Variational principle for ideal fluid in a space with torsion*, *Doklady AN BSSR* **19** (1975) 599-601 (in Russian).

2912. Tuniyak V.N., *Determination of the Lagrangian in the tetrad theory of gravitation*, *Sov. J. Phys.* **18** (1975) 74-76 [*Izvestia VUZov, Fizika*, n. 1 (1975) 91-94 (in Russian)].
2913. Tuniyak V.N., *Determination of the source in the tetrad theory of gravitation*, *Sov. J. Phys.* **18** (1975) 77-79 [*Izvestia VUZov, Fizika*, n. 1 (1975) 94-97 (in Russian)].
2914. Tuniyak V.N., *On defining the spin moment in the tetrad theory of gravitation*, *Sov. J. Phys.* **19** (1976) 599-602 [*Izvestia VUZov, Fizika*, n. 5 (1976) 68-71 (in Russian)].
2915. Tuniyak V.N., *The de Sitter cosmological model in the tetrad theory of gravitation*, *Sov. J. Phys.* **19** (1976) 783-785 [*Izvestia VUZov, Fizika*, n. 6 (1976) 118-120 (in Russian)].
2916. Tuniyak V.N., *Homogeneous isotropic cosmological space-time model in the Einstein-Cartan gravitational theory*, *Sov. J. Phys.* **20** (1977) 297-300 [*Izvestia VUZov, Fizika*, n. 3 (1977) 24-28 (in Russian)].
2917. Tuniyak V.N., *Extended translational invariance and the tetrad theory of gravity*, *Sov. J. Phys.* **20** (1977) 538-541 [*Izvestia VUZov, Fizika*, n. 4 (1977) 137-141 (in Russian)].
2918. Tuniyak V.N., *Localization of the Poincaré group and the tetrad formulation of general relativity*, *Izvestia VUZov, Fizika*, n. 10 (1977) 100-103 (in Russian).
2919. Tuniyak V.N., *Quadratic Lagrangians in space with torsion*, *Sov. J. Phys.* **20** (1977) 1537-1539 [*Izvestia VUZov, Fizika*, n. 12 (1977) 7-10 (in Russian)].
2920. Tuniyak V.N., *Cosserat continuum in space with torsion*, *Sov. J. Phys.* **20** (1977) 1540-1542 [*Izvestia VUZov, Fizika*, n. 12 (1977) 11-13 (in Russian)].
2921. Tuniyak V.N., *Weak equivalence principle and gauge theory of the tetrad gravitational field*, *Sov. J. Phys.* **21** (1978) 426-429 [*Izvestia VUZov, Fizika*, n. 4 (1978) 18-21 (in Russian)].
2922. Tuniyak V.N., *Conservation laws in the tetrad formulation of the general theory of relativity*, *Sov. J. Phys.* **22** (1979) 194-197 [*Izvestia VUZov, Fizika*, n. 2 (1979) 101-105 (in Russian)].
2923. Tuniyak V.N., *Quadratic Lagrangians in space with torsion, and theory of gauge spinor field*, *Sov. J. Phys.* **22** (1979) 985-989 [*Izvestia VUZov, Fizika*, n. 9 (1979) 74-79 (in Russian)].
2924. Tuniyak V.N., *Cosmological term in the general theory of relativity and localization of the de Sitter and Einstein groups*, *Sov. J. Phys.* **27** (1984) 701-705 [*Izvestia VUZov, Fizika*, n. 8 (1983) 107-108 (in Russian)].
2925. Tuniyak V.N., *Theory of a gauge gravitational field with localization of the Einstein group*, *Sov. J. Phys.* **28** (1985) 579-583 [*Izvestia VUZov, Fizika*, n. 7 (1985) 68-73 (in Russian)].
2926. Tuniyak V.N., *Localization of the Einstein group and nonsingular cosmological model of space-time with torsion*, *Sov. J. Phys.* **29** (1986) 783-789 [*Izvestia VUZov, Fizika*, n. 10 (1986) 13-17 (in Russian)].
2927. Tuniyak V.N., *Theory of a gravitational gauge field with localization of the de Sitter group*, *Sov. J. Phys.* **29** (1986) 908-911 [*Izvestia VUZov, Fizika*, n. 11 (1986) 50-53 (in Russian)].
2928. Tuniyak V.N., *Relativistic theory of gravity and a torsion field*, *Theor. Math. Phys.* **118** (1999) 101-106.
2929. Tuniyak V.N. and Gorbatskii O.E., *Spherically symmetric static solution with a completely polarized Cosserat medium in the Einstein-Cartan theory*, *Sov. J. Phys.* **22** (1979) 822-825 [*Izvestia VUZov, Fizika*, n. 8 (1979) 21-24 (in Russian)].
2930. Tuniyak V.N. and Karakura F., *Dynamical torsion field*, *Izvestia AN BSSR, ser. fiz.-mat. nauk*, n. 2 (1981) 114-118 (in Russian).

## U

2931. Ulhoa S.C. and R.G.G. Amorim, *On teleparallel quantum gravity in Schwarzschild space-time*, *Adv. High En. Phys.* **2014** (2014) 812691 (6 pages).
2932. Ulhoa S.C., Khanna F.C., and Santana A.E., *Teleparallel formalism of Galilean gravity*, *Grav. Cosmol.* **17** (2011) 242-251.
2933. Ünal N., *Electromagnetic and gravitational interactions of the spinning particle*, *J. Math. Phys.* **47** (2006) 092501.
2934. Urani J.R. and Kutchko F.J., *Field equations and the tetrad connection*, *J. Math. Phys.* **26** (1985) 946-950.
2935. Urbakh V.Y., *Generalized theory of vector fields*, *Dokl. Akad. Nauk SSSR* **101** (1955) 1043-1046 (in Russian).
2936. Urban Z. and Volna J., *The metrizable problem for Lorentz-invariant affine connections*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1650110 (12 pages).
2937. Urrutia L.F. and Vergara J.D., *Consistent coupling of the gravitino field to a gravitational background with torsion*, *Phys. Rev.* **D44** (1991) 3882-3886.
2938. Urrutia L.F. and Zertuche F., *The Poincaré limit in 2+1 dimensional quantum de Sitter gravity*, *Class. Quant. Grav.* **9** (1992) 641-650.
2939. Utiyama R., *Invariant theoretical interpretation of interaction*, *Phys. Rev.* **101** (1956) 1597-1607.
2940. Utiyama R., *On Weyl's gauge field*, *Progr. Theor. Phys.* **50** (1973) 2080-2090.
2941. Utiyama R., *On Weyl's gauge field. II*, *Progr. Theor. Phys.* **53** (1975) 565-574.
2942. Utiyama R., *Introduction to the theory of general gauge fields*, *Progr. Theor. Phys.* **64** (1980) 2207-2221.
2943. Utiyama R. and Fukuyama T., *Gravitational field as a generalized gauge field*, *Progr. Theor. Phys.* **45** (1971) 612-627.

## V

2944. Vaitekinas A.S., Lavrov P.M., and Shvartsman Sh.M., *Relativistic spinning particle in a gravitational field with torsion*, *Russ. Phys. J.* **36** (1993) 1055-1058 [*Izvestia VUZov, Fizika* n. 11 (1993) 50-54 (in Russian)].
2945. Vakili B. and Jalalzadeh S., *Signature transition in Einstein-Cartan cosmology*, *Phys. Lett.* **B726** (2013) 28-32.
2946. Vanduyck M.A., *Maxwell's equations in spaces with non-metricity and torsion*, *J. Phys. A: Math. Gen.* **A29** (1996) 2245-2255.
2947. Vanduyck M.A., *A comparison between the Weyl method and the invariant formalism for spinorial covariant derivatives*, *Russ. J. Math. Phys.* **4** (1996) 131-136.
2948. Vanhecke F.J., *On a class of exact solutions of Poincaré gauge field theory in vacuum*, *Rev. bras. fis.* **13** (1984) 280-199.
2949. Vargas J.P., *Geometrization of the physics with teleparallelism. I. The classical interactions*, *Found. Phys.* **22** (1992) 507-526.
2950. Vargas J.P., *Affine torsion á la Cartan*, *Annales de la Fondation L. de Broglie* **32** (2007) 409-423.

2951. Vargas J.P. and Torr D.G., *Conservation of vector-valued forms and the question of the existence of gravitational energy-momentum in general relativity*, *Gen. Relat. Grav.* **23** (1991) 713-732.
2952. Vargas J.P. and Torr D.G., *The cornerstone role of the torsion in Finslerian physical worlds*, *Gen. Relat. Grav.* **27** (1995) 629-644.
2953. Vargas J.G. and Torr D.G., *The Cartan-Einstein unification with teleparallelism and the discrepant measurements of Newton's constant  $G$* , *Found. Phys.* **29** (1999) 145-200.
2954. Vargas J.P., Torr D.G., and Lecompte A., *Geometrization of the physics with teleparallelism. II: Towards a fully geometric Dirac equation*, *Found. Phys.* **22** (1992) 527-547.
2955. Vargas-Paredes A.A., Doria M.M., and Helayël-Neto J.A., *The Lichnerowicz-Weitzenböck formula and superconductivity*, *J. Math. Phys.* **54** (2013) 013101.
2956. Vasilic M., *Extra gauge symmetries in Poincaré gauge theory of gravity*, *Fizika (SFRJ)* **18** (1986) 140-144.
2957. Vasilic M., *String background fields and the Riemann-Cartan geometry*, *Class. Quantum Grav.* **28** (2011) 075008.
2958. Vasilic M. and Vojinovic M., *Zero-size objects in Riemann-Cartan spacetime*, *JHEP* **08** (2008) 104.
2959. Vasilic M. and Vojinovic M., *Spinning branes in Riemann-Cartan spacetimes*, *Phys. Rev.* **D78** (2008) 104002 (10 pages).
2960. Vasilic M. and Vojinovic M., *Test membranes in Riemann-Cartan spacetimes*, *Phys. Rev.* **D81** (2010) 024025 (7 pages).
2961. Vassilevich D.V., *On gauge-invariant boundary conditions for 2-D gravity with dynamical torsion*, *Mod. Phys. Lett.* **A10** (1995) 2239-2244.
2962. Vassilevich D.V., *Spectral geometry of symplectic spinors*, *J. Math. Phys.* **56** (2015) 103511 (10 pages).
2963. Vassiliev D., *A metric-affine field model for the neutrino*, in: "Noncommutative Structures in Mathematics and Physics" Eds. Duplij S. and Wess J., NATO Science Series, vol. **22** (Springer: Amsterdam, 2001) 427-439.
2964. Vassiliev D., *Pseudoinstantons in metric-affine field theory*, *Gen. Rel. Grav.* **34** (2002) 1239-1265.
2965. Vassiliev D., *Quadratic metric-affine gravity*, *Ann. d. Phys. (Berlin)* **14** (2005) 231-252.
2966. Vassiliev D., *Teleparallel model for the neutrino*, *Phys. Rev.* **D75** (2007) 025006 (6 pages).
2967. Vazirian R., Tanhayi M.R., and Motahar Z.A., *Weyl-invariant extension of the metric-affine gravity*, *Adv. High En. Phys.* **2015** (2015) 902396 (7 pages).
2968. Vercin A., *Metric-torsion gauge theory of continuum line defects*, *Int. J. Theor. Phys.* **29** (1990) 7-21.
2969. Vereshkov G.M., Dragilev V.M., and Ponomariev V.N., *Quantum gravitational effects in an isotropic universe with torsion*, *Sov. J. Phys.* **31** (1988) 691-694 [*Izvestia VUZov, Fizika* n9 (1988) 5-9 (in Russian)].
2970. Verwimp T., *Unified prescription for the generation of electroweak and gravitational gauge field Lagrangian on a principal fiber bundle*, *J. Math. Phys.* **31** (1990) 3047-3052.
2971. Viennot D., *Holonomy of a principal composite bundle connection, non-Abelian geometric phases, and gauge theory of gravity*, *J. Math. Phys.* **51** (2010) 103501.
2972. Viet N.A. and Wali K.C., *A discretized version of Kaluza-Klein theory with torsion and massive fields*, *Int. J. Mod. Phys.* **A11** (1996) 2403-2418.

2973. Vignolo S., Fabbri L., and Cianci R., *Dirac spinors in Bianchi-I  $f(R)$ -cosmology with torsion*, *J. Math. Phys.* **52** (2011) 112502.
2974. Vignolo S., Fabbri L., and Stornaiolo C., *A square-torsion modification of Einstein-Cartan theory*, *Ann. d. Phys. (Berlin)* **524** (2012) 826-839.
2975. Vignolo S., Carloni S., and Fabbri L., *Torsion gravity with nonminimally coupled fermionic field: Some cosmological models*, *Phys. Rev.* **D91** (2015) 043528 (12 pages).
2976. Vitagliano V., *The role of nonmetricity in metric-affine theories of gravity*, *Class. Quantum Grav.* **31** (2014) 045006.
2977. Vitagliano V., Sotiriou T.P., and Liberati S., *Dynamics of generalized Palatini theories of gravity*, *Phys. Rev.* **D82** (2010) 084007 (6 pages).
2978. Vitagliano V., Sotiriou T.P., and Liberati S., *The dynamics of metric-affine gravity*, *Ann. Phys. (USA)* **326** (2011) 1259-1273.
2979. Vitória R.L.L. and Bakke, K., *Torsion effects on a relativistic position-dependent mass system*, *Gen. Relativ. Gravit.* **48** (2016) 161 (11 pages).
2980. Vlachynsky E.J., Tresguerres R., Obukhov Yu.N., and Hehl F.W., *An axially symmetric solution of metric-affine gravity*, *Class. Quantum Grav.* **13** (1996) 3253-3259.
2981. Vladimirov A.A. and Diakonov D., *Phase transitions in spinor quantum gravity on a lattice*, *Phys. Rev.* **D86** (2012) 104019 (18 pages).
2982. Vladimirov Yu.S. and Popov A.D., *Composite torsion and spontaneous compactification of Kaluza-Klein type theories*, *Moscow University Physics Bulletin*, **29**, n. 4 (1988) 28-32 (in Russian).
2983. Voicu N., *Energy-momentum tensors in classical field theories: A modern perspective*, *Int. J. Geom. Methods Mod. Phys.* **13** (2016) 1640001 (20 pages).
2984. Vollick D.N., *Modified Palatini action that gives the Einstein-Maxwell theory*, *Phys. Rev.* **D93** (2016) 044061 (5 pages).
2985. Volovich I.V. and Katanaev M.O., *Scalar fields and dynamical torsion in the Kaluza-Klein type theories*, *Theor. Math. Phys.* **66** (1986) 53-60 [*Teor. Mat. Fiz.* **66** (1986) 79-89 (in Russian)].
2986. Volovich I.V. and Katanaev M.O., *Quantum strings with dynamical geometry*, *Pisma ZhETF* **43** (1986) 212-213 (in Russian).
2987. Volovik G.E. and Zubkov M.A., *Emergent geometry experienced by fermions in graphene in the presence of dislocations*, *Ann. Phys. (USA)* **356** (2015) 255-268.
2988. Vyblyi Y.P., *Nonsymmetrical tensor potential in the relativistic theory of gravity*, *Theor. Math. Phys.* **123** (2000) 851-857.

## W

2989. Waelbroeck H.,  *$B \wedge F$  theory and flat space-times*, *Commun. Math. Phys.* **170** (1995) 63-78.
2990. Wallner R.P., *Notes on recent  $U_4$  theories of gravitation*, *Gen. Relat. Grav.* **12** (1980) 719-732.
2991. Wallner R.P., *Notes on gauge theory and gravitation*, *Acta Phys. Austr.* **54** (1982) 165-189.
2992. Wallner R.P., *On hyperbolic  $U_4$ -manifolds with local duality*, *Acta Phys. Austr.* **55** (1983) 67-87.
2993. Wallner R.P., *On the structure of gravitational  $U_4$ -field equations*, *Gen. Relat. Grav.* **17** (1985) 1081-1107.

2994. Wallner R.P., *On new variables in gravity theories*, *Phys. Rev.* **D42** (1990) 441-448.
2995. Wallner R.P., *Exact solutions in  $U_4$  gravity. I. The ansatz for self double duality curvature*, *Gen. Relat. Grav.* **23** (1991) 623-639.
2996. Wanas M.I., *Motion of spinning particles in gravitational fields*, *Astrophys. Space Sci.* **258** (1997) 237-248.
2997. Wanas M.I., *Parameterized absolute parallelism: a geometry for physical applications*, *Turk. J. Phys.* **24** (2000) 473-488.
2998. Wanas M.I., *The accelerating expansion of the universe and torsion energy*, *Mod. Phys. Lett.* **A22** (2007) 5709.
2999. Wanas M.I. and Bakry M.A., *Effect of spin-torsion interaction on Raychaudhuri equation*, *Int. J. Mod. Phys.* **A24** (2009) 5025.
3000. Wanas M.I. and Hassan H.A., *Torsion and particle horizons*, *Int. J. Theor. Phys.* **53** (2014) 3901-3909.
3001. Wanas M.I. and Kahil M.E., *Quantum features of non-symmetric geometries*, *Gen. Relat. Grav.* **31** (1999) 1921-1929.
3002. Wanas M.I. and Kamal M.M., *An AP-structure with Finslerian flavor II: Torsion, curvature and other objects*, *Mod. Phys. Lett.* **A26** (2011) 2065.
3003. Wanas M.I. and Melek M., *Two notes about spin in absolute parallelism spaces*, *Astrophys. Space Sci.* **228** (1995) 277-281.
3004. Wanas M.I., Melek M., and Kahil M.E., *New path equations in absolute parallelism geometry*, *Astrophys. Space Sci.* **228** (1995) 273-276.
3005. Wanas M.I., Youssef N.L., and El Hanafy W., *A pure geometric theory of gravity and a material distribution*, *Gravit. Cosmol.* **23** (2017) 105-118.
3006. Wanas M.I., Youssef N.L., and Sid-Ahmed A.M., *Teleparallel Lagrange geometry and a unified field theory*, *Class. Quantum Grav.* **27** (2910) 045005.
3007. Wang C.-C., *On the geometric structures of simple bodies, a mathematical foundation for the theory of continuous distributions of dislocations*, *Archive Rat. Mech. Anal.* **27** (1967) 33-94.
3008. Wang C.-H., *Linearized torsion waves in a tensor-tensor theory of gravity*, *Annales de la Fondation L. de Broglie* **32** (2007) 267-280.
3009. Wang C.-H., *Charged fluid dynamics in scalar-tensor theories of gravity with torsion*, in: *Proc. of the 11th Marcel Grossmann Meeting "On recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories"*, Berlin, Germany, 23-29 July 2006, eds. H. Kleinert, R.T. Jantzen, and R. Ruffini (World Scientific: Singapore, 2008) pt. B, pp. 1278-1280.
3010. Wang C.-H. and Wu Y.-H., *Inflation in  $R + R^2$  gravity with torsion*, *Class. Quantum Grav.* **26** (2009) 045016.
3011. Wang J., Wang Y., and Yang C.L., *Dirac operators with torsion and the noncommutative residue for manifolds with boundary*, *J. Geom. Phys.* **81** (2014) 92-111.
3012. Wang T., *Static solutions with spherical symmetry in  $f(T)$  theories*, *Phys. Rev.* **D84** (2011) 024042 (10 pages).
3013. Wei H., *Dynamics of teleparallel dark energy*, *Phys. Lett.* **B712** (2012) 430-436.
3014. Wei H., Qi H.-Y., and Ma X.-P., *Constraining  $f(T)$  theories with the varying gravitational constant*, *Eur. Phys. J.* **C72** (2012) 2117.
3015. Weinberg S., *An alternative to general relativity*, in: "From  $SU(3)$  to gravity", *Festschrift in honor of Y. Ne'eman*, Eds. E. Gotsman and G. Tauber (Cambridge Univ. Press: Cambridge, 1985) 441-452.

3016. Weinberg S., *Einstein's mistakes*, *Physics Today* **58**, n. 11 (2005) 31-35.
3017. Weinberg S., *Weinberg replies*, *Physics Today* **59**, n. 4 (2006) 15-16.
3018. Weller J.M., *Fermion condensate from torsion in the reheating era after inflation*, *Phys. Rev.* **D88** (2013) 083511 (18 pages).
3019. Westman H.F. and Złošnik T.G., *Cartan gravity, matter fields, and the gauge principle*, *Ann. Phys. (USA)* **334** (2013) 157-197.
3020. Westman H.F. and Złošnik T.G., *Exploring Cartan gravity with dynamical symmetry breaking*, *Class. Quantum Grav.* **31** (2014) 095004.
3021. Westman H.F. and Złošnik T.G., *An introduction to the physics of Cartan gravity*, *Ann. Phys. (USA)* **361** (2015) 330-376.
3022. Wetterich C., *Dimensional reduction of fermions in generalized gravity*, *Nucl. Phys.* **B242** (1984) 473-502.
3023. Weyl H., *Elektron und Gravitation*, *Z. Phys.* **57** (1929) 261-272.
3024. Weyl H., *A remark on the coupling of gravitation and electron*, *Phys. Rev.* **77** (1950) 699-701.
3025. Weyssenhoff J., *Über die klassisch-relativistische Behandlung des Spinproblems*, in: *Max-Planck-Festschrift* (DVB: Berlin, 1958) 155-168.
3026. Weyssenhoff J. and Raabe A., *Relativistic dynamics of spin-fluids and spin-particle*, *Acta Phys. Pol.* **9** (1947) 7-18.
3027. Wheeler J.T., *Gravitational gauge theory and the existence of time*, *J. Phys.: Conf. Ser.* **462** (2013) 012059 (8 pages).
3028. Wiesendanger C., *Translational gauge invariance and classical gravitodynamics*, *Class. Quantum Grav.* **12** (1995) 585-603.
3029. Wiesendanger C., *Poincaré gauge invariance and gravitation in Minkowski spacetime*, *Class. Quantum Grav.* **13** (1996) 681-699.
3030. Wiesendanger C., *A Poincaré gauge theory of gravitation in Minkowski spacetime*, in: "Proc. of the 14th Course of Internat. School of Cosmology and Gravitation: Quantum Gravity" (Erice, 1995), Eds. P.G. Bergmann, V. de Sabbata, and H.-J. Treder (World Scientific: Singapore, 1996) 366-381.
3031. Wise D.K., *MacDowell-Mansouri gravity and Cartan geometry*, *Class. Quantum Grav.* **27** (2010) 155010.
3032. Wolf C., *Photon dispersion as a consequence of the interaction of torsion with an external magnetic field*, *Nuovo Cim.* **B91** (1986) 231-235.
3033. Wolf C., *A charged fluid sphere admitting spin generated torsion*, *Fizika (Croatia)* **B1** (1992) 197-206.
3034. Wolf C., *The effect of spin generated torsion on the early universe's evolution*, *Gen. Relat. Grav.* **27** (1995) 1031-1042.
3035. Wolf C., *Torsion and quadratic bulk viscosity in a homogeneous isotropic universe*, *Nuovo Cim.* **B112** (1997) 951-960.
3036. Wolf J.A., *On the geometry and classification of absolute parallelisms. I*, *J. Diff. Geom.* **6** (1972) 317-342.
3037. Write M., *Conformal transformations in modified teleparallel theories of gravity revisited*, *Phys. Rev.* **D93** (2016) 103002 (11 pages).
3038. Wu Y.-H. and Wang C.-H., *Momentum-space representation of the Feynman propagator in Riemann-Cartan spacetime*, *Phys. Rev.* **D82** (2010) 064007 (12 pages).

3039. Wu Y.-H. and Wang C.-H., *Brans-Dicke theory of gravity with torsion: A possible solution of  $\omega$ -problem*, *Phys. Rev.* **D86** (2012) 123519 (6 pages).
3040. Wu Y.L., *Quantum field theory of gravity with spin and scaling gauge invariance and spacetime dynamics with quantum inflation*, *Phys. Rev.* **D93** (2016) 024012 (35 pages).
3041. Wu Y.L. and Zhang G.M., *Spin interaction in teleparallel theory of gravitation*, *Nuovo Cim.* **B113** (1998) 1061-1065.
3042. Wu Y.-P., *Inflation with teleparallelism: Can torsion generate primordial fluctuations without local Lorentz symmetry?*, *Phys. Lett.* **B762** (2016) 157-161.
3043. Wu Y.-P. and Geng C.-Q., *Primordial fluctuations within teleparallelism*, *Phys. Rev.* **D86** (2012) 104058 (8 pages).
3044. Wu Y.-S. and Zee A., *Massless fermions and Kaluza-Klein theory with torsion*, *J. Math. Phys.* **25** (1984) 2696-2703.

## X

3045. Xie H.-J. and Shirafuji T., *Torsion field equation and spinor source*, *Prog. Theor. Phys.* **97** (1997) 129-140.
3046. Xu J.-Z., *Nonrelativistic approximation of scalar-tensor theory with torsion and intermediate-range force*, *Int. J. Theor. Phys.* **30** (1991) 1679-1689.
3047. Xu J.-Z., *Scalar-tensor theory with torsion and stellar structure*, *Int. J. Theor. Phys.* **37** (1998) 1151-1158.
3048. Xu J.-Z. and Chen Y.-H., *Intermediate range force and the scalar-tensor theory with torsion*, *Gen. Relat. Grav.* **23** (1991) 169-175.
3049. Xu J.-Z. and Chen Y.-H., *Scalar-tensor theory and propagating torsion theory*, *Int. J. Theor. Phys.* **30** (1991) 1263-1266.
3050. Xue S.-S., *The phase structure of Einstein-Cartan theory*, *Phys. Lett.* **B665** (2008) 54-57.
3051. Xue S.-S., *Quantum Regge calculus of Einstein-Cartan theory*, *Phys. Lett.* **B682** (2009) 300-304.
3052. Xue S.-S., *Detailed discussions and calculations of quantum Regge calculus of Einstein-Cartan theory*, *Phys. Rev.* **D82** (2010) 064039 (25 pages).

## Y

3053. Yadav R.B.S. and Prasad U., *Non-static conformally flat spherically symmetric perfect fluid distribution in Einstein-Cartan theory*, *Astropys. and Space Sci.* **203** (1993) 37-42.
3054. Yajima S., *Mixed anomalies in 4 and 6 dimensional space with torsion*, *Progr. Theor. Phys.* **79** (1988) 535-554.
3055. Yajima S., *Evaluation of the heat kernel in Riemann-Cartan space*, *Class. Quantum Grav.* **13** (1996) 2423-2435.
3056. Yajima S., *Evaluation of the heat kernel in Riemann-Cartan space using the covariant Taylor expansion method*, *Class. Quantum Grav.* **14** (1997) 2853-2868.
3057. Yajima S., Higasida Y., Kawano K., and Kubota S.I., *Torsion dependence of the covariant Taylor expansion in Riemann-Cartan space*, *Class. Quantum Grav.* **16** (1999) 1389-1399.
3058. Yajima S. and Kimura T., *Anomalies in four dimensional curved space-time with torsion*, *Progr. Theor. Phys.* **74** (1985) 866-880.

3059. Yajima S., Tokuo S., Fukuda M., Higashida Y., Kamo Y., Kubota S.-I., and Taira H., *Gravitational anomalies in higher dimensional Riemann-Cartan space*, *Class. Quantum Grav.* **24** (2007) 965.
3060. Yakushin I.V., *Torsion and interaction of polarized photons*, *Russ. Phys. J.* **34** (1991) 1090-1094 [*Izvestia VUZov, Fizika*, n. 12 (1991) 50-55 (in Russian)].
3061. Yakushin I.V., *Contribution of torsion to the hyperfine splitting of the hydrogen atom*, *Yad. Fiz.* **55** (1992) 753-757 (in Russian); English translation: *Sov. J. Nucl. Phys.* **55** (1992) 418-420.
3062. Yakushin I.V., *On the contribution of vector torsion component to the interaction effect of polarized photons*, *Vest. Mosk. Univ., Ser. 3 fiz. astr.* **33** (1992) 82-83 (in Russian).
3063. Yamaoka H. and Adachi T., *Continuum dynamics on a vector bundle for a directed medium*, *J. Phys. A: Math. Theor.* **43** (2010) 325209.
3064. Yamazaki K. and Nagahama H., *Hodge duality and continuum theory of defects*, *J. Phys. A: Math. and Gen.* **A32** (1999) L475-L481.
3065. Yan J. and Qiu X.M., *Sinh-Gordon matter field and a solvable model in two-dimensional gravity*, *Gen. Rel. Grav.* **30** (1998) 1319-1329.
3066. Yan M.L., *The renormalizability of general gravity theory with torsion and the spontaneous breaking of Lorentz group*, *Commun. Theor. Phys. (China)* **2** (1983) 1281-1298.
3067. Yang C.N., *Integral formalism for gauge fields*, *Phys. Rev. Lett.* **33** (1974) 445-447.
3068. Yang G.H., Duan Y.-S., and Huang Y.-C., *Topological invariant in Riemann-Cartan manifold and space-time defects*, *Int. J. Theor. Phys.* **37** (1998) 2953-2964.
3069. Yang J., Banerjee K., and Ma Y. *Hamiltonian analysis of  $R + T^2$  action*, *Phys. Rev.* **D85** (2012) 064047 (7 pages).
3070. Yang J., Banerjee K., and Ma Y. *Connection dynamics of a gauge theory of gravity coupled with matter*, *Class. Quantum Grav.* **30** (2013) 205015.
3071. Yang R.-J., *Conformal transformation in  $f(T)$  theories*, *Eur. Phys. Lett.* **93** (2011) 60001.
3072. Yasskin P., *Two kinds of rotation. An argument for torsion*, *Gen. Relat. Grav.* **13** (1981) 463-471.
3073. Yasskin P.B. and Stoeger W.R., *Propagating equations for test bodies with spin and rotation in theories of gravity with torsion*, *Phys. Rev.* **D21** (1980) 2081-2094.
3074. Yavari A., *A Geometric theory of growth mechanics*, *J. Nonlin. Sci.* **20** (2010) 781-830.
3075. Yavari A. and Goriely A., *Weyl geometry and the nonlinear mechanics of distributed point defects*, *Proc. Roy. Soc. London A: Math., Phys. and Eng. Sci.* **468** (2012) 1364-5021.
3076. Yavari A. and Goriely A., *Riemann-Cartan geometry of nonlinear dislocation mechanics*, *Archive for Rational Mechanics and Analysis* **205** (2012) 59-118.
3077. Yavari A. and Goriely A., *Riemann-Cartan geometry of nonlinear disclination mechanics*, *Mathematics and Mechanics of Solids* **18** (2013) 91-102.
3078. Yavari A. and Goriely A., *Non-metricity and the nonlinear mechanics of distributed point defects*, in: "Differential Geometry and Continuum Mechanics", Eds. G.-Q.G. Chen, M. Grinfeld, R.J. Knops, Springer Proceedings in Mathematics & Statistics **137** (2015) 235-251.
3079. Yavari A., and Marsden J.E., *Covariant balance laws in continua with microstructure*, *Rep. Math. Phys.* **63** (2009) 1-42.
3080. Yefremov A.P., *Some stationary cylindrically symmetric solutions of the Einstein-Cartan equations*, *Sov. J. Phys.* **20** (1977) 501-504 [*Izvestia VUZov, Fizika*, n. 4 (1977) 95-99 (in Russian)].

3081. Yefremov A.P., *Static homogeneous cosmological models in the Einstein-Cartan theory*, *Sov. J. Phys.* **20** (1977) 403-404 [*Izvestia VUZov, Fizika*, n. 3 (1977) 147-148 (in Russian)].
3082. Yefremov A.P., *Motion of test particles in a torsion field*, *Sov. J. Phys.* **23** (1980) 721-723 [*Izvestia VUZov, Fizika*, n. 8 (1980) 84-87 (in Russian)].
3083. Yefremov A.P., *Configuration of a neutron star with axially polarized spin*, *Acta Phys. Pol.* **B12** (1981) 185-188.
3084. Yefremov A.P., *Pulsar and spin-spin interaction*, *Exp. Tech. Physik* **29** (1981) 217-219.
3085. Yefremov A.P., *Gravitational field of a neutron star*, *Sov. J. Phys.* **25** (1982) 51-55 [*Izvestia VUZov, Fizika*, n. 1 (1982) 53-57 (in Russian)].
3086. Yo H.J., *The fine-structure constant and the torsion potential*, *Annales de la Fondation L. de Broglie* **32** (2007) 383-288.
3087. Yo H.J. and Nester J.M., *Hamiltonian analysis of Poincaré gauge theory scalar modes*, *Int. J. Mod. Phys.* **D8** (1999) 459-479.
3088. Yo H.J. and Nester J.M., *Hamiltonian analysis of Poincaré gauge theory: Higher spin modes*, *Int. J. Mod. Phys.* **D11** (2002) 747-779.
3089. Yo H.J. and Nester J.M., *Dynamic scalar torsion and an oscillating universe*, *Mod. Phys. Lett.* **A22** (2007) 2057-2069.
3090. Yoon Y.S., *Conformally coupled induced gravity with gradient torsion*, *Phys. Rev.* **D59** (1999) 127501.
3091. Yoshida H., *Geometric structure of the translation gauge theory of gravitation*, *Progr. Theor. Phys.* **67** (1982) 1107-1121.
3092. Yu X., *Mach's principle and torsion in general relativity*, in: "Spin in Gravity: Is it possible to give an experimental basis to torsion?", *Proc. of the 15th Course of Internat. School of Cosmology and Gravitation* (Erice, 1997), Eds. P.G. Bergmann, V. de Sabbata, G.T. Gillies and P.I. Pronin (World Scientific: Singapore, 1998) 239-252.

## Z

3093. Zakharov A.V. and Galiakhmetov A.M., *On equilibrium state nonsingular cosmological model in Einstein-Cartan theory*, *Ukr. Fiz. Zhurn.* **35** (1990) 327-335 (in Russian).
3094. Zakharov O.A. and Sardanashvily G.A., *On Higgs property of the gravitational field*, *Moscow University Physics Bulletin*, **30**, n. 5 (1989) 70-72 (in Russian).
3095. Zardecki A., *Gravity as a gauge theory with Cartan connection*, *J. Math. Phys.* **29** (1988) 1661-1666.
3096. Zardecki A., *A formulation of supergravity based on Cartan's connection*, *J. Math. Phys.* **34** (1993) 1487-1496.
3097. Zaripov F.Sh., Odintsov S.D., and Shapiro I.L., *Asymptotic behavior of the effective non-minimal coupling constants in curved space-time with torsion*, *Sov. Phys. J.* **33** (1990) 886-889 [*Izvestia VUZov, Fizika* n. 10 (1990) 87-91 (in Russian)].
3098. Zastawniak T., *Geometrical stochastic control and quantization*, *J. Math. Phys.* **38** (1997) 173-181.
3099. Zecca A., *Dirac equation in space-time with torsion*, *Int. J. Theor. Phys.* **41** (2002) 421-428.
3100. Zecca A., *Effect of torsion in Dirac equation for Coulomb potential in Robertson-Walker space-time*, *Int. J. Theor. Phys.* **41** (2002) 1145-1154.
3101. Zecca A., *Interacting gravitational and Dirac fields with torsion*, *Int. J. Theor. Phys.* **42** (2003) 2905-2911.

3102. Zecca A., *Field equations of arbitrary spin in space-time with torsion*, *Int. J. Theor. Phys.* **46** (2007) 1045-1054.
3103. Zhang C.M., *Some constraints on torsion detection physics*, *Int. J. Mod. Phys. A* **8** (1993) 5095-5100.
3104. Zhang C.M., *Backgrounds constrain on the torsion detection*, *Chinese Phys. Lett.* **10** ((1993) 645-647.
3105. Zhang C.M., *Rotation intrinsic spin effects and its astrophysical application*, *Mod. Phys. Lett. A* **13** (1998) 173-180.
3106. Zhang C.M., *Mass neutrino flavor evolution in spacetime with torsion*, *Nuovo Cim.* **B115** (2000) 437-444.
3107. Zhang C.M., *Neutrino oscillation in Weitzenböck space-time*, *Int. J. Mod. Phys. A* **15** (2000) 3453-3459.
3108. Zhang C.M., Chen F.P., and Garcia de Andrade L.C., *Torsion singularity in Weitzenböck space-time*, *Nuovo Cim.* **B110** (1995) 231-236.
3109. Zhang C.M., Yang G.C., Chen F.P., and Wu X.J., *Is there evidence for torsion?*, *Gen. Relat. Grav.* **24** (1992) 359-371.
3110. Zhang C.M. and Zhao P.D., *Equation of stellar structure in Hayashi's new general relativity*, *Int. J. Mod. Phys. A* **10** (1995) 2225-2230.
3111. Zhang G.M. and Wu Y.L., *Kaluza-Klein theory and Dirac equation in higher dimensional Riemann-Cartan space*, *Int. J. Theor. Phys.* **35** (1996) 1211-1221.
3112. Zhang G.M., Wu Y.L., and Liao T.H., *Neutrino solution of Dirac equation in Bianchi type V cosmological model*, *Int. J. Theor. Phys.* **36** (1997) 1881-1891.
3113. Zhang G.M., Wu Y.L., Wang X.-Z., and Sun J.-L., *Dirac Field of Kaluza-Klein theory in Weitzenböck space*, *Int. J. Theor. Phys.* **39** (2000) 2051-2062.
3114. Zhang Y., Li H., Gong Y., and Zhu Z.-H., *Notes on  $f(T)$  theories*, *JCAP* **07** (2011) 015.
3115. Zhang Y.Z., *Approximate solutions for general Riemann-Cartan type  $R + R^2$  theories of gravitation*, *Phys. Rev.* **D28** (1983) 1866-1871.
3116. Zhang Y.Z., *Newtonian limit in the Poincaré gauge field theory of gravitation*, *Scientia Sinica Ser. A: Math. Phys. Astron. and Tech. Sci.* **28** (1985) 399-404.
3117. Zhang Y.Z., *Conservation laws and Bianchi identities in  $R + R^2 + Q^2$  theories*, *Commun. Theor. Phys. (China)* **18** (1992) 337-342.
3118. Zhang Y.Z., *On the cosmological model in a spacetime theory with torsion*, *Acta Astrophysica Sinica* **13** (1993) 101.
3119. Zhang Y.Z., *Cosmological model in spacetime containing torsion*, *Chinese Astronomy and Astrophysics* **17** (1993) 257.
3120. Zhang Y.Z. and Guan K.Y., *Approximate solutions for  $R + R^2$  theories of gravitation: torsion components*, *Gen. Relat. Grav.* **16** (1984) 835-845.
3121. Zheng R., and Huang Q.-G., *Growth factor in  $f(T)$  gravity*, *JCAP* **03** (2011) 002.
3122. Zhogin I.L., *Topological quasi-charges in the Einstein theory of teleparallelism and particle combinatorics*, *Sov. Phys. J.* **33** (1990) 562-565 [*Izvestia VUZov, Fizika*, n. 7 (1990) 15-18 (in Russian)].
3123. Zhogin I.L., *Field equation for the Riemann space with torsion*, *Sov. Phys. J.* **33** (1990) 792-796 [*Izvestia VUZov, Fizika*, n. 9 (1990) 83-87 (in Russian)].
3124. Zhogin I.L., *Absolute parallelism: Spherical symmetry and singularities*, *Russ. Phys. J.* **34** (1991) 781-786 [*Izvestia VUZov, Fizika*, n. 9 (1991) 47-52 (in Russian)].

3125. Zhong Z.-Z., *Nonsymmetric gauge theory of gravitation*, *Phys. Rev.* **D26** (1982) 3412-3418.
3126. Zhytnikov V.V., *Conformally invariant Lagrangians in metric-affine and Riemann-Cartan spaces*, *Int. J. Mod. Phys.* **A8** (1993) 5141-5152.
3127. Zhytnikov V.V., *Wavelike exact solutions of  $R + R^2 + Q^2$  gravity*, *J. Math. Phys.* **35** (1994) 6001-6017.
3128. Zhytnikov V.V., *A new Kerr-Newman-De Sitter exact solution of the Poincaré gauge theory of gravitation*, *Chinese J. Phys.* **34** (1996) 1106-1112.
3129. Zhytnikov V.V., *Double duality and hidden gauge freedom in the Poincaré gauge theory of gravitation*, *Gen. Rel. Grav.* **28** (1996) 137-162.
3130. Zhytnikov V.V. and Ponomarev V.N., *On the equivalence of vacuum equations in quadratic gauge theory of gravity and in general relativity*, in: "Probl. Theor. Grav. and Elem. Particles", Ed. K.P. Stanyukovich (Energoatomizdat: Moscow, 1986) **17**, 93-101 (in Russian).
3131. Ziaie A.H., Moniz P.V., Ranjbar A., and Sepangi H.R., *Einstein-Cartan gravitational collapse of a homogeneous Weysenhoff fluid*, *Eur. Phys. J.* **C74** (2014) 3154 (19 pages).
3132. Zubkov M.A., *Teleparallel gravity on the lattice*, *Phys. Lett.* **B582** (2004) 243-248.
3133. Zubkov M.A., *Gauge invariant discretization of Poincaré quantum gravity*, *Phys. Lett.* **B638** (2006) 503-508.
3134. Zubkov M.A., *Torsion instead of technicolor*, *Mod. Phys. Lett.* **A25** (2010) 2885.
3135. Zubkov M.A., *Dynamical torsion as the microscopic origin of the neutrino seesaw*, *Mod. Phys. Lett.* **A29** (2014) 1450111 (16 pages).
3136. Zubkov M.A., *Emergent gravity and chiral anomaly in Dirac semimetals in the presence of dislocations*, *Ann. Phys. (USA)* **360** (2015) 655-678.







Научное издание

*Владимир Николаевич Пономарев,  
Андрей Олегович Барвинский,  
Юрий Николаевич Обухов*

# КАЛИБРОВОЧНЫЙ ПОДХОД И МЕТОДЫ КВАНТОВАНИЯ В ТЕОРИИ ГРАВИТАЦИИ

Издательство «Наука»  
117997, Москва, Профсоюзная ул., 90  
Главный редактор *Т. Е. Филиппова*  
Редактор *Л. В. Филиппова*

Формат 70 x 100  $\frac{1}{16}$ . Бумага офсетная 80 г/м<sup>2</sup>  
Печать офсетная. Гарнитура «СМР»  
Уч.-изд. л. 28,0. Усл.-печ. л. 29,03  
Тираж 1000 экз. Заказ № 734517

Заказное  
Отпечатано ФГУП Издательство «Наука»

Scientific edition

*Vladimir N. Ponomarev*

*Andrei O. Barvinsky*

*Yuri N. Obukhov*

GAUGE APPROACH  
AND QUANTIZATION METHODS  
IN GRAVITY THEORY

Moscow Nauka 2017

Contraria non contradietoria  
sed complementa sunt

Niels Bohr

May 8, 1961.

Physical law should have mathematical beauty

P A M Dirac

3 Oct 1951

Nature is simple  
in its essence.  
July 28, '59, Yulpawa

自然は本質において  
簡単である  
湯川  
(佐藤文隆)

ISBN 978-5-02-040047-4



9 785020 400474